



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 03:18 PM UTC

PDB ID : 4OFP / pdb_00004ofp
Title : Crystal Structure of SYG-2 D3-D4
Authors : Ozkan, E.; Garcia, K.C.
Deposited on : 2014-01-15
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	208	
1	B	208	
1	C	208	
1	D	208	

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 6032 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein SYG-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	197	1499	927	266	299	7	0	0	0
1	B	194	1472	909	262	294	7	0	0	0
1	C	193	1473	909	262	295	7	0	0	0
1	D	197	1504	928	269	300	7	0	0	0

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	229	ALA	-	expression tag	UNP Q9U3P2
A	230	ASP	-	expression tag	UNP Q9U3P2
A	431	HIS	-	expression tag	UNP Q9U3P2
A	432	HIS	-	expression tag	UNP Q9U3P2
A	433	HIS	-	expression tag	UNP Q9U3P2
A	434	HIS	-	expression tag	UNP Q9U3P2
A	435	HIS	-	expression tag	UNP Q9U3P2
A	436	HIS	-	expression tag	UNP Q9U3P2
B	229	ALA	-	expression tag	UNP Q9U3P2
B	230	ASP	-	expression tag	UNP Q9U3P2
B	431	HIS	-	expression tag	UNP Q9U3P2
B	432	HIS	-	expression tag	UNP Q9U3P2
B	433	HIS	-	expression tag	UNP Q9U3P2
B	434	HIS	-	expression tag	UNP Q9U3P2
B	435	HIS	-	expression tag	UNP Q9U3P2
B	436	HIS	-	expression tag	UNP Q9U3P2
C	229	ALA	-	expression tag	UNP Q9U3P2
C	230	ASP	-	expression tag	UNP Q9U3P2
C	431	HIS	-	expression tag	UNP Q9U3P2
C	432	HIS	-	expression tag	UNP Q9U3P2
C	433	HIS	-	expression tag	UNP Q9U3P2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	434	HIS	-	expression tag	UNP Q9U3P2
C	435	HIS	-	expression tag	UNP Q9U3P2
C	436	HIS	-	expression tag	UNP Q9U3P2
D	229	ALA	-	expression tag	UNP Q9U3P2
D	230	ASP	-	expression tag	UNP Q9U3P2
D	431	HIS	-	expression tag	UNP Q9U3P2
D	432	HIS	-	expression tag	UNP Q9U3P2
D	433	HIS	-	expression tag	UNP Q9U3P2
D	434	HIS	-	expression tag	UNP Q9U3P2
D	435	HIS	-	expression tag	UNP Q9U3P2
D	436	HIS	-	expression tag	UNP Q9U3P2

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	O	0	0
			14	8	1	5		
2	C	1	Total	C	N	O	0	0
			14	8	1	5		
2	D	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	D	1	14	8	1	5	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	121.24Å 121.24Å 178.16Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.31 – 3.00 46.31 – 3.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.31-3.00) 99.9 (46.31-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.16	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.99 (at 3.01Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8.4_1492)	Depositor
R, R_{free}	0.242 , 0.283 0.247 , 0.295	Depositor DCC
R_{free} test set	1365 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	106.1	Xtriage
Anisotropy	0.064	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 102.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6032	wwPDB-VP
Average B, all atoms (Å ²)	147.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/1530	0.89	2/2083 (0.1%)
1	B	0.41	0/1502	0.87	2/2046 (0.1%)
1	C	0.37	0/1502	0.82	2/2043 (0.1%)
1	D	0.42	0/1535	0.90	1/2089 (0.0%)
All	All	0.40	0/6069	0.87	7/8261 (0.1%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	260	GLY	N-CA-C	7.05	119.60	111.36
1	A	303	MET	N-CA-C	-6.02	105.59	113.17
1	B	260	GLY	N-CA-C	5.93	118.30	111.36
1	C	374	THR	CA-C-N	5.68	125.66	120.03
1	C	374	THR	C-N-CA	5.68	125.66	120.03

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1499	0	1437	26	0
1	B	1472	0	1410	37	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1473	0	1417	34	0
1	D	1504	0	1448	29	0
2	A	42	0	39	1	0
2	C	14	0	13	0	0
2	D	28	0	26	1	0
All	All	6032	0	5790	104	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 104 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:247:ARG:NH1	1:D:249:GLY:O	2.04	0.88
1:D:255:ALA:HB1	1:D:290:LYS:HE3	1.57	0.86
1:D:282:LEU:HG	1:D:289:ILE:HG12	1.61	0.82
1:B:282:LEU:HG	1:B:289:ILE:HA	1.61	0.82
1:A:247:ARG:NH2	1:D:248:SER:OG	2.13	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	193/208 (93%)	184 (95%)	9 (5%)	0	100	100
1	B	190/208 (91%)	176 (93%)	12 (6%)	2 (1%)	11	43
1	C	189/208 (91%)	178 (94%)	11 (6%)	0	100	100
1	D	193/208 (93%)	181 (94%)	11 (6%)	1 (0%)	24	60
All	All	765/832 (92%)	719 (94%)	43 (6%)	3 (0%)	30	65

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	285	ARG
1	D	243	PRO
1	B	242	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	170/185 (92%)	168 (99%)	2 (1%)	63	82
1	B	167/185 (90%)	167 (100%)	0	100	100
1	C	168/185 (91%)	167 (99%)	1 (1%)	78	88
1	D	172/185 (93%)	170 (99%)	2 (1%)	63	82
All	All	677/740 (92%)	672 (99%)	5 (1%)	76	86

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	285	ARG
1	A	321	LYS
1	C	254	ILE
1	D	284	THR
1	D	288	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	362	GLN
1	D	423	HIS
1	B	423	HIS
1	C	349	ASN
1	C	362	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	502	1	14,14,15	0.64	1 (7%)	17,19,21	1.23	1 (5%)
2	NAG	A	501	1	14,14,15	0.34	0	17,19,21	0.44	0
2	NAG	D	502	1	14,14,15	0.64	1 (7%)	17,19,21	0.68	0
2	NAG	A	503	1	14,14,15	0.35	0	17,19,21	0.41	0
2	NAG	D	501	1	14,14,15	0.27	0	17,19,21	0.49	0
2	NAG	C	501	1	14,14,15	0.34	0	17,19,21	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	502	1	-	2/6/23/26	0/1/1/1
2	NAG	A	501	1	-	0/6/23/26	0/1/1/1
2	NAG	D	502	1	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	503	1	-	1/6/23/26	0/1/1/1
2	NAG	D	501	1	-	2/6/23/26	0/1/1/1
2	NAG	C	501	1	-	2/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	502	NAG	C1-C2	2.25	1.55	1.52
2	D	502	NAG	C1-C2	2.07	1.55	1.52

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	502	NAG	C1-O5-C5	4.62	118.38	112.19

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	502	NAG	O5-C5-C6-O6
2	A	502	NAG	C4-C5-C6-O6
2	C	501	NAG	O5-C5-C6-O6
2	D	501	NAG	C8-C7-N2-C2
2	D	501	NAG	O7-C7-N2-C2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	502	NAG	1	0
2	D	501	NAG	1	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	197/208 (94%)	-0.12	1 (0%) 87 72	77, 120, 222, 280	0
1	B	194/208 (93%)	-0.15	1 (0%) 87 72	63, 122, 272, 334	0
1	C	193/208 (92%)	0.02	4 (2%) 63 40	94, 164, 274, 307	0
1	D	197/208 (94%)	-0.10	2 (1%) 79 59	69, 120, 221, 262	0
All	All	781/832 (93%)	-0.09	8 (1%) 79 59	63, 133, 255, 334	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	302	ASN	3.4
1	D	429	ALA	2.9
1	C	380	PHE	2.6
1	B	293	TYR	2.5
1	C	429	ALA	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	A	503	14/15	0.53	0.13	121,161,203,211	0
2	NAG	A	501	14/15	0.62	0.14	96,141,171,177	0
2	NAG	A	502	14/15	0.79	0.10	92,128,192,195	0
2	NAG	D	501	14/15	0.82	0.10	87,138,183,192	0
2	NAG	D	502	14/15	0.83	0.11	67,98,182,207	0
2	NAG	C	501	14/15	0.84	0.09	107,138,184,189	0

6.5 Other polymers [i](#)

There are no such residues in this entry.