



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 03:18 PM EDT

PDB ID : 6OFO / pdb_00006of0
Title : Crystal structure of split green fluorescent protein (GFP); s10 circular permutant (194-195)
Authors : Lin, C.-Y.; Romei, M.G.; Deller, M.C.; Doukov, T.I.; Boxer, S.G.
Deposited on : 2019-03-31
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	254	34% 77% 11% • 12%
1	B	254	46% 75% 13% 12%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3417 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Green fluorescent protein (GFP); s10 circular permutant (194-195).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	224	Total	C	N	O	S	0	1	2
			1714	1093	291	326	4			
1	B	223	Total	C	N	O	S	0	0	2
			1659	1069	282	304	4			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	23	Total	O	0	0
			23	23		
2	B	21	Total	O	0	0
			21	21		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Green fluorescent protein (GFP); s10 circular permutant (194-195)



- Molecule 1: Green fluorescent protein (GFP); s10 circular permutant (194-195)



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	50.75Å 51.07Å 97.30Å 90.00° 103.67° 90.00°	Depositor
Resolution (Å)	39.04 – 2.60 39.04 – 2.60	Depositor EDS
% Data completeness (in resolution range)	92.2 (39.04-2.60) 84.0 (39.04-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.08 (at 2.61Å)	Xtriage
Refinement program	PHENIX (1.13rc2_2986: ???)	Depositor
R, R_{free}	0.203 , 0.235 0.205 , 0.236	Depositor DCC
R_{free} test set	697 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	27.9	Xtriage
Anisotropy	0.343	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 56.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.016 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.83	EDS
Total number of atoms	3417	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GYS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.26	0/1732	0.42	2/2351 (0.1%)
1	B	0.25	0/1674	0.36	0/2275
All	All	0.25	0/3406	0.39	2/4626 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	130	VAL	N-CA-C	-5.79	94.78	111.00
1	A	130	VAL	CG1-CB-CG2	5.61	123.14	110.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1714	0	1602	21	0
1	B	1659	0	1524	18	0
2	A	23	0	0	4	0
2	B	21	0	0	3	0
All	All	3417	0	3126	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 6.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:81:ASP:OD1	1:B:92:ARG:NH1	2.19	0.76
1:B:43:VAL:HG23	1:B:134:SER:HA	1.69	0.73
1:A:34:ARG:NH1	2:A:303:HOH:O	2.24	0.69
1:A:158:ARG:HD3	1:A:245:GLN:HB2	1.74	0.69
1:B:15:THR:HG23	1:B:41:GLU:HG3	1.76	0.68
1:B:80:LEU:HD23	1:B:185:ILE:HB	1.76	0.67
1:A:246:GLN:NE2	2:A:301:HOH:O	2.21	0.67
1:B:110:SER:HB2	1:B:115:LEU:HD11	1.78	0.64
1:A:160:ILE:HB	1:A:168:TYR:HB2	1.80	0.63
1:A:124:THR:O	1:A:158:ARG:NH2	2.32	0.62
1:A:150:MET:HE3	1:A:181:LEU:HD11	1.80	0.62
1:B:46:ALA:HA	1:B:140:MET:HE1	1.82	0.60
1:A:136:TYR:O	1:A:141:LYS:NZ	2.35	0.59
1:A:182:VAL:HG11	1:A:184:ARG:HH11	1.68	0.58
1:B:34:ARG:NE	2:B:304:HOH:O	2.35	0.58
1:A:89:PHE:HA	1:A:112:THR:HG21	1.86	0.57
1:A:138:ASP:HA	1:A:141:LYS:HD2	1.87	0.56
1:A:130:VAL:CG1	1:A:181:LEU:HD23	2.38	0.54
1:B:147:LYS:NZ	2:B:306:HOH:O	2.41	0.53
1:B:160:ILE:HB	1:B:168:TYR:HB2	1.91	0.53
1:A:92:ARG:NE	2:A:308:HOH:O	2.41	0.53
1:A:157:GLU:OE1	1:A:171:ARG:NH2	2.41	0.50
1:B:223:ILE:HG13	1:B:247:ASN:HB2	1.94	0.50
1:A:41:GLU:HB3	1:A:104:LEU:HB2	1.94	0.49
1:B:70:PHE:CD2	1:B:147:LYS:HG2	2.49	0.47
1:B:155:VAL:HG22	1:B:173:VAL:HG22	1.96	0.47
1:A:182:VAL:HG11	1:A:184:ARG:NH1	2.31	0.46
1:B:95:GLY:HA3	1:B:106:LEU:HD23	1.97	0.46
1:A:128:GYS:N2	1:A:128:GYS:HD2	2.30	0.45
1:A:70:PHE:CD2	1:A:147:LYS:HG2	2.53	0.43
1:A:193:LYS:HB3	1:A:193:LYS:HE3	1.88	0.43
1:B:42:TYR:CE1	1:B:103:LYS:HG3	2.54	0.42
1:B:104:LEU:HD12	1:B:104:LEU:HA	1.88	0.42
1:A:130:VAL:HG11	1:A:181:LEU:HD23	2.02	0.41
1:A:177:GLU:O	2:A:302:HOH:O	2.22	0.41
1:B:78:VAL:HG22	1:B:183:ASN:HB3	2.03	0.41
1:B:124:THR:O	1:B:158:ARG:NH1	2.53	0.41
1:A:223:ILE:HG13	1:A:247:ASN:HB2	2.02	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:49:THR:HG23	2:B:301:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	216/254 (85%)	210 (97%)	6 (3%)	0	100	100
1	B	214/254 (84%)	207 (97%)	7 (3%)	0	100	100
All	All	430/508 (85%)	417 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/214 (81%)	174 (100%)	0	100	100
1	B	156/214 (73%)	156 (100%)	0	100	100
All	All	330/428 (77%)	330 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	8	ASN
1	A	183	ASN
1	A	201	HIS
1	B	201	HIS
1	B	226	ASN
1	B	231	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	GYS	B	128	1	20,22,23	1.75	5 (25%)	25,30,32	1.43	2 (8%)
1	GYS	A	128	1	20,22,23	1.61	3 (15%)	25,30,32	1.66	5 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	GYS	B	128	1	-	4/10/29/30	0/2/2/2
1	GYS	A	128	1	-	3/10/29/30	0/2/2/2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	128	GYS	C1-N3	3.76	1.43	1.37
1	A	128	GYS	O3-C3	3.66	1.40	1.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	128	GYS	O3-C3	3.55	1.40	1.20
1	A	128	GYS	CA2-N2	3.43	1.45	1.38
1	B	128	GYS	CA2-C2	3.07	1.51	1.48
1	A	128	GYS	C1-N3	2.96	1.42	1.37
1	B	128	GYS	CA2-N2	2.66	1.44	1.38
1	B	128	GYS	CG2-CB2	2.26	1.51	1.46

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	128	GYS	O3-C3-CA3	-4.59	104.81	125.47
1	B	128	GYS	O3-C3-CA3	-4.11	106.97	125.47
1	B	128	GYS	CA2-N2-C1	3.15	108.26	105.80
1	A	128	GYS	CA1-C1-N3	-2.88	121.09	124.84
1	A	128	GYS	CG2-CB2-CA2	-2.68	126.68	129.87
1	A	128	GYS	CA2-N2-C1	2.64	107.86	105.80
1	A	128	GYS	C2-N3-C1	2.19	109.08	108.07

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	128	GYS	N1-CA1-CB1-OG1
1	B	128	GYS	N1-CA1-CB1-OG1
1	B	128	GYS	C2-CA2-CB2-CG2
1	B	128	GYS	N2-CA2-CB2-CG2
1	A	128	GYS	C1-CA1-CB1-OG1
1	B	128	GYS	C1-CA1-CB1-OG1
1	A	128	GYS	N2-CA2-CB2-CG2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	128	GYS	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	223/254 (87%)	1.85	86 (38%) 1 0	19, 37, 64, 84	1 (0%)
1	B	222/254 (87%)	2.15	118 (53%) 0 0	24, 45, 71, 93	0
All	All	445/508 (87%)	2.00	204 (45%) 0 0	19, 41, 69, 93	1 (0%)

All (204) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	64	SER	7.9
1	A	50	HIS	6.2
1	B	138	ASP	5.1
1	A	130	VAL	5.1
1	B	33	LYS	5.0
1	A	65	GLN	4.9
1	B	4	MET	4.8
1	B	195	ASP	4.8
1	B	191	ASP	4.7
1	A	191	ASP	4.7
1	B	199	LEU	4.6
1	B	217	ASP	4.4
1	B	41	GLU	4.4
1	B	86	GLY	4.3
1	B	19	LYS	4.3
1	A	186	GLU	4.3
1	A	204	GLU	4.2
1	A	23	GLU	4.2
1	A	251	GLY	4.2
1	A	112	THR	4.0
1	A	41	GLU	4.0
1	B	155	VAL	4.0
1	B	219	GLN	4.0
1	B	251	GLY	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	49	THR	3.9
1	B	89	PHE	3.9
1	A	219	GLN	3.9
1	B	221	ASN	3.8
1	A	148	SER	3.8
1	B	23	GLU	3.7
1	A	195	ASP	3.7
1	B	49	THR	3.7
1	A	83	ASP	3.7
1	B	111	THR	3.6
1	B	66	GLY	3.6
1	B	192	PHE	3.6
1	B	131	GLN	3.6
1	B	98	ASP	3.6
1	B	218	LYS	3.6
1	B	12	SER	3.6
1	B	215	THR	3.5
1	A	81	ASP	3.5
1	A	194	GLU	3.5
1	B	65	GLN	3.5
1	B	18	SER	3.5
1	B	115	LEU	3.5
1	B	71	THR	3.5
1	B	100	THR	3.5
1	B	244	TYR	3.5
1	B	84	VAL	3.5
1	A	164	ASP	3.4
1	B	20	ASP	3.3
1	B	104	LEU	3.3
1	B	22	ASN	3.3
1	A	4	MET	3.3
1	B	87	HIS	3.3
1	A	193	LYS	3.2
1	A	179	ASP	3.2
1	A	209	SER	3.2
1	B	72	GLY	3.2
1	B	204	GLU	3.1
1	B	99	ALA	3.1
1	B	242	ASP	3.1
1	A	100	THR	3.1
1	A	178	GLY	3.0
1	B	117	VAL	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	157	GLU	3.0
1	A	161	SER	3.0
1	B	21	PRO	3.0
1	B	118	PRO	2.9
1	A	226	ASN	2.9
1	A	121	THR	2.9
1	B	113	GLY	2.9
1	B	153	GLY	2.9
1	A	111	THR	2.9
1	B	81	ASP	2.9
1	A	222	GLY	2.9
1	A	73	VAL	2.9
1	A	7	ASP	2.8
1	B	178	GLY	2.8
1	B	235	ASP	2.8
1	A	99	ALA	2.8
1	B	88	LYS	2.8
1	B	119	TRP	2.8
1	A	86	GLY	2.8
1	B	168	TYR	2.8
1	A	35	ASP	2.8
1	A	165	ASP	2.8
1	B	83	ASP	2.8
1	B	112	THR	2.8
1	B	184	ARG	2.8
1	B	148	SER	2.7
1	A	113	GLY	2.7
1	A	145	PHE	2.7
1	B	42	TYR	2.7
1	B	92	ARG	2.7
1	A	12	SER	2.7
1	B	196	GLY	2.7
1	A	170	THR	2.7
1	B	165	ASP	2.7
1	B	14	GLN	2.6
1	A	71	THR	2.6
1	A	190	THR	2.6
1	B	13	THR	2.6
1	A	36	HIS	2.6
1	A	142	ARG	2.6
1	B	189	GLY	2.6
1	B	197	ASN	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	64	SER	2.6
1	A	101[A]	ILE	2.6
1	B	181	LEU	2.6
1	A	118	PRO	2.6
1	B	159	THR	2.5
1	A	160	ILE	2.5
1	A	66	GLY	2.5
1	A	115	LEU	2.5
1	B	179	ASP	2.5
1	A	246	GLN	2.5
1	B	109	ILE	2.5
1	B	164	ASP	2.5
1	B	137	PRO	2.5
1	B	140	MET	2.5
1	B	91	VAL	2.5
1	A	188	LYS	2.4
1	A	104	LEU	2.4
1	B	77	LEU	2.4
1	B	144	ASP	2.4
1	A	177	GLU	2.4
1	B	198	ILE	2.4
1	B	201	HIS	2.4
1	A	5	LEU	2.4
1	A	22	ASN	2.4
1	B	226	ASN	2.4
1	A	217	ASP	2.4
1	B	134	SER	2.4
1	B	152	GLU	2.4
1	A	119	TRP	2.4
1	A	141	LYS	2.4
1	A	151	PRO	2.4
1	A	223	ILE	2.3
1	B	101	ILE	2.3
1	A	132	ALA	2.3
1	B	250	ILE	2.3
1	B	208	ASN	2.3
1	B	110	SER	2.3
1	B	17	LEU	2.3
1	B	146	PHE	2.3
1	A	163	LYS	2.3
1	B	34	ARG	2.3
1	A	162	PHE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	144	ASP	2.3
1	B	116	PRO	2.3
1	B	82	GLY	2.3
1	A	221	ASN	2.2
1	A	198	ILE	2.2
1	B	185	ILE	2.2
1	B	150	MET	2.2
1	B	15	THR	2.2
1	A	249	PRO	2.2
1	A	98	ASP	2.2
1	B	90	SER	2.2
1	B	74	VAL	2.2
1	B	70	PHE	2.2
1	B	183	ASN	2.2
1	B	38	VAL	2.2
1	A	87	HIS	2.2
1	A	180	THR	2.2
1	A	92	ARG	2.2
1	A	171	ARG	2.2
1	A	197	ASN	2.2
1	B	209	SER	2.2
1	A	241	ALA	2.1
1	B	149	ALA	2.1
1	B	177	GLU	2.1
1	A	244	TYR	2.1
1	A	37	MET	2.1
1	B	73	VAL	2.1
1	B	171	ARG	2.1
1	A	200	GLY	2.1
1	A	231	HIS	2.1
1	B	7	ASP	2.1
1	B	133	PHE	2.1
1	A	20	ASP	2.1
1	A	138	ASP	2.1
1	A	15	THR	2.1
1	A	116	PRO	2.1
1	B	126	LEU	2.1
1	A	213	TYR	2.1
1	A	233	VAL	2.1
1	B	16	VAL	2.1
1	B	212	VAL	2.1
1	B	105	THR	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	48	ILE	2.0
1	B	43	VAL	2.0
1	B	194	GLU	2.0
1	B	120	PRO	2.0
1	B	5	LEU	2.0
1	B	122	LEU	2.0
1	B	187	LEU	2.0
1	B	246	GLN	2.0
1	A	109	ILE	2.0
1	B	44	ASN	2.0
1	B	78	VAL	2.0
1	A	154	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	GYS	A	128	21/22	0.85	0.16	23,28,37,43	0
1	GYS	B	128	21/22	0.86	0.14	31,37,45,49	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.