



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 08:10 AM UTC

PDB ID : 6OFS / pdb_00006ofs
Title : The crystal structure of the periplasmic protease PqqL from Escherichia coli
Authors : Grinter, R.
Deposited on : 2019-04-01
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	913	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7238 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Probable zinc protease PqqL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	905	Total	C	N	O	S	0	0	0
			7164	4479	1271	1394	20			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	932	LEU	-	expression tag	UNP P31828
A	933	GLU	-	expression tag	UNP P31828
A	934	HIS	-	expression tag	UNP P31828
A	935	HIS	-	expression tag	UNP P31828
A	936	HIS	-	expression tag	UNP P31828
A	937	HIS	-	expression tag	UNP P31828
A	938	HIS	-	expression tag	UNP P31828
A	939	HIS	-	expression tag	UNP P31828

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Cl	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	71	Total	O	0	0
			71	71		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Probable zinc protease PqqL



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	98.67Å 98.67Å 230.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.13 – 2.60 44.13 – 2.60	Depositor EDS
% Data completeness (in resolution range)	100.0 (44.13-2.60) 99.9 (44.13-2.60)	Depositor EDS
R_{merge}	0.35	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.93 (at 2.61Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.199 , 0.245 0.211 , 0.256	Depositor DCC
R_{free} test set	1737 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	56.6	Xtriage
Anisotropy	0.104	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 26.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7238	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/7289	1.28	19/9891 (0.2%)

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	554	ASP	CA-CB-CG	7.52	120.12	112.60
1	A	755	ASN	N-CA-C	-6.85	102.22	110.44
1	A	930	PRO	CA-C-N	5.73	132.02	121.70
1	A	930	PRO	C-N-CA	5.73	132.02	121.70
1	A	275	ASP	CA-CB-CG	5.58	118.17	112.60
1	A	207	VAL	N-CA-CB	5.45	117.43	110.13
1	A	191	ARG	N-CA-C	-5.41	105.55	111.82
1	A	217	GLN	CA-C-N	5.39	127.50	120.28
1	A	217	GLN	C-N-CA	5.39	127.50	120.28
1	A	282	GLY	N-CA-C	5.36	118.58	110.75
1	A	231	GLY	N-CA-C	5.34	117.96	110.38
1	A	798	GLU	CB-CG-CD	5.28	121.57	112.60
1	A	130	THR	CA-C-N	5.27	127.29	120.44
1	A	130	THR	C-N-CA	5.27	127.29	120.44
1	A	688	ASP	CA-CB-CG	5.21	117.81	112.60
1	A	300	PHE	CA-C-N	5.07	126.95	120.56
1	A	300	PHE	C-N-CA	5.07	126.95	120.56
1	A	32	ASP	CA-CB-CG	5.07	117.67	112.60
1	A	791	ASP	CA-CB-CG	5.01	117.61	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7164	0	7116	33	0
2	A	1	0	0	0	0
3	A	2	0	0	0	0
4	A	71	0	0	0	0
All	All	7238	0	7116	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:788:LEU:HD11	1:A:843:MET:HG2	1.85	0.59
1:A:775:LEU:HG	1:A:779:MET:HE2	1.87	0.56
1:A:765:LYS:HG2	1:A:923:THR:HG22	1.87	0.56
1:A:336:SER:HA	1:A:345:SER:O	2.09	0.52
1:A:291:MET:HE1	1:A:338:LYS:HG3	1.93	0.51
1:A:791:ASP:OD2	1:A:846:ARG:HD2	2.11	0.50
1:A:927:VAL:HG12	1:A:929:LEU:HG	1.95	0.48
1:A:708:VAL:HG12	1:A:874:VAL:HG22	1.95	0.48
1:A:534:LEU:HD23	1:A:707:ILE:HB	1.96	0.47
1:A:796:ILE:HG23	1:A:835:LEU:HD22	1.96	0.47
1:A:53:HIS:HB3	1:A:54:PRO:HD3	1.95	0.46
1:A:562:LYS:HD3	1:A:672:ARG:HB3	1.96	0.46
1:A:596:MET:HE3	1:A:622:LEU:HG	1.97	0.46
1:A:859:TYR:CD1	1:A:904:MET:HE1	2.51	0.46
1:A:596:MET:HE2	1:A:607:LEU:HD11	1.97	0.46
1:A:69:LEU:HG	1:A:119:TYR:HB3	1.98	0.46
1:A:403:ASP:HB3	1:A:406:MET:HE2	1.98	0.45
1:A:838:LEU:HA	1:A:841:GLU:HG2	1.97	0.45
1:A:577:VAL:HG13	1:A:625:GLN:HB3	1.98	0.45
1:A:311:GLN:HE22	1:A:388:ARG:HD3	1.82	0.44
1:A:130:THR:HG21	1:A:232:ASP:O	2.17	0.44
1:A:258:ARG:HB2	1:A:260:TRP:CH2	2.53	0.44
1:A:221:GLN:HB2	1:A:254:ALA:HB2	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:700:PRO:HA	1:A:703:ILE:HD12	1.99	0.43
1:A:806:VAL:HG23	1:A:825:PHE:HB3	2.01	0.42
1:A:548:ALA:HB3	1:A:607:LEU:HB3	2.02	0.42
1:A:64:ILE:HD13	1:A:225:MET:HG2	2.02	0.41
1:A:328:THR:HB	1:A:359:ASP:HB3	2.01	0.41
1:A:663:MET:HE1	1:A:749:VAL:HB	2.03	0.41
1:A:774:ASN:CG	1:A:776:PRO:HD2	2.46	0.41
1:A:182:ARG:HD3	1:A:459:GLU:OE1	2.20	0.40
1:A:823:LEU:HD22	1:A:843:MET:HE2	2.03	0.40
1:A:503:VAL:HG22	1:A:625:GLN:HE21	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	903/913 (99%)	879 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	781/789 (99%)	752 (96%)	29 (4%)	30	57

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	83	GLU
1	A	111	ARG
1	A	132	GLN
1	A	138	GLN
1	A	144	SER
1	A	214	GLN
1	A	297	GLU
1	A	322	GLN
1	A	329	ILE
1	A	336	SER
1	A	343	TYR
1	A	346	LEU
1	A	365	MET
1	A	542	GLN
1	A	563	SER
1	A	577	VAL
1	A	731	ASP
1	A	797	ARG
1	A	799	GLN
1	A	820	SER
1	A	851	ILE
1	A	858	GLU
1	A	867	LEU
1	A	870	GLN
1	A	874	VAL
1	A	875	GLN
1	A	876	GLN
1	A	909	VAL
1	A	918	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	51	HIS
1	A	88	ASN
1	A	114	ASN
1	A	180	GLN
1	A	245	ASN
1	A	257	ASN
1	A	278	ASN
1	A	311	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	358	GLN
1	A	417	ASN
1	A	464	ASN
1	A	542	GLN
1	A	592	ASN
1	A	615	ASN
1	A	625	GLN
1	A	635	ASN
1	A	695	GLN
1	A	754	GLN
1	A	763	GLN
1	A	799	GLN
1	A	879	ASN
1	A	887	GLN
1	A	915	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	905/913 (99%)	-0.12	8 (0%) 81 78	38, 61, 91, 126	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	869	ILE	2.9
1	A	233	ILE	2.8
1	A	27	ALA	2.8
1	A	28	ALA	2.6
1	A	870	GLN	2.4
1	A	810	LEU	2.2
1	A	238	ALA	2.1
1	A	559	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	A	1001	1/1	0.96	0.14	65,65,65,65	0
3	CL	A	1002	1/1	0.96	0.06	73,73,73,73	0
3	CL	A	1003	1/1	0.96	0.14	79,79,79,79	0

6.5 Other polymers [i](#)

There are no such residues in this entry.