



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:05 PM UTC

PDB ID : 4OHT / pdb_00004oht
Title : Crystal structure of succinic semialdehyde dehydrogenase from *Streptococcus pyogenes* in complex with NADP⁺ as the cofactor
Authors : Park, S.A.; Jang, E.H.; Chi, Y.M.; Lee, K.S.
Deposited on : 2014-01-18
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.


Mol	Chain	Length	Quality of chain
1	A	465	4% 83% 14% .
1	B	465	4% 82% 16% .

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Succinate-semialdehyde dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	455	Total 3574	C 2279	N 585	O 697	S 13	0	0	0
1	B	455	Total 3574	C 2279	N 585	O 697	S 13	0	0	0

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 31	C 10	N 5	O 13	P 3	0	0
2	B	1	Total 31	C 10	N 5	O 13	P 3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	324	Total 324	O 324	0	0
3	B	290	Total 290	O 290	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Succinate-semialdehyde dehydrogenase



- Molecule 1: Succinate-semialdehyde dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	134.76Å 134.76Å 173.53Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.89 – 2.10 39.89 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.5 (39.89-2.10) 99.5 (39.89-2.10)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.91 (at 2.10Å)	Xtriage
Refinement program	CNS 1.3	Depositor
R, R_{free}	0.201 , 0.224 0.201 , 0.224	Depositor DCC
R_{free} test set	4665 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	30.9	Xtriage
Anisotropy	0.008	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7824	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.66	8/3653 (0.2%)	0.87	7/4951 (0.1%)
1	B	0.76	15/3653 (0.4%)	0.86	6/4951 (0.1%)
All	All	0.71	23/7306 (0.3%)	0.87	13/9902 (0.1%)

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	438	LEU	C-O	-17.76	1.01	1.24
1	B	438	LEU	C-O	-16.27	1.03	1.24
1	B	441	LEU	C-O	-15.17	1.06	1.24
1	A	437	GLU	C-O	-14.88	1.05	1.23
1	B	440	GLU	C-O	-14.04	1.07	1.24
1	B	437	GLU	C-O	-13.02	1.07	1.23
1	A	439	SER	C-O	-12.43	1.08	1.24
1	B	439	SER	C-O	-11.55	1.09	1.24
1	B	437	GLU	CD-OE2	-7.48	1.11	1.25
1	B	439	SER	CB-OG	-7.48	1.27	1.42
1	A	437	GLU	CD-OE1	-7.24	1.11	1.25
1	B	437	GLU	CA-C	-6.77	1.44	1.52
1	A	439	SER	CA-C	-6.53	1.44	1.52
1	B	440	GLU	CD-OE2	-6.52	1.12	1.25
1	B	440	GLU	CA-C	-6.45	1.44	1.52
1	B	438	LEU	N-CA	-6.45	1.37	1.46
1	B	441	LEU	C-N	-6.06	1.26	1.33
1	A	438	LEU	CA-CB	-6.05	1.43	1.53
1	A	437	GLU	CD-OE2	-5.97	1.14	1.25
1	B	437	GLU	CD-OE1	-5.42	1.15	1.25
1	A	437	GLU	N-CA	-5.40	1.39	1.45
1	B	439	SER	N-CA	-5.32	1.39	1.46
1	B	438	LEU	CA-CB	-5.25	1.44	1.53

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	75	GLY	N-CA-C	7.00	124.67	115.36
1	B	75	GLY	N-CA-C	6.69	124.93	115.30
1	A	437	GLU	CB-CG-CD	6.46	123.58	112.60
1	B	201	VAL	N-CA-C	5.74	117.62	112.17
1	B	154	LEU	N-CA-C	5.62	118.06	108.90
1	B	251	LEU	N-CA-C	5.54	117.76	111.11
1	B	234	ASP	N-CA-C	5.51	118.00	110.24
1	A	234	ASP	N-CA-C	5.47	118.28	110.24
1	A	251	LEU	N-CA-C	5.43	117.62	111.11
1	A	237	ILE	N-CA-C	5.37	115.83	107.99
1	A	267	LYS	N-CA-C	5.33	118.00	111.82
1	A	132	ASN	N-CA-C	5.21	117.71	111.71
1	B	437	GLU	CB-CG-CD	5.09	121.26	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3574	0	3480	35	1
1	B	3574	0	3480	41	0
2	A	31	0	11	1	0
2	B	31	0	11	1	0
3	A	324	0	0	2	0
3	B	290	0	0	5	0
All	All	7824	0	6982	75	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:139:MET:HE2	1:B:143:ALA:HB2	1.38	1.05
1:A:321:ALA:HB2	1:A:356:ILE:HG13	1.72	0.72
1:A:136:TYR:CZ	1:A:140:ARG:HD3	2.25	0.72
1:B:153:VAL:HB	1:B:184:LEU:HD11	1.77	0.65
1:B:136:TYR:CZ	1:B:140:ARG:HD3	2.31	0.65
1:B:139:MET:HE1	1:B:142:PHE:HD2	1.62	0.65
1:A:210:ARG:HB2	1:A:210:ARG:NH1	2.12	0.64
1:A:214:SER:O	1:A:218:GLU:HG3	1.98	0.63
1:B:305:LEU:HD12	1:B:311:LYS:HA	1.79	0.63
1:A:248:GLU:HG3	1:A:282:LEU:HD11	1.81	0.62
1:A:238:ILE:HD12	1:A:271:VAL:HG12	1.80	0.62
1:B:139:MET:HE3	1:B:139:MET:HA	1.82	0.61
1:B:139:MET:HE2	1:B:143:ALA:CB	2.24	0.60
1:B:321:ALA:HB2	1:B:356:ILE:HG13	1.83	0.60
1:A:136:TYR:OH	1:A:140:ARG:HD3	2.03	0.58
1:B:205:LEU:C	1:B:205:LEU:HD23	2.28	0.57
1:B:271:VAL:HG23	3:B:606:HOH:O	2.03	0.57
1:B:47:GLU:HG3	3:B:728:HOH:O	2.05	0.56
1:A:305:LEU:HD12	1:A:311:LYS:HA	1.90	0.54
1:A:375:GLU:HG2	3:A:653:HOH:O	2.08	0.54
1:B:109:GLU:HG3	3:B:819:HOH:O	2.07	0.53
1:A:274:LYS:HE2	3:A:820:HOH:O	2.09	0.53
1:A:239:LEU:HD12	1:A:394:SER:HB2	1.90	0.51
1:B:138:ILE:HD11	1:B:156:HIS:CD2	2.45	0.50
1:A:152:MET:O	1:A:181:ILE:HA	2.12	0.50
1:A:171:VAL:HG12	1:A:176:ALA:HB3	1.94	0.49
1:B:277:ASP:O	1:B:281:GLU:HG3	2.11	0.49
1:A:138:ILE:HD11	1:A:156:HIS:CD2	2.48	0.49
1:A:144:PRO:O	1:A:148:VAL:HG13	2.13	0.49
1:B:211:GLY:HA3	3:B:837:HOH:O	2.13	0.49
1:A:205:LEU:C	1:A:205:LEU:HD23	2.38	0.49
1:B:131:TRP:CG	1:B:159:ILE:HD11	2.47	0.49
1:A:133:PHE:HB3	1:A:136:TYR:HB3	1.94	0.48
1:A:276:TYR:CG	1:A:370:LYS:HD3	2.49	0.48
1:B:8:PRO:HG2	1:B:74:MET:C	2.39	0.48
1:B:109:GLU:HG3	1:B:109:GLU:O	2.14	0.47
1:B:215:ILE:HD11	2:B:501:NAP:C2A	2.45	0.47
1:A:267:LYS:HE2	1:A:360:GLU:HG3	1.95	0.47
1:B:65:LYS:HG3	1:B:66:TYR:N	2.30	0.47
1:B:152:MET:O	1:B:181:ILE:HA	2.15	0.47
1:B:303:ALA:HB1	1:B:304:PRO:HD2	1.97	0.46
1:A:148:VAL:HG12	1:A:443:PHE:CZ	2.50	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:171:VAL:HG12	1:B:176:ALA:HB3	1.98	0.46
1:A:8:PRO:HG2	1:A:74:MET:C	2.41	0.45
1:B:186:ILE:HB	1:B:190:GLN:HB2	1.99	0.45
1:A:186:ILE:HB	1:A:190:GLN:HB2	1.98	0.45
1:A:427:GLY:HA3	1:A:436:ARG:HD3	2.00	0.44
1:A:276:TYR:O	1:A:280:LYS:HG3	2.17	0.43
1:B:139:MET:CE	1:B:142:PHE:HD2	2.29	0.43
1:A:145:ASN:O	1:A:148:VAL:HG22	2.18	0.43
1:B:262:VAL:HB	1:B:265:SER:HB2	2.00	0.43
1:B:160:CYS:N	1:B:161:PRO:CD	2.81	0.43
1:B:58:ILE:HD13	1:B:173:GLU:HG2	2.01	0.43
1:B:65:LYS:O	1:B:69:ILE:HG13	2.20	0.42
1:B:394:SER:HB3	1:B:400:ALA:HB2	2.01	0.42
1:A:153:VAL:HB	1:A:184:LEU:HD11	2.01	0.42
1:B:287:PHE:CD2	1:B:344:PRO:HB3	2.55	0.42
1:A:165:GLN:O	1:A:169:GLU:HG3	2.20	0.42
1:B:262:VAL:HG12	1:B:264:THR:H	1.85	0.42
1:A:215:ILE:HD11	2:A:501:NAP:C2A	2.50	0.41
1:B:117:TYR:CE1	1:B:119:LYS:HE2	2.55	0.41
1:A:454:ILE:HG12	1:B:414:ILE:HB	2.03	0.41
1:B:337:HIS:CG	1:B:338:PRO:HD2	2.55	0.41
1:B:40:ARG:NH2	3:B:679:HOH:O	2.54	0.41
1:A:262:VAL:HG12	1:A:264:THR:H	1.85	0.41
1:A:192:SER:OG	1:A:218:GLU:OE2	2.39	0.41
1:A:210:ARG:HB2	1:A:210:ARG:HH11	1.84	0.41
1:A:371:VAL:HB	1:A:376:GLU:HB3	2.03	0.41
1:B:101:GLU:HA	1:B:104:MET:HE2	2.02	0.41
1:B:375:GLU:O	1:B:379:GLU:HG3	2.21	0.40
1:B:62:ASP:HB3	1:B:65:LYS:HG2	2.03	0.40
1:A:337:HIS:ND1	1:A:338:PRO:HD2	2.37	0.40
1:B:136:TYR:OH	1:B:140:ARG:HD3	2.21	0.40
1:A:160:CYS:N	1:A:161:PRO:CD	2.84	0.40
1:B:138:ILE:HD11	1:B:156:HIS:HD2	1.85	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:43:ASP:OD2	1:A:43:ASP:OD2[8_554]	2.13	0.07

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	453/465 (97%)	441 (97%)	12 (3%)	0	100	100
1	B	453/465 (97%)	440 (97%)	13 (3%)	0	100	100
All	All	906/930 (97%)	881 (97%)	25 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	380/390 (97%)	377 (99%)	3 (1%)	73	81
1	B	380/390 (97%)	373 (98%)	7 (2%)	51	60
All	All	760/780 (97%)	750 (99%)	10 (1%)	61	69

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	281	GLU
1	A	362	PHE
1	A	393	PHE
1	B	155	LYS
1	B	198	LYS
1	B	245	ASP
1	B	322	LEU
1	B	362	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	393	PHE
1	B	419	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	193	GLN
1	A	309	GLN
1	A	317	GLN
1	A	359	GLN
1	A	397	GLN
1	B	4	GLN
1	B	114	GLN
1	B	246	GLN
1	B	317	GLN
1	B	340	HIS
1	B	359	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAP	A	501	-	32,33,52	1.14	2 (6%)	50,52,80	1.73	9 (18%)
2	NAP	B	501	-	32,33,52	1.17	2 (6%)	50,52,80	1.73	8 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	A	501	-	-	1/21/37/67	0/3/3/5
2	NAP	B	501	-	-	0/21/37/67	0/3/3/5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501	NAP	C8A-N7A	2.65	1.36	1.31
2	A	501	NAP	C8A-N7A	2.55	1.36	1.31
2	B	501	NAP	P2B-O2X	-2.09	1.47	1.54
2	A	501	NAP	C5A-C4A	2.08	1.42	1.39

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	NAP	C5A-C4A-N3A	-5.65	118.94	126.72
2	A	501	NAP	C5A-C4A-N3A	-5.56	119.06	126.72
2	B	501	NAP	N3A-C4A-N9A	4.13	134.19	127.17
2	B	501	NAP	N3A-C2A-N1A	-4.03	122.48	128.58
2	A	501	NAP	C4A-C5A-N7A	-4.02	105.98	110.58
2	A	501	NAP	N3A-C2A-N1A	-3.99	122.54	128.58
2	B	501	NAP	C2A-N3A-C4A	3.80	121.11	111.83
2	A	501	NAP	C2A-N3A-C4A	3.76	121.01	111.83
2	A	501	NAP	N3A-C4A-N9A	3.74	133.53	127.17
2	A	501	NAP	C5A-N7A-C8A	3.60	109.10	103.45
2	B	501	NAP	C4A-C5A-N7A	-3.48	106.60	110.58
2	B	501	NAP	C5A-N7A-C8A	3.38	108.76	103.45
2	B	501	NAP	N9A-C8A-N7A	-3.16	109.45	113.94
2	A	501	NAP	N9A-C8A-N7A	-3.14	109.48	113.94
2	B	501	NAP	C4A-N9A-C8A	2.72	108.59	105.74
2	A	501	NAP	C4A-N9A-C8A	2.47	108.33	105.74

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	NAP	C6A-C5A-N7A	2.19	136.32	132.09

There are no chirality outliers.

All (1) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	501	NAP	PN-O3-PA-O5B

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	NAP	1	0
2	B	501	NAP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	455/465 (97%)	0.06	17 (3%) 45 47	24, 33, 48, 64	0
1	B	455/465 (97%)	0.19	20 (4%) 39 41	25, 35, 51, 70	0
All	All	910/930 (97%)	0.13	37 (4%) 41 43	24, 34, 49, 70	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	336	ASP	3.7
1	A	338	PRO	3.5
1	A	111	ASP	3.3
1	B	292	TRP	3.2
1	B	273	ASP	3.0
1	A	109	GLU	3.0
1	A	375	GLU	2.9
1	B	299	GLU	2.9
1	B	109	GLU	2.7
1	A	274	LYS	2.7
1	A	337	HIS	2.6
1	A	90	ASP	2.6
1	A	273	ASP	2.6
1	B	336	ASP	2.6
1	A	18	ASP	2.6
1	B	375	GLU	2.5
1	A	15	HIS	2.5
1	B	246	GLN	2.5
1	B	338	PRO	2.4
1	A	299	GLU	2.4
1	B	18	ASP	2.4
1	B	395	SER	2.4
1	B	337	HIS	2.3
1	B	281	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	2	ALA	2.3
1	A	281	GLU	2.3
1	A	328	LEU	2.3
1	B	313	ASP	2.2
1	B	2	ALA	2.2
1	B	322	LEU	2.2
1	B	15	HIS	2.2
1	B	135	TYR	2.2
1	B	335	ILE	2.1
1	B	334	ALA	2.1
1	B	372	SER	2.1
1	A	333	GLU	2.1
1	A	334	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAP	A	501	31/48	0.91	0.11	36,45,76,78	0
2	NAP	B	501	31/48	0.91	0.10	41,46,78,80	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAP A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.