



wwPDB EM Validation Summary Report ⓘ

Mar 26, 2026 – 01:32 PM UTC

PDB ID : 7OHW / pdb_00007ohw
EMDB ID : EMD-12911
Title : Nog1-TAP associated immature ribosomal particles from *S. cerevisiae* after rpL25 expression shut down, population B
Authors : Milkereit, P.; Poell, G.
Deposited on : 2021-05-11
Resolution : 3.50 Å(reported)
Based on initial model : 6EM1

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.











Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	13950 (3.00 - 4.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1	3396	
2	2	158	
3	3	306	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	4	278	
5	5	463	
6	6	232	
7	A	291	
8	B	387	
9	C	362	
10	D	505	
11	E	176	
12	F	244	
13	G	256	
14	H	191	
15	J	427	
16	K	376	
17	L	199	
18	M	138	
19	N	204	
20	O	199	
21	P	184	
22	Q	186	
23	S	172	
24	V	137	
25	W	236	
26	Y	127	
27	b	647	
28	e	130	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
29	f	107	
30	h	120	
31	i	100	
32	j	88	
33	m	807	
34	n	605	
35	o	220	
36	r	261	
37	t	322	
38	v	231	
39	x	295	
40	y	245	

2 Entry composition

There are 41 unique types of molecules in this entry. The entry contains 171568 atoms, of which 76920 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called 25S rRNA.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	1	1639	Total	C	H	N	O	P	0	0
			52739	15672	17632	6362	11434	1639		

- Molecule 2 is a RNA chain called 5.8S rRNA.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	2	149	Total	C	H	N	O	P	0	0
			4767	1416	1601	558	1043	149		

- Molecule 3 is a protein called Protein MAK16.

Mol	Chain	Residues	Atoms						AltConf	Trace
3	3	173	Total	C	H	N	O	S	0	0
			2908	901	1474	274	250	9		

- Molecule 4 is a protein called Ribosomal RNA-processing protein 1.

Mol	Chain	Residues	Atoms						AltConf	Trace
4	4	217	Total	C	H	N	O	S	0	0
			3744	1208	1891	319	323	3		

- Molecule 5 is a protein called Ribosome biogenesis protein NSA1.

Mol	Chain	Residues	Atoms						AltConf	Trace
5	5	385	Total	C	H	N	O	S	0	0
			6170	1957	3115	514	573	11		

- Molecule 6 is a RNA chain called ITS2.

Mol	Chain	Residues	Atoms						AltConf	Trace
6	6	65	Total	C	H	N	O	P	0	0
			2061	614	691	228	463	65		

- Molecule 7 is a protein called Ribosome biogenesis protein BRX1.

Mol	Chain	Residues	Atoms						AltConf	Trace
7	A	144	Total	C	H	N	O	S	0	0
			2411	775	1208	217	209	2		

- Molecule 8 is a protein called 60S ribosomal protein L3.

Mol	Chain	Residues	Atoms						AltConf	Trace
8	B	333	Total	C	H	N	O	S	0	0
			5374	1680	2728	490	470	6		

- Molecule 9 is a protein called 60S ribosomal protein L4-A.

Mol	Chain	Residues	Atoms						AltConf	Trace
9	C	343	Total	C	H	N	O	S	0	0
			5336	1643	2725	499	466	3		

- Molecule 10 is a protein called ATP-dependent RNA helicase HAS1.

Mol	Chain	Residues	Atoms						AltConf	Trace
10	D	437	Total	C	H	N	O	S	0	0
			7106	2247	3620	600	627	12		

- Molecule 11 is a protein called 60S ribosomal protein L6-A.

Mol	Chain	Residues	Atoms						AltConf	Trace
11	E	151	Total	C	H	N	O	S	0	0
			2497	780	1292	215	209	1		

- Molecule 12 is a protein called 60S ribosomal protein L7-A.

Mol	Chain	Residues	Atoms						AltConf	Trace
12	F	241	Total	C	H	N	O	S	0	0
			3969	1246	2033	351	338	1		

- Molecule 13 is a protein called 60S ribosomal protein L8-A.

Mol	Chain	Residues	Atoms						AltConf	Trace
13	G	159	Total	C	H	N	O	S	0	0
			2520	794	1289	209	226	2		

- Molecule 14 is a protein called 60S ribosomal protein L9-A.

Mol	Chain	Residues	Atoms						AltConf	Trace
14	H	174	Total	C	H	N	O	S	0	0
			2862	890	1468	254	247	3		

- Molecule 15 is a protein called rRNA-processing protein EBP2.

Mol	Chain	Residues	Atoms						AltConf	Trace
15	J	63	Total	C	H	N	O	S	0	0
			1035	329	506	94	105	1		

- Molecule 16 is a protein called Proteasome-interacting protein CIC1.

Mol	Chain	Residues	Atoms						AltConf	Trace
16	K	257	Total	C	H	N	O	S	0	0
			4230	1337	2157	341	392	3		

- Molecule 17 is a protein called 60S ribosomal protein L13-A.

Mol	Chain	Residues	Atoms						AltConf	Trace
17	L	108	Total	C	H	N	O		0	0
			1782	541	918	180	143			

- Molecule 18 is a protein called 60S ribosomal protein L14-A.

Mol	Chain	Residues	Atoms						AltConf	Trace
18	M	134	Total	C	H	N	O	S	0	0
			2179	668	1138	197	174	2		

- Molecule 19 is a protein called 60S ribosomal protein L15-A.

Mol	Chain	Residues	Atoms						AltConf	Trace
19	N	177	Total	C	H	N	O	S	0	0
			3079	948	1566	320	244	1		

- Molecule 20 is a protein called 60S ribosomal protein L16-A.

Mol	Chain	Residues	Atoms						AltConf	Trace
20	O	197	Total	C	H	N	O	S	0	0
			3215	1003	1660	289	262	1		

- Molecule 21 is a protein called 60S ribosomal protein L17-A.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	P	137	Total	C	H	N	O	0	0
			2139	666	1077	198	198		

- Molecule 22 is a protein called 60S ribosomal protein L18-A.

Mol	Chain	Residues	Atoms					AltConf	Trace	
22	Q	131	Total	C	H	N	O	S	0	0
			2101	645	1092	190	173	1		

- Molecule 23 is a protein called 60S ribosomal protein L20-A.

Mol	Chain	Residues	Atoms						AltConf	Trace
23	S	170	Total	C	H	N	O	S	0	0
			2904	922	1472	265	242	3		

- Molecule 24 is a protein called 60S ribosomal protein L23-A.

Mol	Chain	Residues	Atoms						AltConf	Trace
24	V	98	Total	C	H	N	O	S	0	0
			1498	462	760	136	133	7		

- Molecule 25 is a protein called Ribosome assembly factor MRT4.

Mol	Chain	Residues	Atoms						AltConf	Trace
25	W	185	Total	C	H	N	O	S	0	0
			3049	963	1527	263	292	4		

- Molecule 26 is a protein called 60S ribosomal protein L26-A.

Mol	Chain	Residues	Atoms					AltConf	Trace
26	Y	125	Total	C	H	N	O	0	0
			2060	620	1076	191	173		

- Molecule 27 is a protein called Nucleolar GTP-binding protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace	
27	b	259	Total	C	H	N	O	S	0	0
			4221	1344	2116	355	395	11		

- Molecule 28 is a protein called 60S ribosomal protein L32.

Mol	Chain	Residues	Atoms					AltConf	Trace	
28	e	125	Total	C	H	N	O	S	0	0
			2090	641	1081	203	164	1		

- Molecule 29 is a protein called 60S ribosomal protein L33-A.

Mol	Chain	Residues	Atoms						AltConf	Trace
29	f	106	Total	C	H	N	O	S	0	0
			1731	540	881	165	144	1		

- Molecule 30 is a protein called 60S ribosomal protein L35-A.

Mol	Chain	Residues	Atoms					AltConf	Trace	
30	h	119	Total	C	H	N	O	S	0	0
			2048	615	1079	186	167	1		

- Molecule 31 is a protein called 60S ribosomal protein L36-A.

Mol	Chain	Residues	Atoms						AltConf	Trace
31	i	74	Total	C	H	N	O	S	0	0
			1236	367	642	125	101	1		

- Molecule 32 is a protein called 60S ribosomal protein L37-A.

Mol	Chain	Residues	Atoms						AltConf	Trace
32	j	71	Total	C	H	N	O	S	0	0
			1137	344	571	123	94	5		

- Molecule 33 is a protein called Ribosome biogenesis protein ERB1.

Mol	Chain	Residues	Atoms						AltConf	Trace
33	m	156	Total	C	H	N	O	S	0	0
			2643	847	1318	233	241	4		

- Molecule 34 is a protein called Pescadillo homolog.

Mol	Chain	Residues	Atoms						AltConf	Trace
34	n	334	Total	C	H	N	O	S	0	0
			5516	1787	2782	457	482	8		

- Molecule 35 is a protein called Ribosome biogenesis protein 15.

Mol	Chain	Residues	Atoms						AltConf	Trace
35	o	133	Total	C	H	N	O	S	0	0
			2267	716	1160	198	189	4		

- Molecule 36 is a protein called Ribosome biogenesis protein NSA2.

Mol	Chain	Residues	Atoms						AltConf	Trace
36	r	54	Total	C	H	N	O		0	0
			962	294	485	102	81			

- Molecule 37 is a protein called Ribosome biogenesis protein RLP7.

Mol	Chain	Residues	Atoms						AltConf	Trace
37	t	240	Total	C	H	N	O	S	0	0
			3923	1217	2018	340	345	3		

- Molecule 38 is a protein called Nucleolar protein 16.

Mol	Chain	Residues	Atoms						AltConf	Trace
38	v	130	Total	C	H	N	O	S	0	0
			2223	678	1136	211	195	3		

- Molecule 39 is a protein called Ribosome production factor 1.

Mol	Chain	Residues	Atoms						AltConf	Trace
39	x	267	Total	C	H	N	O	S	0	0
			4573	1444	2305	413	407	4		


- Molecule 40 is a protein called Eukaryotic translation initiation factor 6.

Mol	Chain	Residues	Atoms						AltConf	Trace
40	y	217	Total	C	H	N	O	S	0	0
			3262	1016	1630	280	329	7		

- Molecule 41 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		AltConf
41	j	1	Total	Zn	0
			1	1	






• Molecule 2: 5.8S rRNA


Chain 2: 56% 34% 6%




• Molecule 3: Protein MAK16




- Molecule 4: Ribosomal RNA-processing protein 1




- Molecule 5: Ribosome biogenesis protein NSA1




- Molecule 6: ITS2




- Molecule 7: Ribosome biogenesis protein BRX1




- Molecule 8: 60S ribosomal protein L3




- Molecule 9: 60S ribosomal protein L4-A




- Molecule 10: ATP-dependent RNA helicase HAS1





• Molecule 11: 60S ribosomal protein L6-A

Chain E: 79% 7% 14%




• Molecule 12: 60S ribosomal protein L7-A

Chain F: 91% 8%




• Molecule 13: 60S ribosomal protein L8-A

Chain G: 50% 12% 38%



• Molecule 14: 60S ribosomal protein L9-A

Chain H: 34% 77% 12% 9%



ALA
GLU
GLU
GLU
GLU
GLU
LYS
LYS
LYS

- Molecule 18: 60S ribosomal protein L14-A

Chain M:  87% 10%

MET SER THR D4 D5 I6 V7 K8 A9 D40 Q41 K42 K43 N59 R77 W99 T103 R109 D114 R117 R124 R128 K137 ALA


- Molecule 19: 60S ribosomal protein L15-A

Chain N:  75% 12% 13%

MET G2 A3 E8 E9 R12 V18 L19 R33 A39 R50 G69 ASN ARG LYS ARG ARG VAL PRO LYS GLY ALA THR TYR GLY LYS PRO THR ASN GLN VAL VAL ASN GLU LEU LYS TYR GLN R96 V106 Y119 D124 S125 T126 Y127 E131 V132 R144 D145

N149 V155 R162 R172 K179 K184 A185 G186 K189 K204

- Molecule 20: 60S ribosomal protein L16-A

Chain O:  93% 6%

MET SER V3 D10 S44 F57 A63 G69 R85 S86 R87 V88 S89 H90 V118 V126 K134 L156 Y199

- Molecule 21: 60S ribosomal protein L17-A

Chain P:  67% 7% 26%

MET ALA ARG TYR GLY ALA THR SER THR N10 R30 W39 V51 V58 R69 T70 A71 F76 G77 S87 V91 N97 V114 V119 V120 Q121 A122 P123 K124 GLN ARG ARG ARG ARG THR TYR ARG ALA HIS GLY ARG ARG ILE ASN LYS TYR E140 I146 K159

A160 A161 GLU LYS VAL VAL ARG LEU THR SER ARG GLN ARG GLY ILE ALA GLN LYS ARG ILE ALA ALA

- Molecule 22: 60S ribosomal protein L18-A

Chain Q:  66% 5% 30%

MET GLY ILE ASP ALA THR SER LYS GLN HIS LYS ARG ARG SER GLY H15 L32 L36 A37 R38 R39 T40 V61 V101 T123 L124 D125 R144 N145 SER ARG GLU ALA VAL ARG HIS PHE GLY MET GLY PRO HIS LYS LYS ALA PRO ARG ILE LEU SER THR GLY ARG LYS

PHE GLU ARG ARG GLY ARG ARG SER LYS GLY PHE LYS VAL


- Molecule 23: 60S ribosomal protein L20-A

Chain S:  88% 10% ..




- Molecule 24: 60S ribosomal protein L23-A

Chain V:  16% 66% 5% 28%




- Molecule 25: Ribosome assembly factor MRT4

Chain W:  37% 73% 6% 22%



- Molecule 26: 60S ribosomal protein L26-A


Chain Y:  97% ..



- Molecule 27: Nucleolar GTP-binding protein 1

Chain b:  17% 38% 60%





- Molecule 28: 60S ribosomal protein L32

Chain e: 86% 10%



- Molecule 29: 60S ribosomal protein L33-A


Chain f: 86% 13%




- Molecule 30: 60S ribosomal protein L35-A

Chain h: 84% 15%

Chain n:



- Molecule 36: Ribosome biogenesis protein NSA2



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	37557	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	86.09	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	FEI FALCON III (4k x 4k)	Depositor
Maximum map value	0.134	Depositor
Minimum map value	-0.042	Depositor
Average map value	-0.000	Depositor
Map value standard deviation	0.006	Depositor
Recommended contour level	0.021	Depositor
Map size (Å)	425.40002, 425.40002, 425.40002	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.0635, 1.0635, 1.0635	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	1	0.08	0/39278	0.23	0/61189
2	2	0.08	0/3536	0.22	0/5501
3	3	0.12	0/1461	0.29	0/1958
4	4	0.11	0/1895	0.26	0/2549
5	5	0.07	0/3109	0.21	0/4187
6	6	0.08	0/1527	0.26	0/2371
7	A	0.12	0/1235	0.26	0/1669
8	B	0.11	0/2699	0.25	0/3626
9	C	0.10	0/2660	0.25	0/3601
10	D	0.09	0/3552	0.23	0/4789
11	E	0.10	0/1226	0.22	0/1648
12	F	0.10	0/1974	0.23	0/2654
13	G	0.11	0/1252	0.26	0/1695
14	H	0.10	0/1412	0.27	0/1898
15	J	0.11	0/538	0.23	0/723
16	K	0.09	0/2107	0.24	0/2845
17	L	0.14	0/877	0.33	0/1179
18	M	0.10	0/1056	0.22	0/1421
19	N	0.12	0/1544	0.22	0/2065
20	O	0.10	0/1585	0.23	0/2128
21	P	0.09	0/1080	0.20	0/1455
22	Q	0.12	0/1024	0.27	0/1385
23	S	0.10	0/1468	0.24	0/1973
24	V	0.10	0/747	0.22	0/1002
25	W	0.09	0/1543	0.23	0/2068
26	Y	0.10	0/995	0.23	0/1329
27	b	0.09	0/2133	0.26	0/2857
28	e	0.11	0/1030	0.23	0/1379
29	f	0.11	0/868	0.25	0/1168
30	h	0.11	0/978	0.29	0/1301
31	i	0.13	0/599	0.26	0/793
32	j	0.13	0/578	0.25	0/767

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
33	m	0.10	0/1363	0.24	0/1841
34	n	0.10	0/2802	0.23	0/3791
35	o	0.10	0/1129	0.23	0/1502
36	r	0.08	0/485	0.22	0/636
37	t	0.11	0/1930	0.26	0/2596
38	v	0.11	0/1100	0.22	0/1456
39	x	0.11	0/2313	0.27	0/3100
40	y	0.08	0/1650	0.23	0/2242
All	All	0.09	0/100338	0.24	0/144337

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
14	H	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
14	H	22	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	35107	17632	17660	387	0
2	2	3166	1601	1603	42	0
3	3	1434	1474	1473	13	0
4	4	1853	1891	1887	18	0
5	5	3055	3115	3113	15	0
6	6	1370	691	692	27	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	1203	1208	1205	10	0
8	B	2646	2728	2726	45	0
9	C	2611	2725	2724	30	0
10	D	3486	3620	3618	22	0
11	E	1205	1292	1291	10	0
12	F	1936	2033	2032	12	0
13	G	1231	1289	1287	20	0
14	H	1394	1468	1464	20	0
15	J	529	506	504	2	0
16	K	2073	2157	2155	42	0
17	L	864	918	917	16	0
18	M	1041	1138	1137	11	0
19	N	1513	1566	1564	20	0
20	O	1555	1660	1659	9	0
21	P	1062	1077	1075	10	0
22	Q	1009	1092	1091	6	0
23	S	1432	1472	1470	12	0
24	V	738	760	757	5	0
25	W	1522	1527	1520	10	0
26	Y	984	1076	1075	3	0
27	b	2105	2116	2106	8	0
28	e	1009	1081	1080	10	0
29	f	850	881	880	11	0
30	h	969	1079	1078	18	0
31	i	594	642	641	12	0
32	j	566	571	566	10	0
33	m	1325	1318	1317	12	0
34	n	2734	2782	2778	18	0
35	o	1107	1160	1159	6	0
36	r	477	485	484	5	0
37	t	1905	2018	2015	17	0
38	v	1087	1136	1133	12	0
39	x	2268	2305	2302	14	0
40	y	1632	1630	1628	15	0
41	j	1	0	0	0	0
All	All	94648	76920	76866	827	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 827 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:1:3198:U:O2'	1:1:3199:G:OP1	1.81	0.99
1:1:625:G:O2'	1:1:1401:A:OP1	1.85	0.94
1:1:2899:C:O2'	1:1:2901:G:OP2	1.86	0.93
1:1:1347:U:O2'	1:1:1348:U:OP1	1.90	0.90
1:1:3228:C:O2'	1:1:3229:G:OP2	1.90	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	3	171/306 (56%)	161 (94%)	9 (5%)	1 (1%)	21	54
4	4	209/278 (75%)	191 (91%)	18 (9%)	0	100	100
5	5	377/463 (81%)	364 (97%)	13 (3%)	0	100	100
7	A	138/291 (47%)	124 (90%)	14 (10%)	0	100	100
8	B	329/387 (85%)	316 (96%)	13 (4%)	0	100	100
9	C	341/362 (94%)	323 (95%)	18 (5%)	0	100	100
10	D	433/505 (86%)	413 (95%)	20 (5%)	0	100	100
11	E	147/176 (84%)	142 (97%)	5 (3%)	0	100	100
12	F	239/244 (98%)	236 (99%)	3 (1%)	0	100	100
13	G	155/256 (60%)	151 (97%)	4 (3%)	0	100	100
14	H	166/191 (87%)	159 (96%)	6 (4%)	1 (1%)	21	54
15	J	59/427 (14%)	59 (100%)	0	0	100	100
16	K	253/376 (67%)	240 (95%)	13 (5%)	0	100	100
17	L	106/199 (53%)	97 (92%)	7 (7%)	2 (2%)	6	33
18	M	132/138 (96%)	130 (98%)	2 (2%)	0	100	100
19	N	173/204 (85%)	171 (99%)	2 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
20	O	195/199 (98%)	192 (98%)	3 (2%)	0	100	100
21	P	133/184 (72%)	131 (98%)	2 (2%)	0	100	100
22	Q	129/186 (69%)	128 (99%)	1 (1%)	0	100	100
23	S	168/172 (98%)	156 (93%)	12 (7%)	0	100	100
24	V	92/137 (67%)	91 (99%)	1 (1%)	0	100	100
25	W	171/236 (72%)	167 (98%)	4 (2%)	0	100	100
26	Y	123/127 (97%)	122 (99%)	1 (1%)	0	100	100
27	b	237/647 (37%)	230 (97%)	7 (3%)	0	100	100
28	e	123/130 (95%)	120 (98%)	3 (2%)	0	100	100
29	f	104/107 (97%)	99 (95%)	5 (5%)	0	100	100
30	h	117/120 (98%)	109 (93%)	8 (7%)	0	100	100
31	i	72/100 (72%)	70 (97%)	2 (3%)	0	100	100
32	j	69/88 (78%)	65 (94%)	4 (6%)	0	100	100
33	m	152/807 (19%)	145 (95%)	7 (5%)	0	100	100
34	n	326/605 (54%)	317 (97%)	9 (3%)	0	100	100
35	o	131/220 (60%)	127 (97%)	4 (3%)	0	100	100
36	r	50/261 (19%)	50 (100%)	0	0	100	100
37	t	234/322 (73%)	222 (95%)	12 (5%)	0	100	100
38	v	124/231 (54%)	118 (95%)	6 (5%)	0	100	100
39	x	261/295 (88%)	243 (93%)	18 (7%)	0	100	100
40	y	211/245 (86%)	206 (98%)	5 (2%)	0	100	100
All	All	6650/10222 (65%)	6385 (96%)	261 (4%)	4 (0%)	49	79

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	3	134	TYR
17	L	63	VAL
14	H	23	ARG
17	L	61	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	3	155/274 (57%)	155 (100%)	0	100	100
4	4	203/257 (79%)	202 (100%)	1 (0%)	81	80
5	5	343/410 (84%)	343 (100%)	0	100	100
7	A	136/263 (52%)	132 (97%)	4 (3%)	37	60
8	B	280/323 (87%)	277 (99%)	3 (1%)	65	74
9	C	273/289 (94%)	270 (99%)	3 (1%)	65	74
10	D	381/440 (87%)	375 (98%)	6 (2%)	55	70
11	E	131/153 (86%)	131 (100%)	0	100	100
12	F	204/205 (100%)	204 (100%)	0	100	100
13	G	128/208 (62%)	126 (98%)	2 (2%)	55	70
14	H	157/171 (92%)	156 (99%)	1 (1%)	78	79
15	J	58/383 (15%)	58 (100%)	0	100	100
16	K	238/346 (69%)	235 (99%)	3 (1%)	61	72
17	L	87/159 (55%)	86 (99%)	1 (1%)	65	74
18	M	106/109 (97%)	106 (100%)	0	100	100
19	N	153/176 (87%)	153 (100%)	0	100	100
20	O	160/162 (99%)	158 (99%)	2 (1%)	61	72
21	P	109/146 (75%)	109 (100%)	0	100	100
22	Q	107/151 (71%)	107 (100%)	0	100	100
23	S	155/156 (99%)	154 (99%)	1 (1%)	78	79
24	V	77/105 (73%)	76 (99%)	1 (1%)	61	72
25	W	169/213 (79%)	169 (100%)	0	100	100
26	Y	108/110 (98%)	108 (100%)	0	100	100
27	b	231/573 (40%)	230 (100%)	1 (0%)	84	81
28	e	108/111 (97%)	108 (100%)	0	100	100
29	f	90/91 (99%)	90 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
30	h	104/105 (99%)	104 (100%)	0	100	100
31	i	61/82 (74%)	59 (97%)	2 (3%)	33	58
32	j	59/71 (83%)	59 (100%)	0	100	100
33	m	145/723 (20%)	144 (99%)	1 (1%)	76	78
34	n	302/548 (55%)	301 (100%)	1 (0%)	86	83
35	o	118/199 (59%)	117 (99%)	1 (1%)	73	77
36	r	48/229 (21%)	48 (100%)	0	100	100
37	t	213/287 (74%)	212 (100%)	1 (0%)	81	80
38	v	116/205 (57%)	116 (100%)	0	100	100
39	x	252/276 (91%)	249 (99%)	3 (1%)	63	73
40	y	185/211 (88%)	184 (100%)	1 (0%)	81	80
All	All	5950/8920 (67%)	5911 (99%)	39 (1%)	73	78

5 of 39 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
27	b	70	ASN
39	x	163	ILE
31	i	43	LEU
34	n	229	LEU
39	x	175	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 44 such sidechains are listed below:

Mol	Chain	Res	Type
27	b	267	GLN
35	o	133	HIS
28	e	49	ASN
30	h	76	GLN
39	x	102	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	1	1616/3396 (47%)	297 (18%)	27 (1%)
2	2	147/158 (93%)	25 (17%)	1 (0%)

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
6	6	64/232 (27%)	26 (40%)	2 (3%)
All	All	1827/3786 (48%)	348 (19%)	30 (1%)

5 of 348 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	1	2	U
1	1	3	U
1	1	20	A
1	1	26	A
1	1	34	A

5 of 30 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	1	659	G
2	2	114	G
1	1	1128	U
6	6	227	C
1	1	3228	C

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-12911. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 188




Y Index: 146




Z Index: 155


6.3.2 Raw map



X Index: 188



Y Index: 146




Z Index: 155

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.021. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 1238 nm³; this corresponds to an approximate mass of 1119 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.286 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of $0.286 \text{ \AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.50	-	-
Author-provided FSC curve	3.47	4.15	3.55
Unmasked-calculated*	4.36	8.06	4.55

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.36 differs from the reported value 3.5 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-12911 and PDB model 7OHW. Per-residue inclusion information can be found in [section 3](#) on [page 11](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.021 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.021).

9.4 Atom inclusion [i](#)



At the recommended contour level, 85% of all backbone atoms, 87% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.021) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8690	 0.3610
1	 0.9300	 0.3520
2	 0.9720	 0.3850
3	 0.8540	 0.4220
4	 0.8920	 0.3820
5	 0.8880	 0.4480
6	 0.9780	 0.3280
A	 0.7350	 0.2860
B	 0.8410	 0.2870
C	 0.8780	 0.4550
D	 0.8210	 0.3550
E	 0.9090	 0.4400
F	 0.8910	 0.4320
G	 0.8540	 0.4110
H	 0.4820	 0.3300
J	 0.6960	 0.2530
K	 0.8630	 0.3090
L	 0.9010	 0.4410
M	 0.9000	 0.4070
N	 0.8590	 0.4280
O	 0.8590	 0.4000
P	 0.8090	 0.4230
Q	 0.8780	 0.4360
S	 0.8960	 0.3990
V	 0.5940	 0.1830
W	 0.4260	 0.2450
Y	 0.8920	 0.4560
b	 0.5000	 0.2430
e	 0.8470	 0.4660
f	 0.9000	 0.4820
h	 0.8290	 0.3920
i	 0.8080	 0.3600
j	 0.9130	 0.4570
m	 0.8350	 0.3270
n	 0.8480	 0.2980



Continued on next page...

Continued from previous page...

Chain	Atom inclusion	Q-score
o	 0.9070	 0.3430
r	 0.5460	 0.2960
t	 0.8550	 0.3220
v	 0.8760	 0.4130
x	 0.8370	 0.4080
y	 0.6760	 0.1890