



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:43 PM UTC

PDB ID : 8OHW / pdb_00008ohw
Title : Crystal structure of heparanase from Burkholderia pseudomallei in complex with siastatin B derived inhibitor
Authors : Armstrong, Z.; Yurong, C.; Wu, L.; Overkleeft, H.S.; Davies, G.J.
Deposited on : 2023-03-21
Resolution : 1.27 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2836 (1.30-1.26)
Clashscore	190562	2911 (1.30-1.26)
Ramachandran outliers	187476	2841 (1.30-1.26)
Sidechain outliers	187428	2840 (1.30-1.26)
RSRZ outliers	180081	2832 (1.30-1.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	438	4% 92% 7%
1	BBB	438	9% 96% .

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 7464 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycoside hydrolase family 44 domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	437	Total	C	N	O	S	0	14	0
			3246	2039	573	625	9			
1	BBB	437	Total	C	N	O	S	0	7	0
			3195	2010	560	615	10			

- Molecule 2 is (3 {S},4 {S},5 {S},6 {R})-4,5,6-tris(oxidanyl)piperidine-3-carboxylic acid (CCD ID: VON) (formula: C₆H₁₁NO₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	AAA	1	Total	C	N	O	0	0
			12	6	1	5		
2	BBB	1	Total	C	N	O	0	0
			12	6	1	5		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	AAA	1	Total C O 4 2 2	0	0
3	AAA	1	Total C O 4 2 2	0	0
3	AAA	1	Total C O 4 2 2	0	0
3	AAA	1	Total C O 4 2 2	0	0
3	BBB	1	Total C O 4 2 2	0	0
3	BBB	1	Total C O 4 2 2	0	0
3	BBB	1	Total C O 4 2 2	0	0
3	BBB	1	Total C O 4 2 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	AAA	531	Total O 531 531	0	0
4	BBB	436	Total O 436 436	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycoside hydrolase family 44 domain-containing protein



- Molecule 1: Glycoside hydrolase family 44 domain-containing protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	76.37Å 104.61Å 113.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	61.76 – 1.27 61.76 – 1.27	Depositor EDS
% Data completeness (in resolution range)	99.9 (61.76-1.27) 99.9 (61.76-1.27)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.73 (at 1.27Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.160 , 0.181 0.161 , 0.182	Depositor DCC
R_{free} test set	12173 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	15.2	Xtriage
Anisotropy	0.218	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 33.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7464	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, VON

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.12	3/3319 (0.1%)	1.14	4/4535 (0.1%)
1	BBB	1.11	1/3266 (0.0%)	1.18	0/4460
All	All	1.12	4/6585 (0.1%)	1.16	4/8995 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	47	GLY	C-O	5.55	1.31	1.24
1	BBB	151	HIS	CE1-NE2	5.29	1.37	1.32
1	AAA	317	ILE	C-O	5.07	1.29	1.24
1	AAA	333	GLY	C-O	5.07	1.30	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	445	PHE	CA-CB-CG	6.71	120.51	113.80
1	AAA	160	TYR	O-C-N	-5.46	115.62	120.48
1	AAA	385	ARG	N-CA-C	-5.14	106.52	112.89
1	AAA	238	GLN	O-C-N	5.11	127.54	122.12

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	3246	0	3139	30	0
1	BBB	3195	0	3099	18	0
2	AAA	12	0	0	0	0
2	BBB	12	0	0	0	0
3	AAA	16	0	24	4	0
3	BBB	16	0	24	0	0
4	AAA	531	0	0	4	2
4	BBB	436	0	0	5	2
All	All	7464	0	6286	42	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:308:ASN:HB3	4:AAA:974:HOH:O	1.57	1.02
1:AAA:156:ASN:OD1	3:AAA:504:EDO:H11	1.67	0.94
1:BBB:236[B]:MET:SD	4:BBB:689:HOH:O	2.36	0.83
1:AAA:382[B]:PRO:HG3	1:BBB:135:GLN:HG2	1.63	0.81
1:AAA:236:MET:SD	1:AAA:240[B]:LEU:HD12	2.24	0.77
1:AAA:382[B]:PRO:HG3	1:BBB:135:GLN:CG	2.17	0.75
1:AAA:382[B]:PRO:O	1:BBB:101:GLN:OE1	2.05	0.74
1:BBB:101:GLN:HA	1:BBB:104:ARG:NH1	2.11	0.66
1:AAA:446:ALA:OXT	1:BBB:104:ARG:NE	2.35	0.59
1:BBB:308:ASN:HB3	4:BBB:925:HOH:O	2.05	0.55
1:AAA:308:ASN:HB3	4:AAA:978:HOH:O	2.06	0.54
1:AAA:128:ALA:O	1:AAA:132[B]:LEU:HD13	2.08	0.54
1:BBB:17[B]:MET:HE1	1:BBB:361:LEU:HD13	1.90	0.54
1:AAA:236:MET:SD	1:AAA:240[B]:LEU:CD1	2.96	0.54
1:AAA:382[B]:PRO:CG	1:BBB:135:GLN:HG2	2.36	0.53
1:BBB:356:GLY:HA3	4:BBB:603:HOH:O	2.08	0.53
1:AAA:190[B]:TRP:CZ2	3:AAA:505:EDO:H21	2.44	0.53
1:AAA:17[B]:MET:HE1	1:AAA:361:LEU:HD13	1.91	0.53
1:BBB:100:VAL:O	1:BBB:104:ARG:NH1	2.44	0.50
1:AAA:381[A]:VAL:HG12	1:AAA:384:ALA:HB2	1.94	0.50
1:BBB:441:LEU:C	1:BBB:441:LEU:HD12	2.38	0.48
1:AAA:132[B]:LEU:N	1:AAA:132[B]:LEU:CD1	2.76	0.48
1:AAA:132[B]:LEU:N	1:AAA:132[B]:LEU:HD12	2.29	0.47
1:BBB:360:ARG:NH2	4:BBB:603:HOH:O	2.40	0.47
1:AAA:190[B]:TRP:CZ2	1:AAA:260:TRP:CE3	3.03	0.46
1:AAA:383[A]:ASN:ND2	4:AAA:601:HOH:O	2.32	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:210:LEU:HD23	1:AAA:240[B]:LEU:HD23	1.97	0.46
1:AAA:391:THR:CG2	3:AAA:503:EDO:H11	2.45	0.46
1:AAA:17[B]:MET:HE1	1:AAA:361:LEU:HD22	1.98	0.46
1:AAA:106[A]:ARG:HD2	1:AAA:134:VAL:O	2.16	0.46
1:AAA:106[B]:ARG:NH2	4:AAA:607:HOH:O	2.49	0.45
1:AAA:104:ARG:NH1	1:BBB:356:GLY:O	2.49	0.45
1:BBB:136:LEU:HD21	1:BBB:139:PHE:CZ	2.53	0.44
1:AAA:156:ASN:OD1	3:AAA:504:EDO:C1	2.54	0.43
1:BBB:392:LEU:C	1:BBB:392:LEU:HD23	2.44	0.43
1:AAA:136:LEU:HD21	1:AAA:139:PHE:CZ	2.54	0.43
1:AAA:392:LEU:C	1:AAA:392:LEU:HD23	2.44	0.42
1:BBB:356:GLY:CA	4:BBB:603:HOH:O	2.67	0.42
1:AAA:284:VAL:HG12	1:AAA:289:ALA:HB3	2.02	0.42
1:AAA:381[A]:VAL:CG1	1:AAA:384:ALA:HB2	2.50	0.42
1:AAA:210:LEU:O	1:AAA:251:PHE:HA	2.21	0.41
1:BBB:17[B]:MET:HB3	1:BBB:17[B]:MET:HE2	1.75	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:AAA:1028:HOH:O	4:BBB:895:HOH:O[4_455]	2.16	0.04
4:AAA:616:HOH:O	4:BBB:1004:HOH:O[4_455]	2.17	0.03

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	449/438 (102%)	438 (98%)	11 (2%)	0	100	100
1	BBB	440/438 (100%)	431 (98%)	9 (2%)	0	100	100
All	All	889/876 (102%)	869 (98%)	20 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	320/310 (103%)	317 (99%)	3 (1%)	70	40
1	BBB	315/310 (102%)	312 (99%)	3 (1%)	68	35
All	All	635/620 (102%)	629 (99%)	6 (1%)	68	40

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	156	ASN
1	AAA	302	TYR
1	AAA	385	ARG
1	BBB	302	TYR
1	BBB	367	ASP
1	BBB	385	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	AAA	505	-	3,3,3	0.36	0	2,2,2	0.27	0
3	EDO	BBB	502	-	3,3,3	0.17	0	2,2,2	0.23	0
3	EDO	BBB	503	-	3,3,3	0.51	0	2,2,2	0.25	0
2	VON	BBB	501	-	11,12,12	1.67	2 (18%)	11,17,17	1.83	3 (27%)
2	VON	AAA	501	-	11,12,12	1.37	1 (9%)	11,17,17	1.77	4 (36%)
3	EDO	AAA	502	-	3,3,3	0.29	0	2,2,2	0.33	0
3	EDO	BBB	504	-	3,3,3	0.17	0	2,2,2	0.03	0
3	EDO	AAA	503	-	3,3,3	0.48	0	2,2,2	1.16	0
3	EDO	AAA	504	-	3,3,3	0.47	0	2,2,2	0.40	0
3	EDO	BBB	505	-	3,3,3	0.27	0	2,2,2	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	AAA	505	-	-	0/1/1/1	-
3	EDO	BBB	502	-	-	1/1/1/1	-
3	EDO	BBB	503	-	-	1/1/1/1	-
2	VON	BBB	501	-	-	0/4/21/21	0/1/1/1
2	VON	AAA	501	-	-	4/4/21/21	0/1/1/1
3	EDO	AAA	502	-	-	1/1/1/1	-
3	EDO	BBB	504	-	-	0/1/1/1	-
3	EDO	AAA	503	-	-	1/1/1/1	-
3	EDO	AAA	504	-	-	0/1/1/1	-
3	EDO	BBB	505	-	-	0/1/1/1	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	BBB	501	VON	C7-N1	3.76	1.52	1.47
2	AAA	501	VON	C7-N1	2.74	1.51	1.47
2	BBB	501	VON	O12-C6	-2.16	1.23	1.30

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	BBB	501	VON	O11-C6-C5	-3.93	113.72	123.01
2	AAA	501	VON	C7-N1-C2	3.05	114.72	108.90
2	BBB	501	VON	O10-C4-C5	-2.72	105.14	110.59
2	AAA	501	VON	O11-C6-C5	-2.64	116.77	123.01
2	AAA	501	VON	O10-C4-C5	-2.52	105.53	110.59
2	BBB	501	VON	O12-C6-C5	2.43	120.23	113.93
2	AAA	501	VON	C4-C3-C2	2.10	112.94	109.86

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	BBB	502	EDO	O1-C1-C2-O2
3	AAA	502	EDO	O1-C1-C2-O2
2	AAA	501	VON	C4-C5-C6-O11
2	AAA	501	VON	C4-C5-C6-O12
3	AAA	503	EDO	O1-C1-C2-O2
2	AAA	501	VON	C7-C5-C6-O11
2	AAA	501	VON	C7-C5-C6-O12
3	BBB	503	EDO	O1-C1-C2-O2


There are no ring outliers.


3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	AAA	505	EDO	1	0
3	AAA	503	EDO	1	0
3	AAA	504	EDO	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	437/438 (99%)	0.19	18 (4%) 41 44	5, 14, 26, 52	14 (3%)
1	BBB	437/438 (99%)	0.63	39 (8%) 15 16	6, 18, 38, 74	7 (1%)
All	All	874/876 (99%)	0.41	57 (6%) 25 27	5, 16, 33, 74	21 (2%)

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	BBB	446	ALA	7.7
1	BBB	418	PRO	6.3
1	BBB	417	ALA	6.1
1	BBB	420	GLY	5.4
1	AAA	384	ALA	5.1
1	AAA	446	ALA	5.1
1	BBB	9	PRO	5.0
1	BBB	416	GLY	4.9
1	AAA	190[A]	TRP	4.7
1	BBB	220	SER	4.5
1	BBB	430	VAL	4.3
1	AAA	220	SER	4.3
1	BBB	426	CYS	4.2
1	AAA	382[A]	PRO	4.1
1	BBB	383	ASN	4.0
1	BBB	190	TRP	4.0
1	BBB	384	ALA	3.8
1	AAA	385	ARG	3.7
1	BBB	378	CYS	3.7
1	AAA	383[A]	ASN	3.6
1	BBB	425	ALA	3.4
1	AAA	219	ASP	3.2
1	BBB	385	ARG	3.2
1	BBB	356	GLY	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	AAA	426	CYS	2.9
1	BBB	10	SER	2.8
1	BBB	104	ARG	2.8
1	AAA	356	GLY	2.7
1	BBB	219	ASP	2.7
1	BBB	388	ILE	2.7
1	BBB	386	ALA	2.6
1	AAA	10	SER	2.5
1	BBB	387	GLY	2.5
1	BBB	421	GLY	2.5
1	AAA	424	VAL	2.5
1	BBB	429	GLY	2.4
1	BBB	405	ILE	2.4
1	BBB	11	SER	2.3
1	AAA	240[A]	LEU	2.3
1	BBB	374	VAL	2.3
1	BBB	433	VAL	2.3
1	BBB	423	PRO	2.3
1	AAA	194	ARG	2.3
1	BBB	427	VAL	2.3
1	BBB	422	ARG	2.2
1	AAA	425	ALA	2.2
1	BBB	229	PRO	2.1
1	BBB	389	VAL	2.1
1	AAA	379	SER	2.1
1	BBB	379	SER	2.1
1	AAA	156	ASN	2.1
1	AAA	247	ARG	2.1
1	BBB	381	VAL	2.1
1	BBB	357	GLY	2.0
1	BBB	391	THR	2.0
1	BBB	382	PRO	2.0
1	BBB	344	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	AAA	502	4/4	0.77	0.21	36,46,51,60	0
3	EDO	BBB	502	4/4	0.77	0.16	36,44,44,47	0
3	EDO	AAA	503	4/4	0.88	0.10	25,26,28,32	0
3	EDO	BBB	504	4/4	0.91	0.12	28,28,33,33	0
3	EDO	AAA	504	4/4	0.92	0.11	28,29,30,31	0
3	EDO	AAA	505	4/4	0.92	0.11	22,30,30,30	0
3	EDO	BBB	505	4/4	0.92	0.11	26,30,31,33	0
3	EDO	BBB	503	4/4	0.93	0.10	24,27,29,35	0
2	VON	BBB	501	12/12	0.98	0.05	12,13,15,16	0
2	VON	AAA	501	12/12	0.99	0.04	11,12,14,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around VON BBB 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around VON AAA 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.