



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:08 PM UTC

PDB ID : 1OIJ / pdb_00001oij
Title : Crystal structure of the alkylsulfatase AtsK, a non-heme Fe(II) alphaketoglutarate dependent Dioxygenase in complex with alphaketoglutarate
Authors : Mueller, I.; Kahnert, A.; Pape, T.; Dierks, T.; Meyer-Klauke, W.; Kertesz, M.A.; Uson, I.
Deposited on : 2003-06-18
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	6% 78% 5% • 16%
2	B	301	7% 79% 5% • 15%
2	D	301	4% 74% • • 22%
3	C	301	7% 72% 5% • 21%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8458 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PUTATIVE ALKYL SULFATASE ATSK.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	A	254	Total	C	N	O	0	0	0
			1995	1261	370	364			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	165	ARG	GLU	conflict	UNP Q9WWU5

- Molecule 2 is a protein called PUTATIVE ALKYL SULFATASE ATSK.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	255	Total	C	N	O	0	1	0
			2002	1264	368	370			
2	D	236	Total	C	N	O	0	1	0
			1850	1170	340	340			

- Molecule 3 is a protein called PUTATIVE ALKYL SULFATASE ATSK.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	238	Total	C	N	O	0	1	0
			1876	1186	346	344			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	293	TYR	GLN	conflict	UNP Q9WWU5

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Na	0	0
			1	1		
4	C	1	Total	Na	0	0
			1	1		
4	D	1	Total	Na	0	0
			1	1		

- Molecule 5 is 2-OXOGLUTARIC ACID (CCD ID: AKG) (formula: $C_5H_6O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			10	5	5		
5	B	1	Total	C	O	0	0
			10	5	5		
5	C	1	Total	C	O	0	0
			10	5	5		
5	D	1	Total	C	O	0	0
			10	5	5		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	171	Total	O	0	0
			171	171		
6	B	191	Total	O	0	0
			191	191		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	163	Total 163	O 163	0	0
6	D	166	Total 166	O 166	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: PUTATIVE ALKYL SULFATASE ATSK




• Molecule 2: PUTATIVE ALKYL SULFATASE ATSK




• Molecule 2: PUTATIVE ALKYL SULFATASE ATSK



• Molecule 3: PUTATIVE ALKYL SULFATASE ATSK





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.45Å 144.88Å 160.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.92 – 2.10 38.92 – 2.10	Depositor EDS
% Data completeness (in resolution range)	100.0 (38.92-2.10) 99.7 (38.92-2.10)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 2.06Å)	Xtriage
Refinement program	REFMAC 5.1.19	Depositor
R, R_{free}	0.191 , 0.215 (Not available) , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	32.0	Xtriage
Anisotropy	0.414	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8458	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.05 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.5133e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, AKG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.05	3/2039 (0.1%)	1.02	2/2775 (0.1%)
2	B	1.02	1/2050 (0.0%)	1.03	6/2791 (0.2%)
2	D	0.97	0/1897	1.04	4/2585 (0.2%)
3	C	0.99	2/1924 (0.1%)	1.07	7/2621 (0.3%)
All	All	1.01	6/7910 (0.1%)	1.04	19/10772 (0.2%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	149	GLU	CA-CB	7.18	1.62	1.53
1	A	111	VAL	CA-CB	6.86	1.63	1.54
3	C	204	ILE	CA-CB	6.02	1.61	1.55
2	B	111	VAL	CA-CB	5.97	1.62	1.54
3	C	24	ILE	CA-CB	5.07	1.61	1.54
1	A	245	VAL	CA-CB	5.06	1.62	1.54

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	218	ARG	NE-CZ-NH1	-9.21	112.29	121.50
3	C	218	ARG	NE-CZ-NH2	9.15	127.44	119.20
2	B	296	ARG	NE-CZ-NH2	-8.12	111.89	119.20
2	D	51	ARG	NE-CZ-NH2	7.61	126.05	119.20
3	C	51	ARG	NE-CZ-NH2	7.18	125.67	119.20
2	D	299	ARG	NE-CZ-NH2	7.09	125.58	119.20
2	D	51	ARG	NE-CZ-NH1	-6.92	114.58	121.50
3	C	51	ARG	NE-CZ-NH1	-6.70	114.80	121.50
3	C	299	ARG	NE-CZ-NH2	6.56	125.10	119.20
2	B	296	ARG	CD-NE-CZ	6.44	133.42	124.40
2	D	299	ARG	NE-CZ-NH1	-6.37	115.13	121.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	218	ARG	CD-NE-CZ	6.22	133.11	124.40
3	C	299	ARG	NE-CZ-NH1	-6.21	115.29	121.50
2	B	296	ARG	NE-CZ-NH1	6.16	127.66	121.50
1	A	51	ARG	CD-NE-CZ	5.63	132.28	124.40
2	B	51	ARG	CD-NE-CZ	5.21	131.70	124.40
2	B	51	ARG	NE-CZ-NH2	-5.18	114.54	119.20
1	A	296	ARG	NE-CZ-NH1	-5.15	116.35	121.50
2	B	97	ASP	N-CA-C	5.14	116.47	110.41

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1995	0	1975	8	0
2	B	2002	0	1968	8	0
2	D	1850	0	1815	6	0
3	C	1876	0	1840	14	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	10	0	4	0	0
5	B	10	0	4	0	0
5	C	10	0	4	0	0
5	D	10	0	4	0	0
6	A	171	0	0	1	0
6	B	191	0	0	2	0
6	C	163	0	0	0	0
6	D	166	0	0	0	0
All	All	8458	0	7614	36	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:77:GLU:HG2	3:C:283:ALA:HB3	1.62	0.82
3:C:291:ASP:OD1	3:C:293:TYR:CD2	2.40	0.75
3:C:291:ASP:OD1	3:C:293:TYR:CE2	2.46	0.69
2:D:117:TYR:OH	2:D:211[B]:GLN:NE2	2.25	0.69
2:D:105:ASN:HD22	2:D:105:ASN:H	1.43	0.64
1:A:105:ASN:H	1:A:105:ASN:HD22	1.46	0.62
2:B:105:ASN:H	2:B:105:ASN:HD22	1.48	0.61
3:C:77:GLU:CG	3:C:283:ALA:HB3	2.31	0.60
3:C:200:ARG:HB2	3:C:211[A]:GLN:HG2	1.83	0.60
3:C:105:ASN:HD22	3:C:105:ASN:H	1.49	0.60
2:B:103:ARG:HB3	6:B:2169:HOH:O	2.03	0.58
3:C:159:TRP:HZ2	3:C:293:TYR:CD2	2.24	0.56
3:C:193:GLU:HB2	3:C:299:ARG:HB2	1.88	0.56
1:A:193:GLU:HB2	1:A:299:ARG:HB2	1.90	0.54
2:D:147:LEU:HB2	2:D:152:ARG:HG2	1.91	0.53
2:D:193:GLU:HB2	2:D:299:ARG:HB2	1.88	0.53
2:B:105:ASN:HD22	2:B:105:ASN:N	2.08	0.51
3:C:291:ASP:OD2	3:C:293:TYR:CZ	2.63	0.51
2:B:64:ASP:HB2	6:B:2049:HOH:O	2.10	0.50
2:B:193:GLU:HB2	2:B:299:ARG:HB2	1.93	0.50
1:A:105:ASN:HD22	1:A:105:ASN:N	2.07	0.47
1:A:211:GLN:HG3	6:A:2103:HOH:O	2.13	0.47
2:D:105:ASN:HD22	2:D:105:ASN:N	2.12	0.47
2:B:200:ARG:HB2	2:B:211[A]:GLN:HG2	1.96	0.47
1:A:117:TYR:HB2	1:A:118:PRO:HD2	1.98	0.46
2:D:65:ASP:OD1	2:D:278:ARG:HD3	2.15	0.46
3:C:291:ASP:OD2	3:C:293:TYR:CE1	2.69	0.45
3:C:147:LEU:HB2	3:C:152:ARG:HG2	1.97	0.45
1:A:60:GLN:HA	1:A:62:HIS:CE1	2.52	0.45
1:A:34:SER:HB2	1:A:35:PRO:CD	2.47	0.44
3:C:77:GLU:HG2	3:C:283:ALA:CB	2.39	0.42
3:C:105:ASN:HD22	3:C:105:ASN:N	2.14	0.42
2:B:117:TYR:HB2	2:B:118:PRO:HD2	2.01	0.41
3:C:117:TYR:OH	3:C:211[A]:GLN:NE2	2.54	0.41
1:A:117:TYR:HB2	1:A:118:PRO:CD	2.50	0.41
2:B:149:GLU:N	2:B:150:PRO:CD	2.84	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	246/301 (82%)	239 (97%)	7 (3%)	0	100	100
2	B	248/301 (82%)	240 (97%)	8 (3%)	0	100	100
2	D	231/301 (77%)	224 (97%)	7 (3%)	0	100	100
3	C	233/301 (77%)	224 (96%)	9 (4%)	0	100	100
All	All	958/1204 (80%)	927 (97%)	31 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	207/244 (85%)	200 (97%)	7 (3%)	32	35
2	B	208/244 (85%)	203 (98%)	5 (2%)	43	49
2	D	191/244 (78%)	188 (98%)	3 (2%)	55	64
3	C	194/244 (80%)	186 (96%)	8 (4%)	27	29
All	All	800/976 (82%)	777 (97%)	23 (3%)	38	42

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	51	ARG
1	A	63	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	79	VAL
1	A	94	LEU
1	A	105	ASN
1	A	111	VAL
1	A	300	LYS
2	B	51	ARG
2	B	79	VAL
2	B	91	ARG
2	B	105	ASN
2	B	111	VAL
3	C	77	GLU
3	C	79	VAL
3	C	103	ARG
3	C	105	ASN
3	C	111	VAL
3	C	211[A]	GLN
3	C	211[B]	GLN
3	C	293	TYR
2	D	79	VAL
2	D	105	ASN
2	D	111	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	105	ASN
2	B	105	ASN
2	B	214	HIS
3	C	105	ASN
2	D	105	ASN
2	D	145	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	AKG	D	1301	4	9,9,9	1.45	3 (33%)	11,11,11	2.62	3 (27%)
5	AKG	B	1302	4	9,9,9	1.07	1 (11%)	11,11,11	2.22	4 (36%)
5	AKG	A	1302	4	9,9,9	1.38	2 (22%)	11,11,11	3.13	6 (54%)
5	AKG	C	1301	4	9,9,9	1.65	2 (22%)	11,11,11	2.55	4 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	AKG	D	1301	4	-	1/9/9/9	-
5	AKG	B	1302	4	-	1/9/9/9	-
5	AKG	A	1302	4	-	0/9/9/9	-
5	AKG	C	1301	4	-	0/9/9/9	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	1301	AKG	C2-C1	3.13	1.58	1.53
5	C	1301	AKG	C3-C2	3.01	1.54	1.51
5	D	1301	AKG	C3-C2	2.62	1.54	1.51
5	A	1302	AKG	C3-C2	2.57	1.54	1.51
5	A	1302	AKG	C2-C1	2.48	1.57	1.53
5	B	1302	AKG	C3-C2	2.23	1.53	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	1301	AKG	C2-C1	2.21	1.56	1.53
5	D	1301	AKG	O2-C1	2.03	1.36	1.30

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1302	AKG	C3-C2-C1	7.32	128.29	115.86
5	D	1301	AKG	C3-C2-C1	6.55	126.99	115.86
5	C	1301	AKG	C3-C2-C1	6.30	126.56	115.86
5	B	1302	AKG	C3-C2-C1	4.90	124.19	115.86
5	A	1302	AKG	O5-C2-C1	-4.69	112.97	119.67
5	D	1301	AKG	O5-C2-C1	-3.96	114.00	119.67
5	C	1301	AKG	O5-C2-C1	-3.22	115.07	119.67
5	A	1302	AKG	O1-C1-C2	2.87	125.37	121.81
5	A	1302	AKG	O4-C5-O3	-2.82	116.09	123.33
5	B	1302	AKG	O5-C2-C1	-2.76	115.72	119.67
5	C	1301	AKG	O2-C1-O1	-2.65	117.60	123.90
5	A	1302	AKG	O2-C1-O1	-2.56	117.81	123.90
5	B	1302	AKG	C4-C3-C2	-2.53	108.15	112.91
5	A	1302	AKG	O4-C5-C4	2.24	121.06	114.00
5	C	1301	AKG	O2-C1-C2	2.19	119.65	113.59
5	B	1302	AKG	O2-C1-C2	2.14	119.54	113.59
5	D	1301	AKG	O2-C1-C2	2.04	119.24	113.59

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	D	1301	AKG	C3-C4-C5-O4
5	B	1302	AKG	C3-C4-C5-O4

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	254/301 (84%)	0.27	18 (7%)	22 23	21, 32, 57, 85	0
2	B	255/301 (84%)	0.32	22 (8%)	16 17	18, 32, 58, 84	1 (0%)
2	D	236/301 (78%)	0.22	12 (5%)	33 35	18, 31, 49, 84	1 (0%)
3	C	238/301 (79%)	0.18	20 (8%)	17 18	18, 31, 51, 84	1 (0%)
All	All	983/1204 (81%)	0.25	72 (7%)	21 22	18, 31, 57, 85	3 (0%)

All (72) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	293	TYR	6.3
2	B	300	LYS	5.1
2	B	98	GLY	4.9
3	C	192	TYR	4.8
2	B	191	VAL	4.7
1	A	85	PRO	4.5
1	A	62	HIS	4.4
2	D	192	TYR	4.3
2	B	62	HIS	4.2
1	A	192	TYR	4.0
3	C	299	ARG	3.9
1	A	191	VAL	3.8
2	B	192	TYR	3.7
2	B	89	GLY	3.6
1	A	80	ALA	3.6
2	B	80	ALA	3.6
3	C	43	GLU	3.5
3	C	77	GLU	3.4
2	B	94	LEU	3.4
2	B	13	LEU	3.3
2	B	224	LEU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	164	ASN	3.2
2	B	214	HIS	3.1
1	A	13	LEU	3.1
2	B	299	ARG	3.0
1	A	165	ARG	2.8
2	D	299	ARG	2.8
2	D	224	LEU	2.8
2	B	85	PRO	2.8
3	C	165	GLU	2.7
3	C	13	LEU	2.7
3	C	279	ARG	2.7
3	C	224	LEU	2.7
3	C	79	VAL	2.7
1	A	89	GLY	2.7
1	A	94	LEU	2.6
2	D	79	VAL	2.6
1	A	224	LEU	2.6
2	D	13	LEU	2.6
2	D	62	HIS	2.5
1	A	299	ARG	2.5
1	A	97	ASP	2.5
2	B	63	LEU	2.5
2	D	66	GLN	2.5
1	A	87	VAL	2.4
3	C	204	ILE	2.4
1	A	103	ARG	2.4
2	D	63	LEU	2.4
3	C	63	LEU	2.4
2	B	88	ASP	2.4
3	C	214	HIS	2.4
2	D	65	ASP	2.3
2	B	96	LEU	2.3
2	B	91	ARG	2.3
3	C	62	HIS	2.3
3	C	145	GLN	2.3
1	A	300	LYS	2.3
3	C	64	ASP	2.2
2	B	291	ASP	2.2
3	C	65	ASP	2.2
1	A	214	HIS	2.2
2	B	92	TYR	2.2
2	D	298	THR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	296	ARG	2.1
3	C	103	ARG	2.1
3	C	294	LEU	2.1
1	A	77	GLU	2.1
2	B	38	ASP	2.1
3	C	61	THR	2.1
2	B	298	THR	2.0
2	B	14	GLU	2.0
2	D	43	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NA	B	1301	1/1	0.74	0.18	37,37,37,37	1
5	AKG	B	1302	10/10	0.85	0.19	34,44,55,56	10
4	NA	A	1301	1/1	0.86	0.11	34,34,34,34	1
5	AKG	C	1301	10/10	0.86	0.13	38,48,59,63	0
5	AKG	D	1301	10/10	0.86	0.16	31,42,53,56	10
5	AKG	A	1302	10/10	0.89	0.17	34,45,53,56	10
4	NA	D	1300	1/1	0.96	0.10	33,33,33,33	1
4	NA	C	1300	1/1	0.97	0.04	31,31,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.