



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 12:53 PM UTC

PDB ID : 1OIL / pdb_00001oil
Title : STRUCTURE OF LIPASE
Authors : Kim, K.K.; Song, H.K.; Shin, D.H.; Suh, S.W.
Deposited on : 1996-12-06
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	320	80% 18% ..
1	B	320	80% 17% ..

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5071 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LIPASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	320	Total	C	N	O	S	0	0	0
			2337	1465	402	467	3			
1	B	320	Total	C	N	O	S	0	0	0
			2337	1465	402	467	3			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	ASP	ALA	conflict	UNP P22088
A	3	ASN	GLY	conflict	UNP P22088
A	18	THR	SER	conflict	UNP P22088
A	40	ARG	ASN	conflict	UNP P22088
A	92	THR	SER	conflict	UNP P22088
A	125	GLY	ASP	conflict	UNP P22088
A	137	THR	SER	conflict	UNP P22088
A	154	ASN	HIS	conflict	UNP P22088
A	165	LYS	GLN	conflict	UNP P22088
A	171	GLN	ARG	conflict	UNP P22088
A	218	ILE	LEU	conflict	UNP P22088
A	232	ILE	LEU	conflict	UNP P22088
A	240	ALA	VAL	conflict	UNP P22088
A	243	PRO	LEU	conflict	UNP P22088
A	256	VAL	ILE	conflict	UNP P22088
A	266	VAL	LEU	conflict	UNP P22088
A	276	GLN	LYS	conflict	UNP P22088
A	300	ASN	TYR	conflict	UNP P22088
B	2	ASP	ALA	conflict	UNP P22088
B	3	ASN	GLY	conflict	UNP P22088
B	18	THR	SER	conflict	UNP P22088
B	40	ARG	ASN	conflict	UNP P22088
B	92	THR	SER	conflict	UNP P22088
B	125	GLY	ASP	conflict	UNP P22088
B	137	THR	SER	conflict	UNP P22088

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	154	ASN	HIS	conflict	UNP P22088
B	165	LYS	GLN	conflict	UNP P22088
B	171	GLN	ARG	conflict	UNP P22088
B	218	ILE	LEU	conflict	UNP P22088
B	232	ILE	LEU	conflict	UNP P22088
B	240	ALA	VAL	conflict	UNP P22088
B	243	PRO	LEU	conflict	UNP P22088
B	256	VAL	ILE	conflict	UNP P22088
B	266	VAL	LEU	conflict	UNP P22088
B	276	GLN	LYS	conflict	UNP P22088
B	300	ASN	TYR	conflict	UNP P22088

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Ca 1 1	0	0
2	B	1	Total Ca 1 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	198	Total O 198 198	0	0
3	B	197	Total O 197 197	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: LIPASE



• Molecule 1: LIPASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	85.23Å 47.42Å 86.53Å 90.00° 116.11° 90.00°	Depositor
Resolution (Å)	8.00 – 2.10 8.00 – 2.10	Depositor EDS
% Data completeness (in resolution range)	79.1 (8.00-2.10) 77.4 (8.00-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 2.05Å)	Xtriage
Refinement program	X-PLOR	Depositor
R, R_{free}	0.180 , 0.255 0.171 , 0.246	Depositor DCC
R_{free} test set	2846 reflections (9.97%)	wwPDB-VP
Wilson B-factor (Å ²)	15.8	Xtriage
Anisotropy	0.467	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 105.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.004 for l,-k,h	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5071	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 92.85 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.4106e-09. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.74	0/2382	1.12	19/3263 (0.6%)
1	B	0.73	0/2382	1.10	15/3263 (0.5%)
All	All	0.73	0/4764	1.11	34/6526 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	232	ILE	N-CA-C	-11.83	98.46	108.63
1	A	292	GLN	N-CA-C	11.46	129.21	109.96
1	B	292	GLN	N-CA-C	10.90	126.66	110.52
1	A	293	LEU	CB-CA-C	7.68	125.71	110.42
1	A	232	ILE	N-CA-C	-7.65	102.05	108.63
1	A	235	VAL	N-CA-C	6.58	121.64	112.50
1	A	177	GLN	N-CA-C	-6.47	103.74	111.69
1	A	292	GLN	CA-C-N	6.44	133.84	121.54
1	A	292	GLN	C-N-CA	6.44	133.84	121.54
1	A	285	ASN	N-CA-C	-6.10	100.90	110.36
1	B	286	HIS	N-CA-C	6.08	118.70	111.71
1	A	44	VAL	N-CA-C	6.01	117.13	108.48
1	A	286	HIS	N-CA-C	5.99	117.81	111.28
1	A	20	THR	N-CA-C	-5.90	99.56	107.88

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	285	ASN	N-CA-C	-5.79	101.38	110.36
1	B	54	SER	N-CA-C	-5.78	103.07	110.53
1	A	293	LEU	N-CA-C	-5.73	98.59	110.80
1	B	60	GLY	N-CA-C	5.69	121.21	112.51
1	B	146	PHE	N-CA-C	-5.59	105.19	111.28
1	A	87	SER	CB-CA-C	-5.58	110.13	116.54
1	B	44	VAL	N-CA-C	5.57	115.91	108.11
1	B	87	SER	CB-CA-C	-5.45	110.27	116.54
1	B	293	LEU	N-CA-C	-5.38	99.35	110.80
1	A	292	GLN	CA-C-O	5.23	126.94	120.92
1	A	24	ALA	N-CA-C	-5.20	106.52	112.86
1	A	249	PHE	N-CA-C	-5.20	105.79	111.82
1	B	292	GLN	CA-C-N	5.19	131.45	121.54
1	B	292	GLN	C-N-CA	5.19	131.45	121.54
1	A	54	SER	N-CA-C	-5.18	103.85	110.53
1	B	293	LEU	CA-CB-CG	-5.15	98.26	116.30
1	B	26	VAL	N-CA-C	5.15	115.69	111.62
1	A	231	THR	N-CA-C	5.11	117.93	109.85
1	A	31	TYR	N-CA-C	5.08	117.53	109.96
1	B	20	THR	N-CA-C	-5.03	100.09	110.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	293	LEU	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2337	0	2292	32	0
1	B	2337	0	2292	35	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	198	0	0	1	0
3	B	197	0	0	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5071	0	4584	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:263:ASN:HD22	1:A:265:GLY:H	1.04	0.97
1:B:263:ASN:HD22	1:B:265:GLY:H	1.12	0.94
1:B:292:GLN:HE21	1:B:293:LEU:H	1.08	0.91
1:A:292:GLN:HE21	1:A:293:LEU:H	1.22	0.87
1:A:15:HIS:HE1	1:A:49:LEU:H	1.22	0.87
1:B:162:ALA:O	1:B:166:THR:HG23	1.79	0.82
1:B:243:PRO:HD2	1:B:293:LEU:HD23	1.62	0.80
1:B:15:HIS:HE1	1:B:49:LEU:H	1.25	0.80
1:A:311:HIS:HD2	1:A:314:ARG:HH11	1.34	0.73
1:A:15:HIS:CE1	1:A:49:LEU:H	2.07	0.73
1:B:54:SER:HA	1:B:166:THR:HG21	1.70	0.72
1:A:67:ALA:O	1:A:71:THR:HG23	1.90	0.71
1:A:263:ASN:HD22	1:A:265:GLY:N	1.84	0.71
1:B:15:HIS:CE1	1:B:49:LEU:H	2.10	0.68
1:B:292:GLN:NE2	1:B:293:LEU:H	1.88	0.68
1:B:67:ALA:O	1:B:71:THR:HG23	1.94	0.67
1:A:34:GLN:NE2	1:A:46:VAL:HG23	2.09	0.67
1:A:263:ASN:ND2	1:A:265:GLY:H	1.87	0.66
1:B:292:GLN:HE21	1:B:293:LEU:N	1.88	0.66
1:B:311:HIS:HD2	1:B:314:ARG:HH11	1.41	0.66
1:B:108:THR:OG1	1:B:311:HIS:HE1	1.85	0.60
1:A:108:THR:OG1	1:A:311:HIS:HE1	1.85	0.60
1:B:263:ASN:ND2	1:B:265:GLY:H	1.92	0.60
1:A:292:GLN:HE21	1:A:293:LEU:N	1.98	0.59
1:B:232:ILE:HG13	1:B:299:ALA:HB3	1.84	0.59
1:A:1:ALA:HB3	1:A:40:ARG:HH21	1.68	0.58
1:A:34:GLN:O	1:A:38:GLN:HG3	2.04	0.58
1:B:28:GLU:HG3	3:B:891:HOH:O	2.06	0.55
1:B:34:GLN:NE2	1:B:46:VAL:HG23	2.21	0.55
1:A:311:HIS:CD2	1:A:314:ARG:HH11	2.21	0.54
1:A:177:GLN:HG2	1:A:178:ASN:N	2.23	0.54
1:A:266:VAL:HG12	1:A:267:VAL:HG23	1.88	0.54
1:A:217:THR:HG21	1:A:227:THR:HG23	1.90	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:PRO:HG2	1:A:81:VAL:HG12	1.90	0.53
1:A:163:ALA:O	1:A:167:LEU:HD23	2.10	0.52
1:B:206:LEU:HD22	1:B:274:TYR:OH	2.10	0.52
1:B:258:ARG:HD3	3:B:784:HOH:O	2.10	0.52
1:B:12:ILE:HD13	1:B:45:TYR:HB2	1.93	0.50
1:B:139:ILE:O	1:B:143:VAL:HG23	2.11	0.50
1:A:255:MET:HB3	1:A:260:SER:OG	2.11	0.49
1:A:219:SER:HA	1:A:223:VAL:O	2.12	0.49
1:A:73:LEU:HD21	1:A:81:VAL:HG13	1.95	0.49
1:A:242:ASP:CG	1:A:293:LEU:HB2	2.38	0.49
1:B:174:THR:O	1:B:177:GLN:HG2	2.13	0.48
1:B:194:ALA:HB1	3:B:750:HOH:O	2.13	0.48
1:B:311:HIS:CD2	1:B:314:ARG:HH11	2.28	0.48
1:B:1:ALA:HB3	1:B:4:TYR:HB2	1.96	0.48
1:A:66:LEU:HD21	3:A:590:HOH:O	2.13	0.48
1:A:215:GLN:HE22	1:A:262:GLN:HE21	1.61	0.47
1:A:45:TYR:HB3	1:A:68:TYR:OH	2.15	0.47
1:B:284:TRP:HB3	1:B:288:ASP:HB2	1.96	0.47
1:B:284:TRP:HD1	1:B:301:ALA:HB1	1.80	0.46
1:B:66:LEU:HD21	3:B:788:HOH:O	2.16	0.46
1:B:211:GLY:O	1:B:264:ASP:HB3	2.16	0.46
1:B:219:SER:HA	1:B:223:VAL:O	2.17	0.45
1:A:284:TRP:HB3	1:A:288:ASP:HB2	1.98	0.44
1:B:27:LEU:HD21	1:B:294:LEU:CD2	2.47	0.44
1:B:297:ARG:HD3	3:B:840:HOH:O	2.17	0.44
1:B:15:HIS:HD2	1:B:16:GLY:O	2.01	0.43
1:A:236:ASP:HA	1:A:237:PRO:HD3	1.89	0.43
1:A:87:SER:HB3	1:A:88:GLN:H	1.60	0.42
1:A:63:GLU:OE2	1:A:95:TYR:OH	2.34	0.42
1:A:293:LEU:HD22	1:A:293:LEU:HA	1.92	0.42
1:B:15:HIS:HE1	1:B:49:LEU:N	2.04	0.41
1:B:24:ALA:HA	3:B:765:HOH:O	2.20	0.41
1:B:213:ALA:HB2	1:B:284:TRP:C	2.46	0.41
1:A:108:THR:OG1	1:A:311:HIS:CE1	2.70	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	318/320 (99%)	311 (98%)	6 (2%)	1 (0%)	36	36
1	B	318/320 (99%)	306 (96%)	11 (4%)	1 (0%)	36	36
All	All	636/640 (99%)	617 (97%)	17 (3%)	2 (0%)	36	36

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	293	LEU
1	B	293	LEU

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	245/245 (100%)	232 (95%)	13 (5%)	20	19
1	B	245/245 (100%)	229 (94%)	16 (6%)	15	13
All	All	490/490 (100%)	461 (94%)	29 (6%)	18	16

All (29) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	87	SER
1	A	129	TYR
1	A	149	LEU
1	A	151	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	152	SER
1	A	177	GLN
1	A	235	VAL
1	A	246	LEU
1	A	256	VAL
1	A	266	VAL
1	A	276	GLN
1	A	292	GLN
1	A	293	LEU
1	B	2	ASP
1	B	26	VAL
1	B	27	LEU
1	B	35	GLU
1	B	71	THR
1	B	87	SER
1	B	99	VAL
1	B	166	THR
1	B	167	LEU
1	B	199	VAL
1	B	220	VAL
1	B	227	THR
1	B	246	LEU
1	B	276	GLN
1	B	292	GLN
1	B	293	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	HIS
1	A	34	GLN
1	A	48	ASN
1	A	88	GLN
1	A	239	ASN
1	A	262	GLN
1	A	263	ASN
1	A	276	GLN
1	A	292	GLN
1	A	300	ASN
1	A	311	HIS
1	B	15	HIS
1	B	34	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	88	GLN
1	B	158	GLN
1	B	177	GLN
1	B	215	GLN
1	B	262	GLN
1	B	263	ASN
1	B	276	GLN
1	B	292	GLN
1	B	300	ASN
1	B	311	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	320/320 (100%)	-0.47	2 (0%) 85 87	5, 14, 30, 44	0
1	B	320/320 (100%)	-0.47	3 (0%) 81 83	5, 14, 29, 49	0
All	All	640/640 (100%)	-0.47	5 (0%) 82 84	5, 14, 29, 49	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1	ALA	4.5
1	A	1	ALA	3.0
1	B	221	PHE	2.3
1	B	142	PHE	2.2
1	A	177	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CA	B	401	1/1	0.93	0.05	11,11,11,11	0
2	CA	A	401	1/1	0.95	0.03	10,10,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.