



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2026 – 12:16 PM EDT

PDB ID : 4OIN / pdb_00004oin
Title : Crystal structure of Thermus thermophilus transcription initiation complex soaked with GE23077
Authors : Zhang, Y.; Ebright, R.H.; Arnold, E.
Deposited on : 2014-01-20
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	% 58% 15% • 27%
1	B	315	2% 58% 14% 28%
2	C	1119	2% 82% 17% ••
3	D	1524	4% 78% 18% ••
4	E	99	% 82% 13% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	F	443	3% 67% 12% 22%
6	G	19	5% 58% 26% 16%
7	H	27	7% 56% 33% 11%
8	I	7	43% 29% 29%

2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 29180 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	231	Total	C	N	O	S	0	0	0
			1809	1155	315	337	2			
1	B	227	Total	C	N	O	S	0	0	0
			1789	1143	310	334	2			

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	1112	Total	C	N	O	S	0	0	0
			8774	5550	1565	1635	24			

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	1485	Total	C	N	O	S	0	1	0
			11739	7442	2069	2193	35			

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	94	Total	C	N	O	S	0	0	0
			758	483	132	139	4			

- Molecule 5 is a protein called DNA directed RNA polymerase sigma factor A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	F	346	Total	C	N	O	S	0	0	0
			2807	1770	509	524	4			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	-19	MET	-	expression tag	UNP Q5SKW1
F	-18	GLY	-	expression tag	UNP Q5SKW1
F	-17	SER	-	expression tag	UNP Q5SKW1
F	-16	SER	-	expression tag	UNP Q5SKW1
F	-15	HIS	-	expression tag	UNP Q5SKW1
F	-14	HIS	-	expression tag	UNP Q5SKW1
F	-13	HIS	-	expression tag	UNP Q5SKW1
F	-12	HIS	-	expression tag	UNP Q5SKW1
F	-11	HIS	-	expression tag	UNP Q5SKW1
F	-10	HIS	-	expression tag	UNP Q5SKW1
F	-9	SER	-	expression tag	UNP Q5SKW1
F	-8	SER	-	expression tag	UNP Q5SKW1
F	-7	GLY	-	expression tag	UNP Q5SKW1
F	-6	LEU	-	expression tag	UNP Q5SKW1
F	-5	VAL	-	expression tag	UNP Q5SKW1
F	-4	PRO	-	expression tag	UNP Q5SKW1
F	-3	ARG	-	expression tag	UNP Q5SKW1
F	-2	GLY	-	expression tag	UNP Q5SKW1
F	-1	SER	-	expression tag	UNP Q5SKW1
F	0	HIS	-	expression tag	UNP Q5SKW1

- Molecule 6 is a DNA chain called 5'-D(*CP*CP*T*GP*CP*AP*TP*CP*CP*GP*TP*GP*AP*GP*TP*CP*GP*AP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	G	16	Total	C	N	O	P	0	0	0
			328	156	63	94	15			

- Molecule 7 is a DNA chain called 5'-D(*TP*AP*TP*AP*AP*TP*GP*GP*GP*AP*GP*C P*TP*GP*TP*CP*AP*CP*GP*GP*AP*TP*GP*CP*AP*GP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	H	24	Total	C	N	O	P	0	0	0
			495	236	94	142	23			

- Molecule 8 is a protein called GE23077.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
8	I	7	Total	C	N	O	0	0	0
			50	26	9	15			


- Molecule 9 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	B	1	Total 1	Mg 1	0	0
9	D	3	Total 3	Mg 3	0	0
9	F	1	Total 1	Mg 1	0	0

- Molecule 10 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	D	2	Total 2	Zn 2	0	0

- Molecule 11 is (2Z)-2-methylbut-2-enoic acid (CCD ID: MB8) (formula: C₅H₈O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	I	1	Total 2	C 1	O 1	0	0

- Molecule 12 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	27	Total 27	O 27	0	0
12	B	25	Total 25	O 25	0	0


Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	C	221	Total 221	O 221	0	0
12	D	269	Total 269	O 269	0	0
12	E	22	Total 22	O 22	0	0
12	F	36	Total 36	O 36	0	0
12	G	10	Total 10	O 10	0	0
12	H	4	Total 4	O 4	0	0
12	I	8	Total 8	O 8	0	0


- Molecule 1: DNA-directed RNA polymerase subunit alpha






• Molecule 3: DNA-directed RNA polymerase subunit beta'










- Molecule 4: DNA-directed RNA polymerase subunit omega



- Molecule 5: DNA directed RNA polymerase sigma factor A



- Molecule 6: 5'-D(*CP*CP*T*GP*CP*AP*TP*CP*CP*GP*TP*GP*AP*GP*TP*CP*GP*AP*G)-3'



- Molecule 7: 5'-D(*TP*AP*TP*AP*AP*TP*GP*GP*GP*AP*GP*CP*TP*GP*TP*CP*AP*CP*GP*GP*AP*TP*GP*CP*AP*GP*G)-3'



- Molecule 8: GE23077



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	183.77Å 103.20Å 294.77Å 90.00° 99.18° 90.00°	Depositor
Resolution (Å)	48.50 – 2.80 48.50 – 2.80	Depositor EDS
% Data completeness (in resolution range)	98.0 (48.50-2.80) 98.0 (48.50-2.80)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.49 (at 2.81Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.206 , 0.252 0.206 , 0.249	Depositor DCC
R_{free} test set	6582 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	47.8	Xtriage
Anisotropy	0.026	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 45.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.019 for $1/2^*h-3/2^*k,-1/2^*h-1/2^*k,-1/2^*h+1/2^*k-l$ 0.017 for $1/2^*h+3/2^*k,1/2^*h-1/2^*k,-1/2^*h-1/2^*k-l$	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	29180	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 0QZ, MB8, FGL, DSN, MG, ZN, R2T, 2RA, 2TL, DVA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	0/1841	0.81	4/2504 (0.2%)
1	B	0.29	0/1821	0.79	0/2476
2	C	0.31	0/8941	0.76	4/12092 (0.0%)
3	D	0.32	0/11948	0.79	12/16153 (0.1%)
4	E	0.29	0/772	0.70	0/1040
5	F	0.29	0/2852	0.74	3/3837 (0.1%)
6	G	0.10	0/368	0.48	0/567
7	H	0.12	0/556	0.54	0/858
All	All	0.31	0/29099	0.76	23/39527 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
8	I	0	1

There are no bond length outliers.

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1208	ASP	N-CA-C	-6.26	99.67	109.25
3	D	433	GLY	N-CA-C	6.14	119.46	110.80
1	A	8	ALA	CA-C-N	6.11	126.62	120.14
1	A	8	ALA	C-N-CA	6.11	126.62	120.14
2	C	169	GLY	CA-C-N	6.04	125.94	119.85

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
8	I	5	2TL	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1809	0	1863	29	0
1	B	1789	0	1841	27	0
2	C	8774	0	8877	115	0
3	D	11739	0	11977	172	0
4	E	758	0	770	9	0
5	F	2807	0	2882	36	0
6	G	328	0	181	5	0
7	H	495	0	272	9	0
8	I	50	0	37	3	0
9	B	1	0	0	0	0
9	D	3	0	0	0	0
9	F	1	0	0	0	0
10	D	2	0	0	0	0
11	I	2	0	0	0	0
12	A	27	0	0	0	0
12	B	25	0	0	0	0
12	C	221	0	0	5	0
12	D	269	0	0	7	0
12	E	22	0	0	0	0
12	F	36	0	0	0	0
12	G	10	0	0	1	0
12	H	4	0	0	2	0
12	I	8	0	0	0	0
All	All	29180	0	28700	360	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 360 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:802:ARG:HB2	2:C:826:TYR:HB2	1.62	0.81
3:D:1254:GLN:HB3	3:D:1258:ARG:HB2	1.66	0.75
2:C:428:ARG:NH2	2:C:447:ALA:O	2.20	0.75
2:C:409:ARG:HH11	2:C:454:SER:HB2	1.53	0.74
2:C:758:ARG:HH21	2:C:788:THR:HB	1.53	0.74

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	229/315 (73%)	226 (99%)	3 (1%)	0	100	100
1	B	225/315 (71%)	222 (99%)	3 (1%)	0	100	100
2	C	1108/1119 (99%)	1088 (98%)	20 (2%)	0	100	100
3	D	1482/1524 (97%)	1451 (98%)	31 (2%)	0	100	100
4	E	92/99 (93%)	89 (97%)	3 (3%)	0	100	100
5	F	344/443 (78%)	340 (99%)	4 (1%)	0	100	100
All	All	3480/3815 (91%)	3416 (98%)	64 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/273 (73%)	194 (97%)	6 (3%)	36	72
1	B	200/273 (73%)	196 (98%)	4 (2%)	48	80
2	C	936/941 (100%)	912 (97%)	24 (3%)	40	75
3	D	1253/1279 (98%)	1223 (98%)	30 (2%)	43	77
4	E	82/88 (93%)	81 (99%)	1 (1%)	63	87
5	F	301/388 (78%)	295 (98%)	6 (2%)	48	80
All	All	2972/3242 (92%)	2901 (98%)	71 (2%)	43	77

5 of 71 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	D	1305	LEU
3	D	1433	SER
5	F	95	THR
2	C	557	ARG
2	C	513	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
3	D	994	GLN
3	D	1195	GLN
3	D	1124	GLN
3	D	1202	GLN
2	C	1047	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

7 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	2TL	I	5	8	5,6,7	1.08	0	5,7,9	1.00	1 (20%)
8	R2T	I	4	8	8,10,11	1.96	2 (25%)	6,13,15	0.91	0
8	2RA	I	1	11,8	4,5,6	0.55	0	1,5,7	0.39	0
8	FGL	I	7	8	5,6,7	1.05	0	1,7,9	1.98	0
8	0QZ	I	6	8	4,5,6	1.72	1 (25%)	2,5,7	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	2TL	I	5	8	-	1/5/6/8	-
8	R2T	I	4	8	-	7/13/14/16	-
8	2RA	I	1	11,8	-	0/2/4/6	-
8	FGL	I	7	8	-	0/4/6/8	-
8	0QZ	I	6	8	-	1/3/4/6	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	I	4	R2T	CD-NE2	4.50	1.43	1.32
8	I	6	0QZ	OB-CA	-3.34	1.37	1.43
8	I	4	R2T	OB1-CB	-2.21	1.37	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	I	5	2TL	O-C-CA	-2.07	119.45	124.77

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	I	4	R2T	CA-CB-CG-OG1
8	I	4	R2T	OB1-CB-CG-CD
8	I	4	R2T	OB1-CB-CG-OG1
8	I	4	R2T	OE1-CD-CG-OG1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
8	I	4	R2T	NE2-CD-CG-OG1

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	I	4	R2T	2	0
8	I	7	FGL	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 7 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
11	MB8	I	101	8	0,1,6	-	-	-		

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	231/315 (73%)	-0.14	2 (0%) 81 74	16, 33, 61, 102	0
1	B	227/315 (72%)	0.17	5 (2%) 62 52	21, 43, 78, 108	0
2	C	1112/1119 (99%)	-0.18	18 (1%) 70 61	4, 25, 84, 113	0
3	D	1485/1524 (97%)	0.10	67 (4%) 38 30	2, 33, 94, 121	1 (0%)
4	E	94/99 (94%)	-0.32	1 (1%) 78 70	7, 28, 63, 73	0
5	F	346/443 (78%)	0.34	13 (3%) 44 35	14, 47, 87, 113	0
6	G	16/19 (84%)	0.71	1 (6%) 26 19	44, 77, 145, 154	0
7	H	24/27 (88%)	0.65	2 (8%) 17 12	43, 83, 136, 156	0
8	I	0/7	-	-	-	-
All	All	3535/3868 (91%)	0.02	109 (3%) 51 41	2, 34, 89, 156	1 (0%)

The worst 5 of 109 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	449	SER	4.2
3	D	447	VAL	4.2
5	F	141	VAL	4.2
3	D	420	VAL	4.1
1	B	6	LEU	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	2RA	I	1	6/7	0.91	0.10	8,10,16,18	0
8	DVA	I	3	7/8	0.95	0.09	8,9,13,13	0
8	FGL	I	7	7/8	0.96	0.05	8,9,10,11	0
8	R2T	I	4	11/12	0.96	0.07	7,8,9,11	0
8	2TL	I	5	7/8	0.96	0.07	6,7,8,9	0
8	DSN	I	2	6/7	0.97	0.07	7,9,15,15	0
8	0QZ	I	6	6/7	0.98	0.07	7,7,10,11	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
9	MG	B	2001	1/1	0.80	0.11	50,50,50,50	0
11	MB8	I	101	2/7	0.92	0.08	6,6,6,11	0
9	MG	D	2005	1/1	0.93	0.11	31,31,31,31	0
9	MG	F	2001	1/1	0.97	0.04	22,22,22,22	0
10	ZN	D	2002	1/1	0.99	0.03	56,56,56,56	0
9	MG	D	2004	1/1	0.99	0.02	31,31,31,31	0
9	MG	D	2003	1/1	1.00	0.02	5,5,5,5	0
10	ZN	D	2001	1/1	1.00	0.01	10,10,10,10	0

6.5 Other polymers [i](#)

There are no such residues in this entry.