



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:58 PM UTC

PDB ID : 8OIK / pdb_00008oik
Title : D-PHAT domain (NTD) of human SAMD4A
Authors : Kubikova, J.; Jeske, M.
Deposited on : 2023-03-23
Resolution : 1.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6728 (1.64-1.60)
Clashscore	190562	7023 (1.64-1.60)
Ramachandran outliers	187476	6898 (1.64-1.60)
Sidechain outliers	187428	6896 (1.64-1.60)
RSRZ outliers	180081	6727 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	180	11% 78% 14% 8%
1	B	180	21% 79% 9% 11%
1	C	180	22% 76% 9% • 14%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4174 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein Smaug homolog 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	165	Total	C	N	O	S	0	0	0
			1329	846	232	245	6			
1	B	160	Total	C	N	O	S	0	0	0
			1260	804	221	229	6			
1	C	154	Total	C	N	O	S	0	0	0
			1215	777	212	220	6			

There are 75 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-23	MET	-	initiating methionine	UNP Q9UPU9
A	-22	LYS	-	expression tag	UNP Q9UPU9
A	-21	HIS	-	expression tag	UNP Q9UPU9
A	-20	HIS	-	expression tag	UNP Q9UPU9
A	-19	HIS	-	expression tag	UNP Q9UPU9
A	-18	HIS	-	expression tag	UNP Q9UPU9
A	-17	HIS	-	expression tag	UNP Q9UPU9
A	-16	HIS	-	expression tag	UNP Q9UPU9
A	-15	PRO	-	expression tag	UNP Q9UPU9
A	-14	MET	-	expression tag	UNP Q9UPU9
A	-13	SER	-	expression tag	UNP Q9UPU9
A	-12	ASP	-	expression tag	UNP Q9UPU9
A	-11	TYR	-	expression tag	UNP Q9UPU9
A	-10	ASP	-	expression tag	UNP Q9UPU9
A	-9	ILE	-	expression tag	UNP Q9UPU9
A	-8	PRO	-	expression tag	UNP Q9UPU9
A	-7	THR	-	expression tag	UNP Q9UPU9
A	-6	THR	-	expression tag	UNP Q9UPU9
A	-5	GLU	-	expression tag	UNP Q9UPU9
A	-4	ASN	-	expression tag	UNP Q9UPU9
A	-3	LEU	-	expression tag	UNP Q9UPU9
A	-2	TYR	-	expression tag	UNP Q9UPU9
A	-1	PHE	-	expression tag	UNP Q9UPU9

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLN	-	expression tag	UNP Q9UPU9
A	1	SER	-	expression tag	UNP Q9UPU9
B	-23	MET	-	initiating methionine	UNP Q9UPU9
B	-22	LYS	-	expression tag	UNP Q9UPU9
B	-21	HIS	-	expression tag	UNP Q9UPU9
B	-20	HIS	-	expression tag	UNP Q9UPU9
B	-19	HIS	-	expression tag	UNP Q9UPU9
B	-18	HIS	-	expression tag	UNP Q9UPU9
B	-17	HIS	-	expression tag	UNP Q9UPU9
B	-16	HIS	-	expression tag	UNP Q9UPU9
B	-15	PRO	-	expression tag	UNP Q9UPU9
B	-14	MET	-	expression tag	UNP Q9UPU9
B	-13	SER	-	expression tag	UNP Q9UPU9
B	-12	ASP	-	expression tag	UNP Q9UPU9
B	-11	TYR	-	expression tag	UNP Q9UPU9
B	-10	ASP	-	expression tag	UNP Q9UPU9
B	-9	ILE	-	expression tag	UNP Q9UPU9
B	-8	PRO	-	expression tag	UNP Q9UPU9
B	-7	THR	-	expression tag	UNP Q9UPU9
B	-6	THR	-	expression tag	UNP Q9UPU9
B	-5	GLU	-	expression tag	UNP Q9UPU9
B	-4	ASN	-	expression tag	UNP Q9UPU9
B	-3	LEU	-	expression tag	UNP Q9UPU9
B	-2	TYR	-	expression tag	UNP Q9UPU9
B	-1	PHE	-	expression tag	UNP Q9UPU9
B	0	GLN	-	expression tag	UNP Q9UPU9
B	1	SER	-	expression tag	UNP Q9UPU9
C	-23	MET	-	initiating methionine	UNP Q9UPU9
C	-22	LYS	-	expression tag	UNP Q9UPU9
C	-21	HIS	-	expression tag	UNP Q9UPU9
C	-20	HIS	-	expression tag	UNP Q9UPU9
C	-19	HIS	-	expression tag	UNP Q9UPU9
C	-18	HIS	-	expression tag	UNP Q9UPU9
C	-17	HIS	-	expression tag	UNP Q9UPU9
C	-16	HIS	-	expression tag	UNP Q9UPU9
C	-15	PRO	-	expression tag	UNP Q9UPU9
C	-14	MET	-	expression tag	UNP Q9UPU9
C	-13	SER	-	expression tag	UNP Q9UPU9
C	-12	ASP	-	expression tag	UNP Q9UPU9
C	-11	TYR	-	expression tag	UNP Q9UPU9
C	-10	ASP	-	expression tag	UNP Q9UPU9
C	-9	ILE	-	expression tag	UNP Q9UPU9

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	-8	PRO	-	expression tag	UNP Q9UPU9
C	-7	THR	-	expression tag	UNP Q9UPU9
C	-6	THR	-	expression tag	UNP Q9UPU9
C	-5	GLU	-	expression tag	UNP Q9UPU9
C	-4	ASN	-	expression tag	UNP Q9UPU9
C	-3	LEU	-	expression tag	UNP Q9UPU9
C	-2	TYR	-	expression tag	UNP Q9UPU9
C	-1	PHE	-	expression tag	UNP Q9UPU9
C	0	GLN	-	expression tag	UNP Q9UPU9
C	1	SER	-	expression tag	UNP Q9UPU9

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			7	4	3		
3	C	1	Total	C	O	0	0
			7	4	3		
3	C	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	132	Total	O	0	0
			132	132		
4	B	116	Total	O	0	0
			116	116		
4	C	97	Total	O	0	0
			97	97		

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	81.83Å 142.79Å 137.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.45 – 1.62 49.45 – 1.62	Depositor EDS
% Data completeness (in resolution range)	87.4 (49.45-1.62) 88.1 (49.45-1.62)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.03 (at 1.62Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.207 , 0.221 0.208 , 0.221	Depositor DCC
R_{free} test set	2003 reflections (1.95%)	wwPDB-VP
Wilson B-factor (Å ²)	33.5	Xtriage
Anisotropy	0.012	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 48.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.008 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.014 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4174	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/1357	0.59	0/1841
1	B	0.44	0/1286	0.61	0/1745
1	C	0.43	0/1239	0.61	0/1680
All	All	0.43	0/3882	0.60	0/5266

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1329	0	1334	22	0
1	B	1260	0	1247	9	0
1	C	1215	0	1222	18	0
2	B	4	0	6	0	0
3	B	7	0	10	0	0
3	C	14	0	20	3	0
4	A	132	0	0	1	0
4	B	116	0	0	3	0
4	C	97	0	0	0	0
All	All	4174	0	3839	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:-5:GLU:HG2	1:C:123:ARG:HH21	1.34	0.92
1:A:-5:GLU:HG2	1:C:123:ARG:NH2	1.87	0.90
1:A:-8:PRO:HB2	1:C:123:ARG:HD2	1.51	0.89
1:A:120:GLU:OE1	1:A:123:ARG:NH1	2.21	0.74
1:A:105:LEU:O	1:A:109:LEU:HG	1.92	0.69
1:A:-8:PRO:HB2	1:C:123:ARG:CD	2.27	0.64
1:B:79:VAL:HG11	1:B:108:ILE:HD11	1.83	0.61
1:B:109:LEU:HD11	1:B:126:LEU:HD21	1.85	0.57
1:A:29:LEU:HD21	1:C:10:LEU:HG	1.87	0.55
1:A:48:SER:CB	3:C:201:PEG:H32	2.36	0.55
1:B:-7:THR:HG23	1:B:-3:LEU:HD23	1.89	0.55
1:A:92:PRO:HA	1:A:134:ALA:HB2	1.90	0.53
1:C:45:LEU:HD23	3:C:202:PEG:H32	1.93	0.51
1:C:115:HIS:O	1:C:116:ASN:C	2.53	0.50
1:C:45:LEU:HD23	3:C:202:PEG:C3	2.42	0.49
1:A:109:LEU:HD11	1:A:126:LEU:HD21	1.94	0.49
1:B:92:PRO:HG3	1:B:133:PRO:HG2	1.96	0.47
1:A:59:GLU:CD	1:C:39:ARG:HE	2.23	0.46
1:A:120:GLU:CD	1:A:123:ARG:NH1	2.74	0.46
1:B:70:GLN:NE2	4:B:301:HOH:O	2.23	0.45
1:A:150:LEU:O	1:A:154:THR:HG23	2.16	0.45
1:A:-5:GLU:CG	1:C:123:ARG:HH21	2.19	0.45
1:C:111:HIS:C	1:C:111:HIS:CD2	2.95	0.44
1:B:84:LEU:HD11	1:B:121:GLU:HB3	1.99	0.44
4:B:314:HOH:O	1:C:56:HIS:HD2	1.99	0.44
1:A:-2:TYR:O	1:A:2:MET:HG3	2.17	0.43
1:C:80:ILE:HG13	1:C:108:ILE:HD12	2.00	0.43
1:C:117:GLN:O	1:C:118:HIS:C	2.60	0.43
1:B:92:PRO:HA	1:B:134:ALA:HB2	2.01	0.43
4:A:224:HOH:O	1:C:-1:PHE:HD1	2.01	0.43
1:A:-4:ASN:HB3	1:A:-1:PHE:HB2	2.01	0.42
1:A:15:LYS:HE2	1:C:32:ARG:NH1	2.35	0.41
1:A:-7:THR:HB	1:C:21:GLU:HG3	2.01	0.41
1:A:91:LYS:HA	1:A:92:PRO:HD3	1.96	0.41
1:B:15:LYS:NZ	4:B:305:HOH:O	2.42	0.41
1:B:71:TRP:CD1	1:B:82:LEU:HD13	2.55	0.41
1:A:4:ARG:HH11	1:A:4:ARG:HG2	1.86	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:-8:PRO:HG3	1:C:147:LEU:HD21	2.03	0.41
1:A:30:LEU:O	1:A:33:VAL:HG22	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	163/180 (91%)	161 (99%)	2 (1%)	0	100	100
1	B	158/180 (88%)	156 (99%)	2 (1%)	0	100	100
1	C	152/180 (84%)	149 (98%)	2 (1%)	1 (1%)	18	6
All	All	473/540 (88%)	466 (98%)	6 (1%)	1 (0%)	43	25

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	118	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	147/163 (90%)	146 (99%)	1 (1%)	76	62
1	B	134/163 (82%)	132 (98%)	2 (2%)	57	34

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	131/163 (80%)	129 (98%)	2 (2%)	57	34
All	All	412/489 (84%)	407 (99%)	5 (1%)	63	43

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-6	THR
1	B	107	LYS
1	B	150	LEU
1	C	76	LYS
1	C	118	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	115	HIS
1	A	117	GLN
1	A	148	ASN
1	B	0	GLN
1	B	69	ASN
1	B	72	GLN
1	B	111	HIS
1	B	116	ASN
1	C	35	GLN
1	C	72	GLN
1	C	73	GLN
1	C	116	ASN
1	C	118	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	B	201	-	3,3,3	0.10	0	2,2,2	0.07	0
3	PEG	C	202	-	6,6,6	0.11	0	5,5,5	0.06	0
3	PEG	C	201	-	6,6,6	0.12	0	5,5,5	0.12	0
3	PEG	B	202	-	6,6,6	0.09	0	5,5,5	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	B	201	-	-	1/1/1/1	-
3	PEG	C	202	-	-	3/4/4/4	-
3	PEG	C	201	-	-	3/4/4/4	-
3	PEG	B	202	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	202	PEG	O1-C1-C2-O2
3	C	201	PEG	O1-C1-C2-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	C	201	PEG	O2-C3-C4-O4
3	B	202	PEG	O2-C3-C4-O4
3	C	201	PEG	C4-C3-O2-C2
3	C	202	PEG	O1-C1-C2-O2
2	B	201	EDO	O1-C1-C2-O2
3	C	202	PEG	C4-C3-O2-C2
3	B	202	PEG	C4-C3-O2-C2
3	C	202	PEG	C1-C2-O2-C3

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	202	PEG	2	0
3	C	201	PEG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	165/180 (91%)	0.71	20 (12%) 8 9	25, 34, 51, 72	0
1	B	160/180 (88%)	1.22	38 (23%) 2 2	27, 37, 77, 84	0
1	C	154/180 (85%)	1.26	40 (25%) 1 1	26, 38, 74, 90	0
All	All	479/540 (88%)	1.06	98 (20%) 2 3	25, 35, 72, 90	0

All (98) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	-6	THR	6.9
1	B	146	TRP	6.8
1	B	-9	ILE	6.7
1	C	113	ILE	6.1
1	C	-1	PHE	6.0
1	A	-7	THR	5.9
1	C	115	HIS	5.9
1	A	-9	ILE	5.8
1	B	-8	PRO	5.6
1	B	149	HIS	5.5
1	B	113	ILE	5.5
1	C	119	ILE	5.5
1	C	-2	TYR	5.1
1	C	146	TRP	4.8
1	B	-3	LEU	4.8
1	A	-8	PRO	4.8
1	B	150	LEU	4.7
1	C	114	GLU	4.5
1	C	76	LYS	4.5
1	A	-6	THR	4.3
1	B	-4	ASN	4.2
1	C	150	LEU	4.2
1	A	155	SER	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	109	LEU	3.9
1	B	137	LEU	3.9
1	B	-7	THR	3.9
1	C	116	ASN	3.8
1	C	111	HIS	3.8
1	B	-5	GLU	3.7
1	C	149	HIS	3.6
1	C	145	MET	3.6
1	C	147	LEU	3.6
1	B	-1	PHE	3.6
1	B	115	HIS	3.5
1	C	117	GLN	3.5
1	B	114	GLU	3.5
1	B	119	ILE	3.5
1	C	77	ASP	3.4
1	B	108	ILE	3.4
1	A	128	TYR	3.3
1	C	124	GLN	3.3
1	A	152	ASP	3.3
1	B	72	GLN	3.2
1	C	-3	LEU	3.2
1	B	147	LEU	3.1
1	C	137	LEU	3.1
1	B	76	LYS	3.1
1	B	145	MET	3.0
1	B	148	ASN	3.0
1	A	-1	PHE	2.9
1	C	0	GLN	2.9
1	B	118	HIS	2.9
1	C	120	GLU	2.9
1	C	118	HIS	2.9
1	C	138	GLU	2.9
1	C	80	ILE	2.8
1	C	141	SER	2.8
1	A	117	GLN	2.8
1	B	128	TYR	2.7
1	C	73	GLN	2.7
1	B	0	GLN	2.7
1	C	110	ALA	2.7
1	A	-5	GLU	2.6
1	B	112	SER	2.6
1	A	154	THR	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	108	ILE	2.6
1	A	148	ASN	2.6
1	C	126	LEU	2.6
1	B	144	ALA	2.5
1	B	80	ILE	2.4
1	C	75	SER	2.4
1	A	0	GLN	2.4
1	B	142	ALA	2.4
1	B	111	HIS	2.4
1	B	104	LEU	2.3
1	C	15	LYS	2.3
1	A	92	PRO	2.3
1	A	-3	LEU	2.3
1	B	116	ASN	2.3
1	B	122	SER	2.3
1	A	73	GLN	2.2
1	B	73	GLN	2.2
1	C	72	GLN	2.2
1	A	10	LEU	2.2
1	C	148	ASN	2.2
1	C	112	SER	2.2
1	B	77	ASP	2.1
1	C	70	GLN	2.1
1	C	144	ALA	2.1
1	B	143	LEU	2.1
1	C	128	TYR	2.1
1	B	123	ARG	2.1
1	C	4	ARG	2.1
1	C	84	LEU	2.1
1	C	79	VAL	2.0
1	A	137	LEU	2.0
1	A	31	LYS	2.0
1	A	93	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	PEG	B	202	7/7	0.69	0.24	44,45,54,57	0
3	PEG	C	201	7/7	0.83	0.18	49,50,55,55	0
3	PEG	C	202	7/7	0.83	0.19	46,47,53,57	0
2	EDO	B	201	4/4	0.85	0.17	49,54,54,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.