



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:06 PM UTC

PDB ID : 1OJ9 / pdb_00001oj9
Title : HUMAN MONOAMINE OXIDASE B IN COMPLEX WITH 1,4-DIPHENYL-2-BUTENE
Authors : Binda, C.; Edmondson, D.E.; Mattevi, A.
Deposited on : 2003-07-08
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	520	11% 88% 7% . .
1	B	520	9% 87% 7% . 5%

2 Entry composition ⓘ


There are 4 unique types of molecules in this entry. The entry contains 8279 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AMINE OXIDASE [FLAVIN-CONTAINING] B.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	499	Total	C	N	O	S	0	0	0
			3971	2538	681	728	24			
1	B	494	Total	C	N	O	S	0	0	0
			3940	2519	676	721	24			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: C₂₇H₃₃N₉O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is 1,4-DIPHENYL-2-BUTENE (CCD ID: 1PB) (formula: C₁₆H₁₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C 16 16	0	0
3	B	1	Total C 16 16	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	100	Total O 100 100	0	0
4	B	130	Total O 130 130	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: AMINE OXIDASE [FLAVIN-CONTAINING] B



• Molecule 1: AMINE OXIDASE [FLAVIN-CONTAINING] B



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	132.10Å 224.02Å 86.83Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 2.30 40.00 – 2.30	Depositor EDS
% Data completeness (in resolution range)	90.6 (40.00-2.30) 90.2 (40.00-2.30)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.88 (at 2.29Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.212 , 0.253 0.206 , 0.243	Depositor DCC
R_{free} test set	1333 reflections (2.57%)	wwPDB-VP
Wilson B-factor (Å ²)	19.8	Xtriage
Anisotropy	0.583	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 52.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.003 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.015 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8279	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 1PB, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/4068	0.83	0/5522
1	B	0.53	1/4037 (0.0%)	0.83	0/5479
All	All	0.53	1/8105 (0.0%)	0.83	0/11001

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	489	VAL	CA-CB	5.99	1.57	1.54

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3971	0	3967	23	0
1	B	3940	0	3937	25	0
2	A	53	0	29	0	0
2	B	53	0	29	0	0
3	A	16	0	16	0	0
3	B	16	0	16	0	0
4	A	100	0	0	1	0
4	B	130	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	8279	0	7994	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:28:LEU:HD11	1:B:454:MET:HE1	1.50	0.91
1:A:28:LEU:HD11	1:A:454:MET:HE1	1.51	0.91
1:A:353:ALA:HB1	1:B:280:MET:HE1	1.57	0.83
1:A:280:MET:HE1	1:B:353:ALA:HB1	1.63	0.80
1:B:117:ASN:HD22	1:B:120:ARG:HH21	1.31	0.77
1:A:117:ASN:HD22	1:A:120:ARG:HH21	1.37	0.71
1:A:3:ASN:HB3	1:A:4:LYS:HG2	1.89	0.55
1:B:3:ASN:HB3	1:B:4:LYS:HG2	1.90	0.54
1:A:389:CYS:HB2	1:B:280:MET:HG3	1.93	0.50
1:B:117:ASN:HD22	1:B:120:ARG:NH2	2.04	0.50
1:A:270:MET:HE1	1:B:268:LEU:HD23	1.93	0.50
1:B:445:ARG:HD2	1:B:463:TRP:CZ2	2.48	0.48
1:A:91:HIS:HE1	4:A:2067:HOH:O	1.96	0.48
1:A:353:ALA:CB	1:B:280:MET:HE1	2.35	0.48
1:A:268:LEU:HD23	1:B:270:MET:HE1	1.96	0.47
1:A:445:ARG:HD2	1:A:463:TRP:CZ2	2.50	0.47
1:B:278:LEU:HD13	1:B:282:ARG:HG2	1.98	0.46
1:B:280:MET:HB3	1:B:280:MET:HE3	1.70	0.46
1:A:117:ASN:HD22	1:A:120:ARG:NH2	2.09	0.45
1:A:278:LEU:HD13	1:A:282:ARG:HG2	1.98	0.45
1:A:116:ASN:HD21	1:A:483:GLU:HG2	1.82	0.45
1:B:308:LYS:HB2	1:B:308:LYS:HE2	1.85	0.45
1:A:445:ARG:HD2	1:A:463:TRP:CH2	2.52	0.44
1:B:28:LEU:HD21	1:B:456:LYS:HE3	1.99	0.44
1:A:28:LEU:HD21	1:A:456:LYS:HE3	2.00	0.44
1:B:281:MET:HB3	1:B:413:VAL:HG21	2.00	0.43
1:A:108:ASN:HA	1:A:109:PRO:HD3	1.89	0.43
1:A:280:MET:HG3	1:B:389:CYS:HB2	2.01	0.43
1:A:249:THR:CG2	1:A:253:GLU:HB2	2.49	0.43
1:B:445:ARG:HD2	1:B:463:TRP:CH2	2.53	0.42
1:A:489:VAL:N	1:A:490:PRO:HD2	2.34	0.42
1:A:23:LEU:HD22	1:A:28:LEU:HD12	2.02	0.42
1:B:55:ASP:O	1:B:296:LYS:HE2	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:416:GLN:HA	1:B:417:PRO:HD3	1.93	0.41
1:A:237:TYR:HB3	1:A:248:GLU:HB3	2.02	0.41
1:B:237:TYR:HB3	1:B:248:GLU:HB3	2.02	0.41
1:B:489:VAL:N	1:B:490:PRO:HD2	2.35	0.41
1:B:108:ASN:HA	1:B:109:PRO:HD3	1.87	0.41
1:B:238:ILE:HG21	1:B:421:ILE:HD11	2.02	0.41
1:A:280:MET:HB3	1:A:280:MET:HE3	1.73	0.41
1:B:249:THR:CG2	1:B:253:GLU:HB2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	497/520 (96%)	483 (97%)	14 (3%)	0	100	100
1	B	492/520 (95%)	481 (98%)	11 (2%)	0	100	100
All	All	989/1040 (95%)	964 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	427/444 (96%)	410 (96%)	17 (4%)	28	42

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	424/444 (96%)	408 (96%)	16 (4%)	29	44
All	All	851/888 (96%)	818 (96%)	33 (4%)	28	43

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	A	49	GLN
1	A	149	LYS
1	A	209	LYS
1	A	242	ARG
1	A	249	THR
1	A	280	MET
1	A	308	LYS
1	A	318	ASP
1	A	321	GLU
1	A	332	LYS
1	A	379	GLU
1	A	397	CYS
1	A	445	ARG
1	A	464	GLN
1	A	495	LEU
1	A	498	LEU
1	B	3	ASN
1	B	49	GLN
1	B	149	LYS
1	B	209	LYS
1	B	242	ARG
1	B	249	THR
1	B	280	MET
1	B	308	LYS
1	B	318	ASP
1	B	321	GLU
1	B	332	LYS
1	B	379	GLU
1	B	397	CYS
1	B	445	ARG
1	B	464	GLN
1	B	495	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	116	ASN
1	A	117	ASN
1	A	431	HIS
1	B	116	ASN
1	B	117	ASN
1	B	431	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	1PB	A	1503	-	17,17,17	1.12	2 (11%)	20,20,20	1.30	3 (15%)
2	FAD	A	1502	1	58,58,58	1.11	5 (8%)	85,89,89	1.61	14 (16%)
2	FAD	B	1497	1	58,58,58	1.11	5 (8%)	85,89,89	1.58	14 (16%)
3	1PB	B	1498	-	17,17,17	1.14	2 (11%)	20,20,20	1.27	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	1PB	A	1503	-	-	5/7/7/7	0/2/2/2
2	FAD	A	1502	1	-	0/34/50/50	0/6/6/6
2	FAD	B	1497	1	-	0/34/50/50	0/6/6/6
3	1PB	B	1498	-	-	5/7/7/7	0/2/2/2

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1502	FAD	C4X-N5	3.59	1.38	1.30
2	B	1497	FAD	C4X-N5	3.55	1.38	1.30
2	A	1502	FAD	C10-N1	3.06	1.39	1.33
2	B	1497	FAD	C10-N1	3.05	1.39	1.33
2	B	1497	FAD	C2A-N3A	2.72	1.38	1.33
3	B	1498	1PB	CD-CC	-2.70	1.39	1.52
2	A	1502	FAD	C2A-N1A	2.68	1.38	1.33
3	A	1503	1PB	CD-CC	-2.68	1.39	1.52
3	A	1503	1PB	CA-CB	2.59	1.39	1.31
3	B	1498	1PB	CA-CB	2.58	1.39	1.31
2	B	1497	FAD	C2A-N1A	2.53	1.38	1.33
2	A	1502	FAD	C2A-N3A	2.37	1.38	1.33
2	A	1502	FAD	C8A-N7A	2.31	1.36	1.31
2	B	1497	FAD	C8A-N7A	2.17	1.35	1.31

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1502	FAD	N3A-C2A-N1A	-5.83	119.75	128.58
2	B	1497	FAD	N3A-C2A-N1A	-5.80	119.80	128.58
2	B	1497	FAD	C5A-C4A-N3A	-4.15	121.00	126.72
2	A	1502	FAD	C5A-C4A-N3A	-3.98	121.24	126.72
2	A	1502	FAD	C4X-C10-N1	-3.90	115.02	124.59
2	B	1497	FAD	N9A-C8A-N7A	-3.81	108.54	113.94
2	A	1502	FAD	N9A-C8A-N7A	-3.74	108.63	113.94
2	A	1502	FAD	O4-C4-C4X	-3.67	116.84	126.53
2	B	1497	FAD	C4X-C10-N1	-3.64	115.67	124.59
3	A	1503	1PB	CC-CD-C1	3.58	121.89	113.21
3	B	1498	1PB	CC-CD-C1	3.57	121.85	113.21
2	B	1497	FAD	O4-C4-C4X	-3.45	117.42	126.53
2	A	1502	FAD	C9A-C5X-N5	-3.42	118.82	122.45
2	B	1497	FAD	C5A-N7A-C8A	3.35	108.72	103.45
2	B	1497	FAD	C2A-N3A-C4A	3.30	119.89	111.83

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1502	FAD	C2A-N3A-C4A	3.30	119.89	111.83
2	A	1502	FAD	C5A-N7A-C8A	3.28	108.60	103.45
2	B	1497	FAD	C9A-C5X-N5	-3.16	119.10	122.45
2	A	1502	FAD	C4-N3-C2	-3.12	120.11	125.64
2	B	1497	FAD	C4-N3-C2	-3.01	120.30	125.64
2	A	1502	FAD	C10-C4X-N5	-2.87	118.95	124.81
2	B	1497	FAD	C10-C4X-N5	-2.81	119.06	124.81
3	A	1503	1PB	CC-CB-CA	-2.75	119.39	125.48
3	B	1498	1PB	CC-CB-CA	-2.72	119.45	125.48
2	B	1497	FAD	N3A-C4A-N9A	2.71	131.77	127.17
2	A	1502	FAD	N3A-C4A-N9A	2.44	131.31	127.17
3	A	1503	1PB	CG-CA-CB	-2.41	121.08	126.99
2	A	1502	FAD	C4A-C5A-N7A	-2.40	107.83	110.58
2	B	1497	FAD	C4A-C5A-N7A	-2.40	107.84	110.58
2	B	1497	FAD	C4X-C4-N3	2.30	119.10	113.25
2	A	1502	FAD	C4X-C4-N3	2.27	119.03	113.25
3	B	1498	1PB	CG-CA-CB	-2.23	121.53	126.99
2	A	1502	FAD	C4-C4X-C10	2.05	120.46	116.93
2	B	1497	FAD	C5'-C4'-C3'	-2.05	108.34	112.22

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	1498	1PB	CB-CA-CG-CL
3	B	1498	1PB	CB-CA-CG-CH
3	A	1503	1PB	CB-CA-CG-CL
3	A	1503	1PB	CB-CA-CG-CH
3	B	1498	1PB	CA-CB-CC-CD
3	A	1503	1PB	CA-CB-CC-CD
3	A	1503	1PB	CN-C1-CD-CC
3	B	1498	1PB	CN-C1-CD-CC
3	A	1503	1PB	CR-C1-CD-CC
3	B	1498	1PB	CR-C1-CD-CC


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	499/520 (95%)	0.78	58 (11%) 9 10	30, 42, 58, 82	0
1	B	494/520 (95%)	0.78	47 (9%) 14 15	30, 41, 56, 73	0
All	All	993/1040 (95%)	0.78	105 (10%) 11 12	30, 42, 57, 82	0

All (105) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	501	ILE	7.5
1	B	496	ILE	6.3
1	A	498	LEU	5.3
1	B	107	TRP	5.1
1	A	500	THR	4.8
1	B	163	GLN	4.3
1	B	460	ASP	4.1
1	A	497	GLY	3.8
1	B	354	ARG	3.8
1	A	460	ASP	3.7
1	B	92	VAL	3.7
1	A	495	LEU	3.7
1	A	163	GLN	3.7
1	B	3	ASN	3.7
1	A	93	LYS	3.7
1	B	495	LEU	3.7
1	A	107	TRP	3.6
1	A	95	LYS	3.6
1	A	49	GLN	3.6
1	A	481	PHE	3.6
1	B	318	ASP	3.5
1	A	321	GLU	3.5
1	B	493	LEU	3.5
1	B	49	GLN	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	499	THR	3.4
1	A	3	ASN	3.4
1	A	308	LYS	3.2
1	B	321	GLU	3.2
1	A	318	ASP	3.2
1	B	479	THR	3.2
1	B	106	VAL	3.1
1	A	494	ARG	3.0
1	B	476	PRO	3.0
1	A	242	ARG	2.9
1	B	141	GLU	2.9
1	A	470	VAL	2.9
1	B	242	ARG	2.8
1	B	224	LEU	2.8
1	B	494	ARG	2.8
1	B	455	GLY	2.8
1	A	224	LEU	2.8
1	A	29	ASN	2.7
1	B	26	SER	2.7
1	B	243	GLU	2.7
1	B	227	ASP	2.7
1	B	249	THR	2.7
1	A	319	GLY	2.6
1	B	246	LEU	2.6
1	A	320	GLU	2.6
1	B	226	GLY	2.6
1	B	414	LEU	2.6
1	A	370	LYS	2.6
1	B	93	LYS	2.6
1	A	468	GLU	2.5
1	A	412	ARG	2.5
1	B	319	GLY	2.5
1	A	492	LEU	2.5
1	B	482	LEU	2.5
1	A	302	LYS	2.5
1	A	474	ALA	2.5
1	A	243	GLU	2.4
1	A	280	MET	2.4
1	B	254	MET	2.4
1	B	370	LYS	2.4
1	A	226	GLY	2.4
1	A	477	ILE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	493	LEU	2.4
1	A	483	GLU	2.4
1	B	374	SER	2.4
1	A	254	MET	2.4
1	B	103	PHE	2.3
1	A	109	PRO	2.3
1	A	101	GLY	2.3
1	A	482	LEU	2.3
1	A	154	LYS	2.3
1	A	227	ASP	2.3
1	B	27	GLY	2.3
1	A	480	THR	2.3
1	A	128	GLU	2.3
1	A	475	GLN	2.3
1	B	223	ASP	2.3
1	B	485	HIS	2.3
1	B	481	PHE	2.2
1	A	104	PRO	2.2
1	A	354	ARG	2.2
1	A	484	ARG	2.2
1	B	303	GLU	2.2
1	B	475	GLN	2.2
1	A	249	THR	2.2
1	B	478	THR	2.2
1	A	463	TRP	2.2
1	A	141	GLU	2.2
1	A	478	THR	2.2
1	B	52	LYS	2.1
1	B	302	LYS	2.1
1	A	252	HIS	2.1
1	B	480	THR	2.1
1	A	159	GLU	2.1
1	A	251	ASN	2.1
1	A	489	VAL	2.1
1	A	479	THR	2.1
1	B	483	GLU	2.1
1	B	241	THR	2.0
1	B	355	LEU	2.0
1	A	491	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	1PB	A	1503	16/16	0.86	0.18	59,60,61,61	0
3	1PB	B	1498	16/16	0.89	0.18	59,60,61,61	0
2	FAD	A	1502	53/53	0.97	0.08	36,38,45,46	0
2	FAD	B	1497	53/53	0.97	0.08	36,38,45,46	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FAD A 1502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD B 1497:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.