



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 07:45 PM UTC

PDB ID : 3OJ3 / pdb_00003oj3
Title : Crystal structure of the A20 ZnF4 and ubiquitin complex
Authors : Bosanac, I.; Hymowitz, S.G.
Deposited on : 2010-08-20
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	79	71% 20% • 8%
1	B	79	62% 29% 9%
1	C	79	66% 25% • 8%
1	D	79	67% 24% • 8%
1	E	79	66% 25% • 8%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	79	
1	G	79	
1	H	79	
2	I	49	
2	J	49	
2	K	49	
2	L	49	
2	M	49	
2	N	49	
2	O	49	
2	P	49	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6729 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ubiquitin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	73	Total	C	N	O	S	0	0	0
			582	368	99	114	1			
1	B	72	Total	C	N	O	S	0	0	0
			574	362	98	113	1			
1	C	73	Total	C	N	O	S	0	0	0
			582	368	99	114	1			
1	D	73	Total	C	N	O	S	0	0	0
			582	368	99	114	1			
1	E	73	Total	C	N	O	S	0	0	0
			582	368	99	114	1			
1	F	73	Total	C	N	O	S	0	0	0
			582	368	99	114	1			
1	G	71	Total	C	N	O	S	0	0	0
			563	356	94	112	1			
1	H	71	Total	C	N	O	S	0	0	0
			563	356	94	112	1			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P0CG47
A	-1	SER	-	expression tag	UNP P0CG47
A	0	HIS	-	expression tag	UNP P0CG47
B	-2	GLY	-	expression tag	UNP P0CG47
B	-1	SER	-	expression tag	UNP P0CG47
B	0	HIS	-	expression tag	UNP P0CG47
C	-2	GLY	-	expression tag	UNP P0CG47
C	-1	SER	-	expression tag	UNP P0CG47
C	0	HIS	-	expression tag	UNP P0CG47
D	-2	GLY	-	expression tag	UNP P0CG47
D	-1	SER	-	expression tag	UNP P0CG47
D	0	HIS	-	expression tag	UNP P0CG47
E	-2	GLY	-	expression tag	UNP P0CG47

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-1	SER	-	expression tag	UNP P0CG47
E	0	HIS	-	expression tag	UNP P0CG47
F	-2	GLY	-	expression tag	UNP P0CG47
F	-1	SER	-	expression tag	UNP P0CG47
F	0	HIS	-	expression tag	UNP P0CG47
G	-2	GLY	-	expression tag	UNP P0CG47
G	-1	SER	-	expression tag	UNP P0CG47
G	0	HIS	-	expression tag	UNP P0CG47
H	-2	GLY	-	expression tag	UNP P0CG47
H	-1	SER	-	expression tag	UNP P0CG47
H	0	HIS	-	expression tag	UNP P0CG47

- Molecule 2 is a protein called Tumor necrosis factor alpha-induced protein 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	33	Total	C	N	O	S	0	0	0
			260	164	45	47	4			
2	J	30	Total	C	N	O	S	0	0	0
			240	152	41	43	4			
2	K	31	Total	C	N	O	S	0	0	0
			249	158	43	44	4			
2	L	31	Total	C	N	O	S	0	0	0
			249	158	43	44	4			
2	M	31	Total	C	N	O	S	0	0	0
			249	158	43	44	4			
2	N	30	Total	C	N	O	S	0	0	0
			240	152	41	43	4			
2	O	32	Total	C	N	O	S	0	0	0
			256	162	44	46	4			
2	P	31	Total	C	N	O	S	0	0	0
			249	158	43	44	4			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	587	GLY	-	expression tag	UNP P21580
I	588	SER	-	expression tag	UNP P21580
I	589	PRO	-	expression tag	UNP P21580
I	590	GLU	-	expression tag	UNP P21580
I	591	PHE	-	expression tag	UNP P21580
J	587	GLY	-	expression tag	UNP P21580
J	588	SER	-	expression tag	UNP P21580

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
J	589	PRO	-	expression tag	UNP P21580
J	590	GLU	-	expression tag	UNP P21580
J	591	PHE	-	expression tag	UNP P21580
K	587	GLY	-	expression tag	UNP P21580
K	588	SER	-	expression tag	UNP P21580
K	589	PRO	-	expression tag	UNP P21580
K	590	GLU	-	expression tag	UNP P21580
K	591	PHE	-	expression tag	UNP P21580
L	587	GLY	-	expression tag	UNP P21580
L	588	SER	-	expression tag	UNP P21580
L	589	PRO	-	expression tag	UNP P21580
L	590	GLU	-	expression tag	UNP P21580
L	591	PHE	-	expression tag	UNP P21580
M	587	GLY	-	expression tag	UNP P21580
M	588	SER	-	expression tag	UNP P21580
M	589	PRO	-	expression tag	UNP P21580
M	590	GLU	-	expression tag	UNP P21580
M	591	PHE	-	expression tag	UNP P21580
N	587	GLY	-	expression tag	UNP P21580
N	588	SER	-	expression tag	UNP P21580
N	589	PRO	-	expression tag	UNP P21580
N	590	GLU	-	expression tag	UNP P21580
N	591	PHE	-	expression tag	UNP P21580
O	587	GLY	-	expression tag	UNP P21580
O	588	SER	-	expression tag	UNP P21580
O	589	PRO	-	expression tag	UNP P21580
O	590	GLU	-	expression tag	UNP P21580
O	591	PHE	-	expression tag	UNP P21580
P	587	GLY	-	expression tag	UNP P21580
P	588	SER	-	expression tag	UNP P21580
P	589	PRO	-	expression tag	UNP P21580
P	590	GLU	-	expression tag	UNP P21580
P	591	PHE	-	expression tag	UNP P21580

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	I	1	Total Zn 1 1	0	0
3	J	1	Total Zn 1 1	0	0
3	K	1	Total Zn 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	L	1	Total 1	Zn 1	0	0
3	M	1	Total 1	Zn 1	0	0
3	N	1	Total 1	Zn 1	0	0
3	O	1	Total 1	Zn 1	0	0
3	P	1	Total 1	Zn 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	12	Total 12	O 12	0	0
4	B	11	Total 11	O 11	0	0
4	C	18	Total 18	O 18	0	0
4	D	7	Total 7	O 7	0	0
4	E	9	Total 9	O 9	0	0
4	F	15	Total 15	O 15	0	0
4	G	12	Total 12	O 12	0	0
4	H	7	Total 7	O 7	0	0
4	I	4	Total 4	O 4	0	0
4	J	3	Total 3	O 3	0	0
4	K	3	Total 3	O 3	0	0
4	L	2	Total 2	O 2	0	0
4	M	4	Total 4	O 4	0	0
4	N	4	Total 4	O 4	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	O	6	Total	O	0	0
			6	6		
4	P	2	Total	O	0	0
			2	2		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ubiquitin

Chain A: 




- Molecule 1: Ubiquitin

Chain B: 




- Molecule 1: Ubiquitin

Chain C: 



- Molecule 1: Ubiquitin

Chain D: 



- Molecule 1: Ubiquitin

Chain E: 




- Molecule 1: Ubiquitin

Chain F: 




- Molecule 1: Ubiquitin

Chain G: 68% 19% 10%



- Molecule 1: Ubiquitin

Chain H: 70% 18% 10%



- Molecule 2: Tumor necrosis factor alpha-induced protein 3

Chain I: 29% 37% 33%



- Molecule 2: Tumor necrosis factor alpha-induced protein 3

Chain J: 39% 18% 39%



- Molecule 2: Tumor necrosis factor alpha-induced protein 3

Chain K: 31% 29% 37%




- Molecule 2: Tumor necrosis factor alpha-induced protein 3

Chain L: 35% 24% 37%





- Molecule 2: Tumor necrosis factor alpha-induced protein 3



Chain M: 33% 24% 6% 37%




● Molecule 2: Tumor necrosis factor alpha-induced protein 3



● Molecule 2: Tumor necrosis factor alpha-induced protein 3



● Molecule 2: Tumor necrosis factor alpha-induced protein 3



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	42.83Å 170.03Å 66.24Å 90.00° 90.10° 90.00°	Depositor
Resolution (Å)	43.06 – 2.50 43.06 – 2.50	Depositor EDS
% Data completeness (in resolution range)	96.1 (43.06-2.50) 95.7 (43.06-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.18 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.6.4_486, REFMAC 5.2.0019	Depositor
R, R_{free}	0.203 , 0.226 0.202 , 0.221	Depositor DCC
R_{free} test set	1630 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	50.9	Xtriage
Anisotropy	0.453	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 61.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.24$	Xtriage
Estimated twinning fraction	0.317 for h,-k,-l	Xtriage
Reported twinning fraction	0.351 for h,-k,-l	Depositor
Outliers	0 of 31426 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6729	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/588	0.84	2/792 (0.3%)
1	B	0.60	0/580	0.85	0/781
1	C	0.60	0/588	0.83	2/792 (0.3%)
1	D	0.62	0/588	0.84	2/792 (0.3%)
1	E	0.58	0/588	0.84	2/792 (0.3%)
1	F	0.58	0/588	0.82	2/792 (0.3%)
1	G	0.55	0/569	0.81	2/767 (0.3%)
1	H	0.57	0/569	0.84	2/767 (0.3%)
2	I	0.60	0/265	0.87	0/352
2	J	0.50	0/245	0.84	0/326
2	K	0.55	0/254	0.87	0/337
2	L	0.52	0/254	0.91	1/337 (0.3%)
2	M	0.55	0/254	0.91	0/337
2	N	0.57	0/245	0.87	1/326 (0.3%)
2	O	0.58	0/261	0.98	0/347
2	P	0.50	0/254	0.86	0/337
All	All	0.57	0/6690	0.85	16/8974 (0.2%)

There are no bond length outliers.

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	37	PRO	CA-C-N	5.90	125.61	119.82
1	C	37	PRO	C-N-CA	5.90	125.61	119.82
1	D	37	PRO	CA-C-N	5.69	125.40	119.82
1	D	37	PRO	C-N-CA	5.69	125.40	119.82
1	F	37	PRO	CA-C-N	5.65	125.36	119.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	582	0	610	14	0
1	B	574	0	599	24	0
1	C	582	0	610	18	0
1	D	582	0	610	17	0
1	E	582	0	610	21	0
1	F	582	0	610	29	0
1	G	563	0	586	13	0
1	H	563	0	586	13	0
2	I	260	0	249	20	0
2	J	240	0	226	8	0
2	K	249	0	239	17	0
2	L	249	0	239	9	0
2	M	249	0	240	14	0
2	N	240	0	226	18	0
2	O	256	0	246	27	0
2	P	249	0	239	12	0
3	I	1	0	0	0	0
3	J	1	0	0	0	0
3	K	1	0	0	0	0
3	L	1	0	0	0	0
3	M	1	0	0	0	0
3	N	1	0	0	0	0
3	O	1	0	0	0	0
3	P	1	0	0	0	0
4	A	12	0	0	2	0
4	B	11	0	0	8	0
4	C	18	0	0	1	0
4	D	7	0	0	3	0
4	E	9	0	0	1	0
4	F	15	0	0	6	0
4	G	12	0	0	3	0
4	H	7	0	0	1	0
4	I	4	0	0	3	0
4	J	3	0	0	0	0
4	K	3	0	0	1	0
4	L	2	0	0	0	0
4	M	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	N	4	0	0	0	0
4	O	6	0	0	1	0
4	P	2	0	0	0	0
All	All	6729	0	6725	223	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 223 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:O:618:PRO:HD2	4:O:28:HOH:O	1.52	1.08
2:O:629:ILE:O	2:O:633:GLU:HG2	1.76	0.84
2:M:631:TYR:HE2	2:M:635:LYS:HZ1	1.28	0.80
2:O:607:CYS:SG	2:O:624:CYS:N	2.55	0.79
1:G:9:THR:HG23	4:G:109:HOH:O	1.84	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	71/79 (90%)	70 (99%)	1 (1%)	0	100	100
1	B	70/79 (89%)	68 (97%)	2 (3%)	0	100	100
1	C	71/79 (90%)	69 (97%)	2 (3%)	0	100	100
1	D	71/79 (90%)	70 (99%)	1 (1%)	0	100	100
1	E	71/79 (90%)	69 (97%)	2 (3%)	0	100	100
1	F	71/79 (90%)	69 (97%)	2 (3%)	0	100	100
1	G	69/79 (87%)	68 (99%)	1 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	69/79 (87%)	67 (97%)	2 (3%)	0	100	100
2	I	31/49 (63%)	27 (87%)	3 (10%)	1 (3%)	3	4
2	J	28/49 (57%)	25 (89%)	2 (7%)	1 (4%)	2	3
2	K	29/49 (59%)	26 (90%)	2 (7%)	1 (3%)	3	4
2	L	29/49 (59%)	26 (90%)	2 (7%)	1 (3%)	3	4
2	M	29/49 (59%)	25 (86%)	3 (10%)	1 (3%)	3	4
2	N	28/49 (57%)	25 (89%)	2 (7%)	1 (4%)	2	3
2	O	30/49 (61%)	25 (83%)	4 (13%)	1 (3%)	3	4
2	P	29/49 (59%)	26 (90%)	2 (7%)	1 (3%)	3	4
All	All	796/1024 (78%)	755 (95%)	33 (4%)	8 (1%)	12	24

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	O	610	ALA
2	I	610	ALA
2	J	610	ALA
2	K	610	ALA
2	L	610	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	67/70 (96%)	66 (98%)	1 (2%)	57	80
1	B	66/70 (94%)	65 (98%)	1 (2%)	57	80
1	C	67/70 (96%)	65 (97%)	2 (3%)	36	64
1	D	67/70 (96%)	65 (97%)	2 (3%)	36	64
1	E	67/70 (96%)	66 (98%)	1 (2%)	57	80
1	F	67/70 (96%)	66 (98%)	1 (2%)	57	80
1	G	65/70 (93%)	64 (98%)	1 (2%)	57	80

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	65/70 (93%)	64 (98%)	1 (2%)	57	80
2	I	28/40 (70%)	27 (96%)	1 (4%)	31	58
2	J	26/40 (65%)	25 (96%)	1 (4%)	29	56
2	K	27/40 (68%)	26 (96%)	1 (4%)	30	57
2	L	27/40 (68%)	26 (96%)	1 (4%)	30	57
2	M	27/40 (68%)	25 (93%)	2 (7%)	13	27
2	N	26/40 (65%)	25 (96%)	1 (4%)	29	56
2	O	28/40 (70%)	25 (89%)	3 (11%)	6	13
2	P	27/40 (68%)	26 (96%)	1 (4%)	30	57
All	All	747/880 (85%)	726 (97%)	21 (3%)	38	66

5 of 21 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	M	617	THR
2	O	617	THR
2	P	617	THR
2	O	625	THR
2	N	617	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 27 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	62	GLN
1	G	62	GLN
2	M	620	ASN
1	G	60	ASN
1	H	60	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	73/79 (92%)	-1.30	0 100 100	36, 54, 71, 78	0
1	B	72/79 (91%)	-1.31	0 100 100	36, 53, 72, 80	0
1	C	73/79 (92%)	-1.34	0 100 100	35, 52, 71, 81	0
1	D	73/79 (92%)	-1.36	0 100 100	36, 52, 74, 79	0
1	E	73/79 (92%)	-1.33	0 100 100	37, 53, 73, 80	0
1	F	73/79 (92%)	-1.31	0 100 100	36, 53, 74, 79	0
1	G	71/79 (89%)	-1.18	0 100 100	36, 54, 74, 79	0
1	H	71/79 (89%)	-1.27	0 100 100	36, 53, 73, 80	0
2	I	33/49 (67%)	-1.19	0 100 100	44, 57, 71, 79	0
2	J	30/49 (61%)	-1.19	0 100 100	43, 57, 71, 74	0
2	K	31/49 (63%)	-1.29	0 100 100	43, 58, 70, 72	0
2	L	31/49 (63%)	-1.20	0 100 100	44, 56, 72, 82	0
2	M	31/49 (63%)	-1.25	0 100 100	45, 58, 72, 75	0
2	N	30/49 (61%)	-1.24	0 100 100	44, 57, 71, 72	0
2	O	32/49 (65%)	-1.13	0 100 100	43, 67, 79, 84	0
2	P	31/49 (63%)	-1.25	0 100 100	44, 58, 72, 78	0
All	All	828/1024 (80%)	-1.28	0 100 100	35, 56, 74, 84	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ZN	I	901	1/1	1.00	0.01	46,46,46,46	0
3	ZN	J	902	1/1	1.00	0.02	53,53,53,53	0
3	ZN	K	903	1/1	1.00	0.02	53,53,53,53	0
3	ZN	L	904	1/1	1.00	0.01	48,48,48,48	0
3	ZN	M	905	1/1	1.00	0.01	50,50,50,50	0
3	ZN	N	906	1/1	1.00	0.02	55,55,55,55	0
3	ZN	O	907	1/1	1.00	0.01	73,73,73,73	0
3	ZN	P	908	1/1	1.00	0.01	60,60,60,60	0

6.5 Other polymers [i](#)

There are no such residues in this entry.