



wwPDB X-ray Structure Validation Summary Report ⓘ

May 7, 2026 – 09:09 AM EDT

PDB ID : 3OJY / pdb_00003ojy
Title : Crystal Structure of Human Complement Component C8
Authors : Lovelace, L.L.; Cooper, C.L.; Sodetz, J.M.; Lebioda, L.
Deposited on : 2010-08-23
Resolution : 2.51 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	554	
2	B	537	
3	C	182	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9234 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Complement component C8 alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	478	Total	C	N	O	S	0	0	0
			3799	2359	666	736	38			

- Molecule 2 is a protein called Complement component C8 beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	504	Total	C	N	O	P S	0	0	0
			4050	2521	722	771	1 35			


- Molecule 3 is a protein called Complement component C8 gamma chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	165	Total	C	N	O	S	0	0	0
			1295	828	224	239	4			

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Ca	0	0
			1	1		
4	B	1	Total	Ca	0	0
			1	1		

- Molecule 5 is beta-D-mannopyranose (CCD ID: BMA) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	A	1	Total	C	O	0	0
			11	6	5		
5	B	1	Total	C	O	0	0
			11	6	5		
5	B	1	Total	C	O	0	0
			11	6	5		
5	B	1	Total	C	O	0	0
			11	6	5		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Complement component C8 alpha chain




- Molecule 2: Complement component C8 beta chain





• Molecule 3: Complement component C8 gamma chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	139.57Å 139.57Å 127.16Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	45.69 – 2.51 45.69 – 2.51	Depositor EDS
% Data completeness (in resolution range)	93.1 (45.69-2.51) 93.1 (45.69-2.51)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.27 (at 2.51Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.249 , 0.337 0.248 , 0.335	Depositor DCC
R_{free} test set	2267 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	50.6	Xtriage
Anisotropy	0.087	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 28.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.038 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9234	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, CA, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.89	4/3873 (0.1%)	1.09	6/5207 (0.1%)
2	B	0.95	6/4131 (0.1%)	1.20	29/5574 (0.5%)
3	C	0.85	0/1323	1.13	5/1794 (0.3%)
All	All	0.91	10/9327 (0.1%)	1.14	40/12575 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	2
3	C	0	1
All	All	0	3

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	234	THR	CA-CB	6.77	1.61	1.52
2	B	16	TRP	CG-CD1	5.42	1.50	1.36
1	A	515	TRP	CG-CD1	5.38	1.50	1.36
1	A	518	TRP	CG-CD1	5.29	1.50	1.36
1	A	512	TRP	CG-CD1	5.11	1.49	1.36

The worst 5 of 40 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	255	ARG	N-CA-C	11.04	123.45	107.88
3	C	111	ASP	N-CA-C	-9.95	97.70	110.53
2	B	221	GLY	N-CA-C	-8.76	102.00	115.66

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	38	LYS	N-CA-C	7.78	121.28	111.24
2	B	108	CYS	N-CA-C	7.57	120.34	110.43

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	107	LYS	Peptide
2	B	96	SER	Peptide
3	C	110	THR	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3799	0	3596	131	1
2	B	4050	0	3853	234	1
3	C	1295	0	1259	47	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	44	0	40	1	0
5	B	44	0	40	2	0
All	All	9234	0	8788	397	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 397 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:93:GLY:O	2:B:261:TYR:CD2	2.04	1.09
2:B:520:GLN:NE2	2:B:520:GLN:HA	1.52	1.08
1:A:322:GLU:HB3	1:A:392:ARG:HB2	1.34	1.08
2:B:516:ASN:O	2:B:518:PRO:HD3	1.56	1.04
2:B:146:CYS:O	2:B:148:PRO:HD3	1.58	1.03

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:10:GLN:NE2	2:B:520:GLN:O[2_545]	2.07	0.13

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	464/554 (84%)	386 (83%)	55 (12%)	23 (5%)	1	2
2	B	495/537 (92%)	412 (83%)	55 (11%)	28 (6%)	1	1
3	C	161/182 (88%)	146 (91%)	11 (7%)	4 (2%)	4	7
All	All	1120/1273 (88%)	944 (84%)	121 (11%)	55 (5%)	1	2

5 of 55 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	194	GLY
1	A	369	CYS
2	B	202	PHE
2	B	218	SER
2	B	350	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	412/466 (88%)	343 (83%)	69 (17%)	2	4
2	B	446/472 (94%)	352 (79%)	94 (21%)	1	2
3	C	134/149 (90%)	108 (81%)	26 (19%)	1	2
All	All	992/1087 (91%)	803 (81%)	189 (19%)	1	3

5 of 189 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	230	ARG
2	B	411	VAL
2	B	255	ARG
2	B	356	ILE
2	B	463	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	48	HIS
3	C	175	HIS
3	C	60	GLN
3	C	95	GLN
2	B	110	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TPO	B	364	2	8,10,11	1.16	1 (12%)	10,14,16	1.14	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TPO	B	364	2	-	5/9/11/13	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	364	TPO	P-OG1	2.44	1.63	1.59

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	364	TPO	O3P-P-O2P	2.04	115.47	107.80

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	364	TPO	N-CA-CB-CG2
2	B	364	TPO	N-CA-CB-OG1
2	B	364	TPO	C-CA-CB-CG2
2	B	364	TPO	CA-CB-OG1-P
2	B	364	TPO	CG2-CB-OG1-P

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	364	TPO	2	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	BMA	A	556	1	11,11,12	0.68	0	15,15,17	1.38	2 (13%)
5	BMA	B	540	2	11,11,12	0.68	0	15,15,17	1.55	3 (20%)
5	BMA	A	558	1	11,11,12	0.68	0	15,15,17	1.59	3 (20%)
5	BMA	A	559	1	11,11,12	1.00	0	15,15,17	2.86	5 (33%)
5	BMA	B	539	2	11,11,12	0.49	0	15,15,17	1.05	1 (6%)
5	BMA	A	557	1	11,11,12	0.71	0	15,15,17	1.84	3 (20%)
5	BMA	B	541	2	11,11,12	0.68	0	15,15,17	1.17	1 (6%)
5	BMA	B	542	2	11,11,12	0.81	0	15,15,17	0.91	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BMA	A	556	1	-	2/2/19/22	0/1/1/1
5	BMA	B	540	2	-	0/2/19/22	0/1/1/1
5	BMA	A	558	1	-	2/2/19/22	0/1/1/1
5	BMA	A	559	1	-	2/2/19/22	0/1/1/1
5	BMA	B	539	2	-	0/2/19/22	0/1/1/1
5	BMA	A	557	1	-	2/2/19/22	0/1/1/1
5	BMA	B	541	2	-	2/2/19/22	0/1/1/1
5	BMA	B	542	2	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	559	BMA	C1-O5-C5	-8.19	101.22	112.19
5	A	557	BMA	C1-O5-C5	-5.14	105.29	112.19
5	A	559	BMA	C3-C4-C5	4.05	117.57	110.23
5	A	558	BMA	C1-O5-C5	-3.74	107.17	112.19
5	A	559	BMA	O5-C5-C6	3.61	114.68	107.66

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	557	BMA	O5-C5-C6-O6
5	B	541	BMA	O5-C5-C6-O6
5	A	557	BMA	C4-C5-C6-O6
5	A	559	BMA	O5-C5-C6-O6
5	B	541	BMA	C4-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	558	BMA	1	0
5	B	541	BMA	1	0
5	B	542	BMA	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	478/554 (86%)	0.69	40 (8%) 17 15	8, 54, 78, 94	1 (0%)
2	B	503/537 (93%)	0.70	59 (11%) 9 7	26, 48, 74, 86	0
3	C	165/182 (90%)	0.65	8 (4%) 35 31	28, 53, 77, 83	0
All	All	1146/1273 (90%)	0.69	107 (9%) 14 12	8, 50, 77, 94	1 (0%)

The worst 5 of 107 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	281	ASP	10.8
2	B	93	GLY	8.0
2	B	346	VAL	5.3
1	A	106	ILE	4.6
1	A	386	VAL	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	TPO	B	364	11/12	0.53	0.17	69,71,79,79	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	BMA	B	542	11/12	0.52	0.16	56,60,62,62	0
5	BMA	A	559	11/12	0.62	0.18	79,81,81,82	0
5	BMA	B	540	11/12	0.67	0.13	63,67,67,70	0
5	BMA	A	556	11/12	0.84	0.11	47,51,56,59	0
5	BMA	B	539	11/12	0.84	0.10	56,59,62,64	0
5	BMA	A	557	11/12	0.86	0.11	58,61,61,62	0
5	BMA	A	558	11/12	0.86	0.11	63,65,67,67	0
5	BMA	B	541	11/12	0.93	0.07	39,42,44,46	0
4	CA	A	555	1/1	0.99	0.03	45,45,45,45	0
4	CA	B	538	1/1	0.99	0.03	38,38,38,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.