



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 2, 2026 – 02:55 AM UTC

PDB ID : 4OJN / pdb_00004ojn
Title : Crystal structure of human muscle L-lactate dehydrogenase
Authors : Kolappan, S.; Craig, L.
Deposited on : 2014-01-21
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	337	3% 87% 10% ..
1	B	337	% 86% 11% ..
1	C	337	2% 84% 13% ..
1	D	337	3% 83% 15% ..
1	E	337	3% 83% 14% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	337	
1	G	337	
1	H	337	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	1PE	H	400	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 21736 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-lactate dehydrogenase A chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	332	Total	C	N	O	S	0	10	0
			2661	1699	454	495	13			
1	B	331	Total	C	N	O	S	0	4	0
			2604	1663	444	484	13			
1	C	332	Total	C	N	O	S	0	7	0
			2637	1681	451	492	13			
1	D	332	Total	C	N	O	S	0	10	0
			2664	1696	458	497	13			
1	E	331	Total	C	N	O	S	0	5	0
			2613	1668	445	487	13			
1	F	331	Total	C	N	O	S	0	11	0
			2666	1702	458	493	13			
1	G	331	Total	C	N	O	S	0	8	0
			2639	1679	454	493	13			
1	H	331	Total	C	N	O	S	0	1	0
			2579	1648	440	478	13			

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	333	HIS	-	expression tag	UNP P00338
A	334	HIS	-	expression tag	UNP P00338
A	335	HIS	-	expression tag	UNP P00338
A	336	HIS	-	expression tag	UNP P00338
A	337	HIS	-	expression tag	UNP P00338
A	338	HIS	-	expression tag	UNP P00338
B	333	HIS	-	expression tag	UNP P00338
B	334	HIS	-	expression tag	UNP P00338
B	335	HIS	-	expression tag	UNP P00338
B	336	HIS	-	expression tag	UNP P00338
B	337	HIS	-	expression tag	UNP P00338
B	338	HIS	-	expression tag	UNP P00338
C	333	HIS	-	expression tag	UNP P00338

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	334	HIS	-	expression tag	UNP P00338
C	335	HIS	-	expression tag	UNP P00338
C	336	HIS	-	expression tag	UNP P00338
C	337	HIS	-	expression tag	UNP P00338
C	338	HIS	-	expression tag	UNP P00338
D	333	HIS	-	expression tag	UNP P00338
D	334	HIS	-	expression tag	UNP P00338
D	335	HIS	-	expression tag	UNP P00338
D	336	HIS	-	expression tag	UNP P00338
D	337	HIS	-	expression tag	UNP P00338
D	338	HIS	-	expression tag	UNP P00338
E	333	HIS	-	expression tag	UNP P00338
E	334	HIS	-	expression tag	UNP P00338
E	335	HIS	-	expression tag	UNP P00338
E	336	HIS	-	expression tag	UNP P00338
E	337	HIS	-	expression tag	UNP P00338
E	338	HIS	-	expression tag	UNP P00338
F	333	HIS	-	expression tag	UNP P00338
F	334	HIS	-	expression tag	UNP P00338
F	335	HIS	-	expression tag	UNP P00338
F	336	HIS	-	expression tag	UNP P00338
F	337	HIS	-	expression tag	UNP P00338
F	338	HIS	-	expression tag	UNP P00338
G	333	HIS	-	expression tag	UNP P00338
G	334	HIS	-	expression tag	UNP P00338
G	335	HIS	-	expression tag	UNP P00338
G	336	HIS	-	expression tag	UNP P00338
G	337	HIS	-	expression tag	UNP P00338
G	338	HIS	-	expression tag	UNP P00338
H	333	HIS	-	expression tag	UNP P00338
H	334	HIS	-	expression tag	UNP P00338
H	335	HIS	-	expression tag	UNP P00338
H	336	HIS	-	expression tag	UNP P00338
H	337	HIS	-	expression tag	UNP P00338
H	338	HIS	-	expression tag	UNP P00338

- Molecule 2 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: C₁₀H₂₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			16	10	6		
2	B	1	Total	C	O	0	0
			16	10	6		
2	C	1	Total	C	O	0	0
			16	10	6		
2	D	1	Total	C	O	0	0
			16	10	6		
2	E	1	Total	C	O	0	0
			16	10	6		
2	F	1	Total	C	O	0	0
			16	10	6		
2	G	1	Total	C	O	0	0
			16	10	6		
2	H	1	Total	C	O	0	0
			16	10	6		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	C	O	0	0
			6	3	3		
3	D	1	Total	C	O	0	0
			6	3	3		
3	E	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	57	Total	O	0	0
			57	57		
4	B	66	Total	O	0	0
			66	66		
4	C	70	Total	O	0	0
			70	70		
4	D	84	Total	O	0	0
			84	84		
4	E	50	Total	O	0	0
			50	50		
4	F	66	Total	O	0	0
			66	66		
4	G	75	Total	O	0	0
			75	75		
4	H	59	Total	O	0	0
			59	59		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: L-lactate dehydrogenase A chain




- Molecule 1: L-lactate dehydrogenase A chain



- Molecule 1: L-lactate dehydrogenase A chain



- Molecule 1: L-lactate dehydrogenase A chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	147.14Å 147.14Å 333.24Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.84 – 2.40 19.84 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.5 (19.84-2.40) 99.5 (19.84-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.00 (at 2.41Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.213 , 0.250 0.214 , 0.247	Depositor DCC
R_{free} test set	8138 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	35.6	Xtriage
Anisotropy	0.409	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.037 for -h,-k,l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	21736	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 1PE, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.66	0/2709	0.85	2/3663 (0.1%)
1	B	0.67	0/2649	0.90	2/3584 (0.1%)
1	C	0.67	0/2682	0.85	2/3628 (0.1%)
1	D	0.66	0/2709	0.82	0/3661
1	E	0.68	0/2658	0.87	1/3596 (0.0%)
1	F	0.66	0/2711	0.84	1/3665 (0.0%)
1	G	0.69	0/2683	0.86	1/3626 (0.0%)
1	H	0.68	0/2624	0.88	0/3550
All	All	0.67	0/21425	0.86	9/28973 (0.0%)

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	121	ILE	CA-C-N	-6.71	111.89	119.28
1	C	121	ILE	C-N-CA	-6.71	111.89	119.28
1	G	200	VAL	N-CA-C	5.81	112.42	106.21
1	B	74	THR	CA-C-N	5.67	125.20	119.19
1	B	74	THR	C-N-CA	5.67	125.20	119.19

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2661	0	2733	25	0
1	B	2604	0	2681	28	0
1	C	2637	0	2707	42	0
1	D	2664	0	2735	39	0
1	E	2613	0	2686	35	0
1	F	2666	0	2749	40	0
1	G	2639	0	2712	19	0
1	H	2579	0	2661	42	0
2	A	16	0	22	4	0
2	B	16	0	22	1	0
2	C	16	0	22	2	0
2	D	16	0	22	2	0
2	E	16	0	22	4	0
2	F	16	0	22	1	0
2	G	16	0	22	2	0
2	H	16	0	22	8	0
3	D	12	0	16	1	0
3	E	6	0	8	0	0
4	A	57	0	0	0	0
4	B	66	0	0	0	0
4	C	70	0	0	4	0
4	D	84	0	0	6	0
4	E	50	0	0	1	0
4	F	66	0	0	1	0
4	G	75	0	0	2	0
4	H	59	0	0	5	0
All	All	21736	0	21864	246	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 246 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:70[A]:LEU:HD13	1:C:70[A]:LEU:O	1.21	1.30
1:B:239[A]:TYR:HD2	1:B:239[A]:TYR:N	1.52	1.07
1:C:70[A]:LEU:C	1:C:70[A]:LEU:CD1	2.30	1.04
1:C:70[A]:LEU:HD13	1:C:70[A]:LEU:C	1.79	1.03
1:C:70[A]:LEU:O	1:C:70[A]:LEU:CD1	2.06	1.02

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/337 (101%)	329 (97%)	9 (3%)	1 (0%)	36	50
1	B	333/337 (99%)	320 (96%)	13 (4%)	0	100	100
1	C	337/337 (100%)	331 (98%)	6 (2%)	0	100	100
1	D	340/337 (101%)	321 (94%)	18 (5%)	1 (0%)	36	50
1	E	334/337 (99%)	321 (96%)	12 (4%)	1 (0%)	36	50
1	F	340/337 (101%)	328 (96%)	12 (4%)	0	100	100
1	G	337/337 (100%)	328 (97%)	8 (2%)	1 (0%)	36	50
1	H	330/337 (98%)	317 (96%)	10 (3%)	3 (1%)	14	22
All	All	2690/2696 (100%)	2595 (96%)	88 (3%)	7 (0%)	36	50

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	14	LYS
1	H	17	GLN
1	H	15	GLU
1	E	17	GLN
1	H	18	THR

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	298/293 (102%)	285 (96%)	13 (4%)	25	43

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	291/293 (99%)	280 (96%)	11 (4%)	29	49
1	C	295/293 (101%)	279 (95%)	16 (5%)	20	35
1	D	298/293 (102%)	288 (97%)	10 (3%)	32	54
1	E	292/293 (100%)	276 (94%)	16 (6%)	19	34
1	F	298/293 (102%)	282 (95%)	16 (5%)	20	35
1	G	295/293 (101%)	277 (94%)	18 (6%)	17	30
1	H	288/293 (98%)	273 (95%)	15 (5%)	21	36
All	All	2355/2344 (100%)	2240 (95%)	115 (5%)	24	39

5 of 115 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	128	SER
1	H	277	ILE
1	F	116[B]	ILE
1	H	212	LYS
1	G	330	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 88 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	111	GLN
1	G	226	GLN
1	F	205	ASN
1	G	100	GLN
1	H	20	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	1PE	F	400	-	15,15,15	0.35	0	14,14,14	0.52	0
2	1PE	E	401	-	15,15,15	0.52	0	14,14,14	0.72	0
2	1PE	C	400	-	15,15,15	0.48	0	14,14,14	0.49	0
2	1PE	H	400	-	15,15,15	0.46	0	14,14,14	0.64	0
3	GOL	E	402	-	5,5,5	0.32	0	5,5,5	0.23	0
2	1PE	G	400	-	15,15,15	0.32	0	14,14,14	0.75	0
2	1PE	A	400	-	15,15,15	0.33	0	14,14,14	0.81	0
2	1PE	D	401	-	15,15,15	0.52	0	14,14,14	0.51	0
2	1PE	B	400	-	15,15,15	0.38	0	14,14,14	0.57	0
3	GOL	D	403	-	5,5,5	0.26	0	5,5,5	0.38	0
3	GOL	D	402	-	5,5,5	0.50	0	5,5,5	1.16	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1PE	F	400	-	-	10/13/13/13	-
2	1PE	E	401	-	-	8/13/13/13	-
2	1PE	C	400	-	-	5/13/13/13	-
2	1PE	H	400	-	-	8/13/13/13	-
3	GOL	E	402	-	-	4/4/4/4	-
2	1PE	G	400	-	-	6/13/13/13	-
2	1PE	A	400	-	-	8/13/13/13	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1PE	D	401	-	-	6/13/13/13	-
2	1PE	B	400	-	-	2/13/13/13	-
3	GOL	D	403	-	-	0/4/4/4	-
3	GOL	D	402	-	-	0/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	402	GOL	O3-C3-C2	2.08	119.73	110.38

There are no chirality outliers.

5 of 57 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	E	402	GOL	O1-C1-C2-C3
2	E	401	1PE	OH4-C13-C23-OH3
2	F	400	1PE	OH4-C13-C23-OH3
2	E	401	1PE	OH5-C14-C24-OH4
2	B	400	1PE	OH4-C13-C23-OH3

There are no ring outliers.

9 monomers are involved in 25 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	400	1PE	1	0
2	E	401	1PE	4	0
2	C	400	1PE	2	0
2	H	400	1PE	8	0
2	G	400	1PE	2	0
2	A	400	1PE	4	0
2	D	401	1PE	2	0
2	B	400	1PE	1	0
3	D	402	GOL	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	332/337 (98%)	-0.04	10 (3%)	52	48	13, 35, 54, 84	10 (3%)
1	B	331/337 (98%)	-0.16	2 (0%)	85	83	19, 34, 50, 60	4 (1%)
1	C	332/337 (98%)	-0.19	6 (1%)	67	63	14, 33, 50, 88	7 (2%)
1	D	332/337 (98%)	-0.09	9 (2%)	56	52	15, 34, 52, 94	10 (3%)
1	E	331/337 (98%)	-0.03	9 (2%)	56	52	17, 38, 64, 84	5 (1%)
1	F	331/337 (98%)	-0.06	10 (3%)	52	48	15, 34, 52, 88	11 (3%)
1	G	331/337 (98%)	-0.26	4 (1%)	76	73	14, 31, 45, 54	8 (2%)
1	H	331/337 (98%)	-0.12	9 (2%)	56	52	18, 37, 56, 101	1 (0%)
All	All	2651/2696 (98%)	-0.12	59 (2%)	62	58	13, 34, 54, 101	56 (2%)

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	333[A]	HIS	5.5
1	H	15	GLU	4.7
1	F	13	LEU	4.7
1	B	239[A]	TYR	4.6
1	D	17	GLN	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	E	402	6/6	0.68	0.25	67,72,72,74	0
2	1PE	E	401	16/16	0.83	0.15	37,43,66,68	0
3	GOL	D	403	6/6	0.87	0.11	62,64,64,64	0
2	1PE	H	400	16/16	0.87	0.12	37,44,52,52	0
3	GOL	D	402	6/6	0.88	0.24	35,36,43,48	0
2	1PE	A	400	16/16	0.89	0.12	35,41,58,63	0
2	1PE	F	400	16/16	0.90	0.13	37,42,59,59	0
2	1PE	G	400	16/16	0.90	0.12	32,38,47,47	0
2	1PE	C	400	16/16	0.90	0.11	37,43,56,60	0
2	1PE	D	401	16/16	0.91	0.11	42,48,58,59	0
2	1PE	B	400	16/16	0.92	0.10	35,42,55,59	0

6.5 Other polymers [i](#)

There are no such residues in this entry.