



wwPDB EM Validation Summary Report ⓘ

Mar 27, 2026 – 03:01 AM UTC

PDB ID : 6OJ5 / pdb_00006oj5
EMDB ID : EMD-20088
Title : In situ structure of rotavirus VP1 RNA-dependent RNA polymerase (TLP_RNA)
Authors : Jenni, S.; Salgado, E.N.; Herrmann, T.; Li, Z.; Grant, T.; Grigorieff, N.; Trapani, S.; Estrozi, L.F.; Harrison, S.C.
Deposited on : 2019-04-10
Resolution : 5.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	887	50% 78% 12% 10%
1	F	887	53% 81% 10% 10%
1	G	887	44% 81% 10% 10%
1	H	887	42% 78% 12% 10%
1	I	887	50% 80% 10% 10%
1	J	887	55% 78% 14% 8%
2	P	1088	53% 87% 12%

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Inner capsid protein VP2








• Molecule 1: Inner capsid protein VP2


Chain D: 50% 82% 11% 7%








• Molecule 1: Inner capsid protein VP2












• Molecule 1: Inner capsid protein VP2








• Molecule 1: Inner capsid protein VP2








• Molecule 1: Inner capsid protein VP2





• Molecule 2: RNA-directed RNA polymerase





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	82008	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI POLARA 300	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	64	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	10.245	Depositor
Minimum map value	-6.056	Depositor
Average map value	0.015	Depositor
Map value standard deviation	0.486	Depositor
Recommended contour level	2	Depositor
Map size (Å)	369.0, 369.0, 369.0	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.23, 1.23, 1.23	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.48	0/6495	0.85	0/8810
1	B	0.47	0/6495	0.84	2/8810 (0.0%)
1	C	0.47	0/6487	0.83	0/8799
1	D	0.46	0/6885	0.84	0/9332
1	E	0.48	0/6618	0.83	0/8979
1	F	0.47	0/6683	0.84	1/9064 (0.0%)
1	G	0.45	0/6683	0.83	2/9064 (0.0%)
1	H	0.46	0/6683	0.84	2/9064 (0.0%)
1	I	0.46	0/6665	0.81	0/9041
1	J	0.46	0/6798	0.82	0/9217
2	P	0.45	0/8963	0.84	2/12120 (0.0%)
All	All	0.47	0/75455	0.83	9/102300 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	2
1	F	0	1
1	G	0	3
1	H	0	1
1	I	0	3
1	J	0	3
2	P	0	2
All	All	0	18

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	P	783	TYR	CA-CB-CG	5.40	123.61	113.90
1	B	378	ASN	N-CA-C	5.25	117.07	110.24
1	G	143	ALA	CA-C-N	5.16	131.39	121.54
1	G	143	ALA	C-N-CA	5.16	131.39	121.54
2	P	84	ALA	N-CA-C	5.10	121.66	110.80

There are no chirality outliers.

5 of 18 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	371	THR	Peptide
1	B	143	ALA	Peptide
1	C	143	ALA	Peptide
1	D	102	ILE	Peptide
1	D	143	ALA	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6379	6408	6408	61	0
1	B	6379	6408	6408	67	0
1	C	6372	6400	6400	66	0
1	D	6765	6809	6809	62	0
1	E	6499	6529	6529	66	0
1	F	6563	6588	6588	53	0
1	G	6563	6588	6588	46	0
1	H	6563	6588	6588	63	0
1	I	6545	6569	6569	54	0
1	J	6678	6713	6713	70	0
2	P	8789	8862	8861	79	0
All	All	74095	74462	74461	637	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 637 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:P:359:MET:SD	2:P:614:ARG:NH2	2.61	0.73
1:F:744:GLU:O	1:F:748:THR:OG1	2.06	0.72
1:G:281:ILE:O	1:G:286:ARG:NH1	2.23	0.72
1:F:281:ILE:O	1:F:286:ARG:NH1	2.23	0.71
1:F:309:ASN:O	1:F:622:ASN:ND2	2.23	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	779/887 (88%)	712 (91%)	67 (9%)	0	100	100
1	B	779/887 (88%)	714 (92%)	65 (8%)	0	100	100
1	C	778/887 (88%)	706 (91%)	72 (9%)	0	100	100
1	D	825/887 (93%)	752 (91%)	72 (9%)	1 (0%)	48	83
1	E	793/887 (89%)	720 (91%)	73 (9%)	0	100	100
1	F	800/887 (90%)	720 (90%)	80 (10%)	0	100	100
1	G	800/887 (90%)	721 (90%)	79 (10%)	0	100	100
1	H	800/887 (90%)	722 (90%)	77 (10%)	1 (0%)	48	83
1	I	798/887 (90%)	724 (91%)	74 (9%)	0	100	100
1	J	814/887 (92%)	739 (91%)	75 (9%)	0	100	100
2	P	1084/1088 (100%)	985 (91%)	96 (9%)	3 (0%)	36	71
All	All	9050/9958 (91%)	8215 (91%)	830 (9%)	5 (0%)	49	83

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	P	82	TYR
2	P	84	ALA

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	P	942	PRO
1	H	107	PRO
1	D	103	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	717/818 (88%)	717 (100%)	0	100	100
1	B	717/818 (88%)	717 (100%)	0	100	100
1	C	716/818 (88%)	716 (100%)	0	100	100
1	D	762/818 (93%)	762 (100%)	0	100	100
1	E	731/818 (89%)	731 (100%)	0	100	100
1	F	738/818 (90%)	738 (100%)	0	100	100
1	G	738/818 (90%)	737 (100%)	1 (0%)	88	88
1	H	738/818 (90%)	738 (100%)	0	100	100
1	I	736/818 (90%)	736 (100%)	0	100	100
1	J	752/818 (92%)	752 (100%)	0	100	100
2	P	986/989 (100%)	986 (100%)	0	100	100
All	All	8331/9169 (91%)	8330 (100%)	1 (0%)	100	100

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	268	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 45 such sidechains are listed below:

Mol	Chain	Res	Type
1	H	368	GLN
2	P	51	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	H	396	GLN
1	J	267	GLN
2	P	143	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-20088. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y



Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)

6.2.1 Primary map



X Index: 150




Y Index: 150




Z Index: 150


6.2.2 Raw map



X Index: 150



Y Index: 150



Z Index: 150


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)

6.3.1 Primary map



X Index: 146




Y Index: 189




Z Index: 147


6.3.2 Raw map



X Index: 152



Y Index: 155



Z Index: 150

The images above show the largest variance slices of the map in three orthogonal directions.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 2.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency

A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_20088_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.

7.2 Volume estimate [i](#)




The volume at the recommended contour level is 523 nm³; this corresponds to an approximate mass of 473 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.192 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	5.20	-	-
Author-provided FSC curve	5.20	5.63	5.24
Unmasked-calculated*	5.34	6.00	5.40

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-20088 and PDB model 6OJ5. Per-residue inclusion information can be found in [section 3](#) on [page 5](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 2.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (2).

9.4 Atom inclusion [i](#)



At the recommended contour level, 39% of all backbone atoms, 39% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (2) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.3920	0.0760
A	0.3660	0.0520
B	0.4110	0.0810
C	0.4340	0.1090
D	0.4070	0.0890
E	0.3840	0.0650
F	0.3660	0.0470
G	0.4380	0.1050
H	0.4320	0.1090
I	0.3870	0.0660
J	0.3460	0.0360
P	0.4040	0.0770

1.0

0.0

<0.0