



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 20, 2026 – 12:01 AM UTC

PDB ID : 4OKI / pdb\_00004oki  
Title : X-ray structure of the nucleotide-binding subdomain of the enoylreductase domain of PpsC from Mycobacterium tuberculosis  
Authors : Faille, A.; Mourey, L.; Pedelacq, J.D.  
Deposited on : 2014-01-22  
Resolution : 1.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	216	 10% (Poor fit), 61% (0 outliers), 21% (1 outlier), 16% (2+ outliers)
1	B	216	 10% (Poor fit), 66% (0 outliers), 14% (1 outlier), 17% (2+ outliers)

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3033 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phthiocerol synthesis polyketide synthase type I PpsC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	181	1369	865	242	257	5	0	3	0
1	B	179	1348	855	236	253	4	0	4	0

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	636	MET	-	expression tag	UNP P96202
A	637	GLY	-	expression tag	UNP P96202
A	638	SER	-	expression tag	UNP P96202
A	639	SER	-	expression tag	UNP P96202
A	640	HIS	-	expression tag	UNP P96202
A	641	HIS	-	expression tag	UNP P96202
A	642	HIS	-	expression tag	UNP P96202
A	643	HIS	-	expression tag	UNP P96202
A	644	HIS	-	expression tag	UNP P96202
A	645	HIS	-	expression tag	UNP P96202
A	646	SER	-	expression tag	UNP P96202
A	647	SER	-	expression tag	UNP P96202
A	648	GLY	-	expression tag	UNP P96202
A	649	LEU	-	expression tag	UNP P96202
A	650	VAL	-	expression tag	UNP P96202
A	651	PRO	-	expression tag	UNP P96202
A	652	ARG	-	expression tag	UNP P96202
A	653	GLY	-	expression tag	UNP P96202
A	654	SER	-	expression tag	UNP P96202
A	655	HIS	-	expression tag	UNP P96202
A	656	MET	-	expression tag	UNP P96202
A	850	GLY	-	expression tag	UNP P96202
A	851	SER	-	expression tag	UNP P96202
B	636	MET	-	expression tag	UNP P96202
B	637	GLY	-	expression tag	UNP P96202

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	638	SER	-	expression tag	UNP P96202
B	639	SER	-	expression tag	UNP P96202
B	640	HIS	-	expression tag	UNP P96202
B	641	HIS	-	expression tag	UNP P96202
B	642	HIS	-	expression tag	UNP P96202
B	643	HIS	-	expression tag	UNP P96202
B	644	HIS	-	expression tag	UNP P96202
B	645	HIS	-	expression tag	UNP P96202
B	646	SER	-	expression tag	UNP P96202
B	647	SER	-	expression tag	UNP P96202
B	648	GLY	-	expression tag	UNP P96202
B	649	LEU	-	expression tag	UNP P96202
B	650	VAL	-	expression tag	UNP P96202
B	651	PRO	-	expression tag	UNP P96202
B	652	ARG	-	expression tag	UNP P96202
B	653	GLY	-	expression tag	UNP P96202
B	654	SER	-	expression tag	UNP P96202
B	655	HIS	-	expression tag	UNP P96202
B	656	MET	-	expression tag	UNP P96202
B	850	GLY	-	expression tag	UNP P96202
B	851	SER	-	expression tag	UNP P96202

- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	2	Total Na 2 2	0	0
2	B	5	Total Na 5 5	0	0

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 8 6 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	158	Total O 158 158	0	0
4	B	143	Total O 143 143	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phthiocerol synthesis polyketide synthase type I PpsC



- Molecule 1: Phthiocerol synthesis polyketide synthase type I PpsC



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.88Å 78.74Å 88.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.00 – 1.50 42.00 – 1.50	Depositor EDS
% Data completeness (in resolution range)	90.3 (42.00-1.50) 90.3 (42.00-1.50)	Depositor EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.22 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.182 , 0.218 0.191 , 0.224	Depositor DCC
$R_{free}$ test set	3445 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.2	Xtrriage
Anisotropy	0.119	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 36.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3033	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.07% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MPD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.75	25/1393 (1.8%)	1.46	5/1881 (0.3%)
1	B	1.66	15/1378 (1.1%)	1.38	13/1865 (0.7%)
All	All	1.71	40/2771 (1.4%)	1.42	18/3746 (0.5%)

The worst 5 of 40 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	840	HIS	CG-ND1	-7.99	1.29	1.38
1	A	693	SER	C-O	-7.39	1.18	1.24
1	B	684	HIS	ND1-CE1	7.20	1.39	1.32
1	A	842	ALA	N-CA	7.18	1.55	1.46
1	A	700	ILE	C-O	-7.14	1.16	1.24

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	749	VAL	N-CA-CB	-7.26	102.62	112.28
1	B	749	VAL	CB-CA-C	6.66	120.30	111.85
1	B	749	VAL	N-CA-C	-6.49	106.01	112.17
1	B	710	ALA	N-CA-C	-6.35	104.43	111.36
1	B	836	HIS	N-CA-C	6.13	117.62	111.07

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1369	0	1399	16	0
1	B	1348	0	1382	13	0
2	A	2	0	0	0	0
2	B	5	0	0	0	0
3	A	8	0	14	0	0
4	A	158	0	0	5	0
4	B	143	0	0	2	0
All	All	3033	0	2795	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 27 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:698:VAL:HG23	1:B:766[A]:VAL:HG13	1.36	1.03
1:A:777:GLN:HG2	4:A:1051:HOH:O	1.73	0.87
1:A:749:VAL:CG1	1:A:778:ARG:HG3	2.14	0.78
1:B:698:VAL:CG2	1:B:766[A]:VAL:HG13	2.20	0.70
1:A:749:VAL:HG11	1:A:778:ARG:HG3	1.74	0.69

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	182/216 (84%)	181 (100%)	1 (0%)	0	100	100
1	B	181/216 (84%)	179 (99%)	2 (1%)	0	100	100
All	All	363/432 (84%)	360 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	140/168 (83%)	135 (96%)	5 (4%)	31	6
1	B	139/168 (83%)	133 (96%)	6 (4%)	26	4
All	All	279/336 (83%)	268 (96%)	11 (4%)	28	5

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	753	ASP
1	B	796	ASP
1	B	832	GLN
1	B	803	LEU
1	A	840	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	826	GLN
1	B	832	GLN
1	A	826	GLN
1	A	835	GLN
1	B	701	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 7 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	MPD	A	902	-	7,7,7	0.34	0	9,10,10	0.73	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	A	902	-	-	4/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	902	MPD	C2-C3-C4-O4
3	A	902	MPD	O2-C2-C3-C4
3	A	902	MPD	C1-C2-C3-C4
3	A	902	MPD	CM-C2-C3-C4

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	181/216 (83%)	0.52	21 (11%) 9   9	12, 23, 41, 50	3 (1%)
1	B	179/216 (82%)	0.72	21 (11%) 9   9	11, 25, 53, 71	4 (2%)
All	All	360/432 (83%)	0.62	42 (11%) 9   9	11, 25, 48, 71	7 (1%)

The worst 5 of 42 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	734	MET	5.5
1	B	797	VAL	5.4
1	B	673	ALA	4.8
1	B	730	ALA	4.8
1	A	670	ASN	4.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NA	A	901	1/1	0.86	0.17	45,45,45,45	0
2	NA	B	904	1/1	0.86	0.12	47,47,47,47	0
3	MPD	A	902	8/8	0.92	0.13	40,49,53,53	0
2	NA	A	903	1/1	0.93	0.12	46,46,46,46	0
2	NA	B	903	1/1	0.93	0.11	41,41,41,41	0
2	NA	B	902	1/1	0.94	0.07	43,43,43,43	0
2	NA	B	901	1/1	0.95	0.15	40,40,40,40	0
2	NA	B	905	1/1	0.97	0.05	36,36,36,36	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.