



wwPDB EM Validation Summary Report ⓘ

Mar 5, 2026 – 07:10 PM UTC

PDB ID : 7OKX / pdb_00007okx
EMDB ID : EMD-12969
Title : Structure of active transcription elongation complex Pol II-DSIF (SPT5-KO W5)-ELL2-EAF1 (composite structure)
Authors : Chen, Y.; Vos, S.M.; Dienemann, C.; Ninov, M.; Urlaub, H.; Cramer, P.
Deposited on : 2021-05-18
Resolution : 3.30 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	15087 (2.80 - 3.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1970	
2	B	1251	
3	C	275	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	184	
5	E	210	
6	F	127	
7	G	172	
8	H	150	
9	I	125	
10	J	67	
11	K	117	
12	L	58	
13	M	640	
14	N	48	
15	O	268	
16	P	47	
17	T	48	
18	Z	1087	

2 Entry composition

There are 20 unique types of molecules in this entry. The entry contains 34834 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase II subunit RPB1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	1407	Total	C	N	O	S	0	0
			11149	7018	1998	2062	71		

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	1131	Total	C	N	O	S	0	0
			9047	5721	1592	1670	64		

- Molecule 3 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	258	Total	C	N	O	S	0	0
			2072	1300	356	410	6		

- Molecule 4 is a protein called RNA polymerase II subunit D.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	118	Total	C	N	O	S	0	0
			967	608	167	188	4		

- Molecule 5 is a protein called DNA-directed RNA polymerase II subunit E.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	209	Total	C	N	O	S	0	0
			1721	1089	300	324	8		

- Molecule 6 is a protein called DNA-directed RNA polymerase II subunit F.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	78	Total	C	N	O	S	0	0
			626	401	106	114	5		

- Molecule 7 is a protein called DNA-directed RNA polymerase II subunit RPB7.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	G	171	Total	C	N	O	S	0	0
			1347	872	218	249	8		

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	H	148	Total	C	N	O	S	0	0
			1186	750	194	237	5		

- Molecule 9 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	I	116	Total	C	N	O	S	0	0
			932	577	165	179	11		

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	J	66	Total	C	N	O	S	0	0
			524	339	88	91	6		

- Molecule 11 is a protein called RNA_pol_L_2 domain-containing protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	K	115	Total	C	N	O	S	0	0
			920	593	152	173	2		

- Molecule 12 is a protein called RNA polymerase II subunit K.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	L	44	Total	C	N	O	S	0	0
			367	228	69	64	6		

- Molecule 13 is a protein called RNA polymerase II elongation factor ELL2.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	M	139	Total	C	N	O	S	0	0
			1106	686	201	213	6		

- Molecule 14 is a DNA chain called Non-template DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	N	27	Total	C	N	O	P	0	0
			549	264	96	162	27		

- Molecule 15 is a protein called ELL-associated factor 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	O	106	Total	C	N	O	S	0	0
			832	523	149	156	4		

- Molecule 16 is a RNA chain called RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	P	13	Total	C	N	O	P	0	0
			279	125	54	87	13		

- Molecule 17 is a DNA chain called Template DNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	T	39	Total	C	N	O	P	0	0
			808	385	149	235	39		

- Molecule 18 is a protein called Transcription elongation factor SPT5.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	Z	51	Total	C	N	O	S	0	0
			393	241	72	79	1		

- Molecule 19 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
19	A	1	Total	Mg	0
			1	1	

- Molecule 20 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
20	A	2	Total	Zn	0
			2	2	
20	B	1	Total	Zn	0
			1	1	
20	C	1	Total	Zn	0
			1	1	

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
20	I	2	Total 2	Zn 2	0
20	J	1	Total 1	Zn 1	0
20	L	1	Total 1	Zn 1	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA-directed RNA polymerase II subunit RPB1




[illegible]

- Molecule 2: DNA-directed RNA polymerase subunit beta

Chain B: 75% 15% 10%





- Molecule 3: DNA-directed RNA polymerase II subunit RPB3

Chain C: 79% 15% 6%



- Molecule 4: RNA polymerase II subunit D

Chain D: 55% 9% 36%




- Molecule 5: DNA-directed RNA polymerase II subunit E

Chain E: 88% 11%



- Molecule 6: DNA-directed RNA polymerase II subunit F

Chain F: 55% 6% 39%



- Molecule 7: DNA-directed RNA polymerase II subunit RPB7

Chain G: 82% 17%



- Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC3

Chain H:  81% 18%



- Molecule 9: DNA-directed RNA polymerase II subunit RPB9

Chain I:  10% 75% 18% 7%




- Molecule 10: DNA-directed RNA polymerases I, II, and III subunit RPABC5

Chain J:  81% 18%



- Molecule 11: RNA_pol_L_2 domain-containing protein

Chain K:  92% 6%




- Molecule 12: RNA polymerase II subunit K


Chain L:  67% 9% 24%



- Molecule 13: RNA polymerase II elongation factor ELL2

Chain M:  18% 78%



[illegible]

THR
GLY
VAL
LEU
LEU
SER
ILE
ASP
GLY
GLU
ASP
GLY
ILE
VAL
ARG
MET
ASP
LEU
ASP
GLU
GLN
LEU
LYS
ILE
LEU
ASN
LEU
ARG
PHE
LEU
GLY
LYS
LEU
LEU
GLU
ALA

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	364641	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	43.21	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	56.269	Depositor
Minimum map value	-31.934	Depositor
Average map value	0.008	Depositor
Map value standard deviation	1.104	Depositor
Recommended contour level	3	Depositor
Map size (Å)	315.0, 315.0, 315.0	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.05, 1.05, 1.05	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.19	0/11350	0.43	0/15314
2	B	0.20	0/9227	0.42	0/12454
3	C	0.20	0/2115	0.41	0/2873
4	D	0.23	0/979	0.54	0/1312
5	E	0.18	0/1752	0.40	0/2366
6	F	0.19	0/636	0.40	0/859
7	G	0.27	0/1378	0.55	0/1870
8	H	0.18	0/1207	0.43	0/1628
9	I	0.21	0/954	0.57	2/1293 (0.2%)
10	J	0.30	0/533	0.43	0/719
11	K	0.19	0/939	0.36	0/1271
12	L	0.21	0/372	0.52	0/493
13	M	0.21	0/1121	0.55	0/1507
14	N	0.23	0/613	0.47	0/940
15	O	0.23	0/849	0.53	0/1145
16	P	0.11	0/312	0.19	0/484
17	T	0.24	0/907	0.43	0/1400
18	Z	0.14	0/396	0.38	0/536
All	All	0.20	0/35640	0.44	2/48464 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
7	G	0	2

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	I	20	CYS	CA-C-N	5.18	130.20	122.74
9	I	20	CYS	C-N-CA	5.18	130.20	122.74

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
7	G	117	MET	Peptide
7	G	124	ASN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11149	0	11300	155	0
2	B	9047	0	9080	132	0
3	C	2072	0	2019	32	0
4	D	967	0	973	13	0
5	E	1721	0	1737	15	0
6	F	626	0	657	6	0
7	G	1347	0	1347	18	0
8	H	1186	0	1147	17	0
9	I	932	0	856	16	0
10	J	524	0	540	11	0
11	K	920	0	942	5	0
12	L	367	0	367	4	0
13	M	1106	0	1082	17	0
14	N	549	0	308	4	0
15	O	832	0	832	14	0
16	P	279	0	142	2	0
17	T	808	0	442	12	0
18	Z	393	0	400	6	0
19	A	1	0	0	0	0
20	A	2	0	0	0	0
20	B	1	0	0	0	0
20	C	1	0	0	0	0
20	I	2	0	0	0	0
20	J	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
20	L	1	0	0	0	0
All	All	34834	0	34171	398	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 398 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:91:GLU:HG2	3:C:92:GLU:OE1	1.46	1.14
15:O:31:HIS:ND1	15:O:104:SER:OG	2.10	0.84
3:C:91:GLU:CG	3:C:92:GLU:OE1	2.29	0.80
3:C:92:GLU:HG2	3:C:93:PHE:N	1.97	0.77
2:B:134:ARG:HD3	2:B:304:ASN:HB3	1.71	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1393/1970 (71%)	1360 (98%)	33 (2%)	0	100	100
2	B	1123/1251 (90%)	1092 (97%)	31 (3%)	0	100	100
3	C	254/275 (92%)	249 (98%)	5 (2%)	0	100	100
4	D	114/184 (62%)	109 (96%)	5 (4%)	0	100	100
5	E	207/210 (99%)	204 (99%)	3 (1%)	0	100	100
6	F	76/127 (60%)	76 (100%)	0	0	100	100
7	G	169/172 (98%)	154 (91%)	15 (9%)	0	100	100
8	H	146/150 (97%)	144 (99%)	2 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
9	I	114/125 (91%)	109 (96%)	5 (4%)	0	100	100
10	J	64/67 (96%)	64 (100%)	0	0	100	100
11	K	113/117 (97%)	111 (98%)	2 (2%)	0	100	100
12	L	42/58 (72%)	41 (98%)	1 (2%)	0	100	100
13	M	131/640 (20%)	128 (98%)	3 (2%)	0	100	100
15	O	104/268 (39%)	99 (95%)	5 (5%)	0	100	100
18	Z	49/1087 (4%)	49 (100%)	0	0	100	100
All	All	4099/6701 (61%)	3989 (97%)	110 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1239/1749 (71%)	1239 (100%)	0	100	100
2	B	991/1084 (91%)	990 (100%)	1 (0%)	88	90
3	C	235/252 (93%)	235 (100%)	0	100	100
4	D	109/160 (68%)	109 (100%)	0	100	100
5	E	191/192 (100%)	191 (100%)	0	100	100
6	F	68/111 (61%)	68 (100%)	0	100	100
7	G	151/153 (99%)	151 (100%)	0	100	100
8	H	129/131 (98%)	129 (100%)	0	100	100
9	I	102/112 (91%)	102 (100%)	0	100	100
10	J	55/56 (98%)	55 (100%)	0	100	100
11	K	104/106 (98%)	104 (100%)	0	100	100
12	L	40/55 (73%)	40 (100%)	0	100	100
13	M	124/575 (22%)	124 (100%)	0	100	100
15	O	93/240 (39%)	93 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
18	Z	45/940 (5%)	45 (100%)	0	100	100
All	All	3676/5916 (62%)	3675 (100%)	1 (0%)	100	100

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	465	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
13	M	87	ASN
13	M	113	ASN
2	B	629	ASN
2	B	602	ASN
13	M	132	GLN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
16	P	12/47 (25%)	1 (8%)	0

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
16	P	36	G

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-12969. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 128



Y Index: 194




Z Index: 161


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 3.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 293 nm³; this corresponds to an approximate mass of 264 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.303 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-12969 and PDB model 7OKX. Per-residue inclusion information can be found in [section 3](#) on [page 8](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 3.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (3).

9.4 Atom inclusion [i](#)



At the recommended contour level, 94% of all backbone atoms, 89% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (3) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8890	 0.5710
A	 0.8940	 0.5890
B	 0.9100	 0.5940
C	 0.9320	 0.6140
D	 0.9610	 0.4870
E	 0.8700	 0.5660
F	 0.9320	 0.6000
G	 0.9700	 0.5240
H	 0.8900	 0.5910
I	 0.8030	 0.5190
J	 0.9530	 0.6260
K	 0.9400	 0.6140
L	 0.8580	 0.5780
M	 0.8350	 0.4680
N	 0.5430	 0.3580
O	 0.9550	 0.5210
P	 0.7710	 0.5400
T	 0.6730	 0.4140
Z	 0.6890	 0.5810

