



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:05 PM UTC

PDB ID : 3OLG / pdb_00003olg
Title : Structures of human pancreatic alpha-amylase in complex with acarviosatin III03
Authors : Qin, X.; Ren, L.
Deposited on : 2010-08-26
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	496	2% 81% 17% .
2	B	2	50% 50%
3	C	3	33% 67%
4	D	4	50% 50%
5	E	3	67% 33%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GLC	E	2	X	-	-	-

2 Entry composition [i](#)

There are 12 unique types of molecules in this entry. The entry contains 4384 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pancreatic alpha-amylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	495	Total	C	N	O	S	0	0	0
			3939	2492	695	731	21			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	287	MET	VAL	engineered mutation	UNP P04746

- Molecule 2 is an oligosaccharide called alpha-D-quinovopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	B	2	Total	C	O	0	0	0
			20	12	8			

- Molecule 3 is an oligosaccharide called alpha-D-quinovopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
3	C	3	Total	C	O	0	0	0
			32	18	14			

- Molecule 4 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
4	D	4	Total	C	O	0	0	0
			45	24	21			

- Molecule 5 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
5	E	3	Total	C	O	0	0	0
			34	18	16			


- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Cl	0	0
			1	1		

- Molecule 8 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 9 is (1S,2S,3R,6R)-6-amino-4-(hydroxymethyl)cyclohex-4-ene-1,2,3-triol (CCD ID: HSD) (formula: C₇H₁₃NO₄).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
9	A	1	Total	C	N	O	0	0
			12	7	1	4		
9	A	1	Total	C	N	O	0	0
			12	7	1	4		

- Molecule 10 is PYROGLUTAMIC ACID (CCD ID: PCA) (formula: $C_5H_7NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	A	1	Total	C	N	O	0	0
			8	5	1	2		

- Molecule 11 is alpha-D-glucopyranose (CCD ID: GLC) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	A	1	Total	C	O	0	0
			12	6	6		


- Molecule 12 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	263	Total 263	O 263	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pancreatic alpha-amylase



- Molecule 2: alpha-D-quinovopyranose-(1-4)-alpha-D-glucopyranose




- Molecule 3: alpha-D-quinovopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose



- Molecule 4: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose



- Molecule 5: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose



BGL1
GLC2
GLP3

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.38Å 74.53Å 135.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.87 – 2.30 40.87 – 2.30	Depositor EDS
% Data completeness (in resolution range)	94.6 (40.87-2.30) 94.6 (40.87-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.93 (at 2.29Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.193 , 0.216 0.191 , 0.216	Depositor DCC
R_{free} test set	1184 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	21.6	Xtriage
Anisotropy	0.138	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4384	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.57% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: G6D, CA, SO4, CL, GLC, BGC, HSD, PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.44	0/4053	0.96	19/5503 (0.3%)

There are no bond length outliers.

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	352	ASN	N-CA-C	7.58	121.85	109.72
1	A	216	ASN	N-CA-C	-6.42	101.03	110.52
1	A	195	ARG	N-CA-C	-6.36	98.37	108.73
1	A	336	THR	N-CA-C	6.27	120.50	109.96
1	A	11	THR	N-CA-C	6.25	120.95	112.88
1	A	293	LEU	N-CA-C	-5.71	97.88	107.99
1	A	55	PHE	N-CA-C	5.59	117.20	109.15
1	A	161	ARG	N-CA-C	5.51	117.68	109.25
1	A	435	SER	N-CA-C	5.36	118.22	110.28
1	A	101	HIS	N-CA-C	5.36	115.61	108.38
1	A	59	TRP	N-CA-C	5.35	119.42	113.01
1	A	92	ARG	N-CA-C	5.29	117.69	110.55
1	A	126	PHE	CA-C-N	5.17	124.83	119.56
1	A	126	PHE	C-N-CA	5.17	124.83	119.56
1	A	400	VAL	N-CA-C	5.15	115.87	110.62
1	A	95	VAL	N-CA-C	5.12	115.94	108.58
1	A	146	GLY	N-CA-C	-5.09	108.59	115.21
1	A	133	GLY	N-CA-C	5.02	118.95	112.83
1	A	185	HIS	N-CA-C	-5.01	105.71	111.07

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3939	0	3711	59	0
2	B	20	0	10	0	0
3	C	32	0	21	0	0
4	D	45	0	39	1	0
5	E	34	0	30	1	0
6	A	1	0	0	0	0
7	A	1	0	0	0	0
8	A	5	0	0	0	0
9	A	24	0	21	4	0
10	A	8	0	5	0	0
11	A	12	0	12	1	0
12	A	263	0	0	7	0
All	All	4384	0	3849	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:163:THR:HB	9:A:500:HSD:H2	1.40	0.87
1:A:150:ASN:ND2	1:A:152:ASN:H	1.95	0.65
1:A:257:LYS:O	1:A:261:LYS:HG2	1.97	0.65
1:A:264:THR:HG22	1:A:311:SER:CB	2.28	0.63
1:A:260:ALA:O	1:A:264:THR:HG23	2.04	0.58
1:A:416:GLN:HG3	1:A:432:ASP:OD2	2.04	0.57
1:A:484:GLU:HB3	12:A:1059:HOH:O	2.04	0.57
1:A:273:LYS:HA	1:A:415:ASN:HD21	1.71	0.56
1:A:393:ASN:O	1:A:396:ILE:HG23	2.05	0.56
1:A:451:ASP:OD2	1:A:454:SER:HB2	2.06	0.56
1:A:20:ARG:NH1	1:A:23:ASP:OD2	2.40	0.55
1:A:352:ASN:OD1	1:A:352:ASN:N	2.40	0.54
1:A:462:CYS:HB2	1:A:466:LYS:HD2	1.89	0.54
1:A:415:ASN:HB3	1:A:431:ASN:HB3	1.91	0.53
1:A:53:ASN:HA	1:A:54:PRO:C	2.35	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:233:GLU:OE2	9:A:503:HSD:N1	2.43	0.52
1:A:103:CYS:SG	1:A:121:PRO:HG3	2.50	0.51
1:A:351:GLY:C	1:A:352:ASN:OD1	2.54	0.51
1:A:162:LEU:HD22	1:A:201:HIS:CD2	2.45	0.51
1:A:404:GLN:HG3	1:A:424:ARG:HG3	1.93	0.51
1:A:227:LYS:HB2	12:A:965:HOH:O	2.10	0.50
12:A:990:HOH:O	5:E:1:BGC:H1	2.10	0.50
1:A:140:LYS:NZ	12:A:991:HOH:O	2.43	0.50
1:A:274:MET:HG3	12:A:572:HOH:O	2.11	0.50
1:A:393:ASN:O	1:A:396:ILE:CG2	2.60	0.49
1:A:320:LEU:HA	1:A:323:MET:HE3	1.93	0.49
1:A:301:ASN:HB2	1:A:308:GLY:HA3	1.94	0.49
1:A:393:ASN:HA	1:A:396:ILE:HG22	1.95	0.49
1:A:273:LYS:HB2	1:A:414:SER:O	2.13	0.48
4:D:1:BGC:H6C2	4:D:2:GLC:O5	2.14	0.48
1:A:150:ASN:HD22	1:A:150:ASN:C	2.22	0.47
1:A:274:MET:H	1:A:415:ASN:HD22	1.62	0.47
1:A:374:PRO:HG2	12:A:1066:HOH:O	2.14	0.47
1:A:410:TYR:C	1:A:410:TYR:CD1	2.93	0.46
1:A:396:ILE:HG23	12:A:715:HOH:O	2.15	0.46
1:A:264:THR:HG22	1:A:311:SER:OG	2.15	0.46
1:A:199:SER:CB	1:A:232:GLN:HB3	2.46	0.46
1:A:462:CYS:SG	1:A:466:LYS:HD2	2.55	0.45
1:A:163:THR:HB	9:A:500:HSD:N1	2.21	0.45
1:A:150:ASN:HD21	1:A:152:ASN:HB2	1.82	0.45
1:A:3:SER:HA	1:A:4:PRO:HD3	1.82	0.44
1:A:186:LEU:O	1:A:191:VAL:HG13	2.17	0.44
1:A:150:ASN:HD22	1:A:151:TYR:N	2.16	0.43
1:A:323:MET:HE3	1:A:323:MET:HB2	1.89	0.43
1:A:274:MET:H	1:A:415:ASN:ND2	2.17	0.42
1:A:357:TRP:CE3	1:A:358:VAL:HA	2.53	0.42
1:A:436:PHE:C	1:A:436:PHE:CD1	2.98	0.42
1:A:10:ARG:NH1	1:A:36:GLY:HA2	2.34	0.41
1:A:380:ASN:O	1:A:381:ASP:HB2	2.19	0.41
1:A:273:LYS:HD3	1:A:276:TYR:CZ	2.55	0.41
1:A:449:TYR:CE2	1:A:495:LYS:HB2	2.56	0.41
1:A:261:LYS:HA	1:A:261:LYS:HD2	1.78	0.41
1:A:273:LYS:HA	1:A:415:ASN:ND2	2.36	0.41
1:A:331:HIS:CG	1:A:332:PRO:HD2	2.56	0.41
1:A:465:ILE:CG2	1:A:466:LYS:N	2.83	0.41
1:A:300:ASP:OD2	9:A:503:HSD:O4	2.30	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:ASN:CA	1:A:54:PRO:C	2.94	0.40
1:A:388:TRP:CD1	11:A:512:GLC:H61	2.56	0.40
1:A:459:ASN:CG	1:A:459:ASN:O	2.64	0.40
1:A:458:ILE:HD12	1:A:463:THR:HG21	2.03	0.40
1:A:462:CYS:CB	1:A:466:LYS:HD2	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	493/496 (99%)	471 (96%)	21 (4%)	1 (0%)	43 55

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	308	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	418/419 (100%)	403 (96%)	15 (4%)	31 47

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	115	CYS
1	A	125	ASP
1	A	130	PRO
1	A	140	LYS
1	A	191	VAL
1	A	227	LYS
1	A	264	THR
1	A	270	ASN
1	A	273	LYS
1	A	278	LYS
1	A	341	SER
1	A	352	ASN
1	A	444	LEU
1	A	454	SER
1	A	466	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	8	GLN
1	A	15	HIS
1	A	150	ASN
1	A	152	ASN
1	A	216	ASN
1	A	270	ASN
1	A	415	ASN
1	A	476	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	B	1	2,9	11,11,12	0.54	0	15,15,17	0.92	0
2	G6D	B	2	2,9	9,9,11	0.65	0	11,12,16	2.25	5 (45%)
3	BGC	C	1	3	12,12,12	0.62	0	17,17,17	1.08	2 (11%)
3	GLC	C	2	3	11,11,12	0.55	0	15,15,17	0.78	1 (6%)
3	G6D	C	3	3,9	9,9,11	0.53	0	11,12,16	0.88	0
4	BGC	D	1	4	12,12,12	1.10	1 (8%)	17,17,17	1.90	4 (23%)
4	GLC	D	2	4	11,11,12	0.42	0	15,15,17	0.75	1 (6%)
4	GLC	D	3	4	11,11,12	0.30	0	15,15,17	0.51	0
4	GLC	D	4	4	11,11,12	0.42	0	15,15,17	0.35	0
5	BGC	E	1	5	12,12,12	1.38	3 (25%)	17,17,17	2.98	8 (47%)
5	GLC	E	2	5	11,11,12	1.19	2 (18%)	15,15,17	1.50	3 (20%)
5	GLC	E	3	5	11,11,12	0.76	0	15,15,17	1.40	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	B	1	2,9	-	0/2/19/22	0/1/1/1
2	G6D	B	2	2,9	-	-	0/1/1/1
3	BGC	C	1	3	-	2/2/22/22	0/1/1/1
3	GLC	C	2	3	-	0/2/19/22	0/1/1/1
3	G6D	C	3	3,9	-	-	0/1/1/1
4	BGC	D	1	4	-	1/2/22/22	0/1/1/1
4	GLC	D	2	4	-	0/2/19/22	0/1/1/1
4	GLC	D	3	4	-	0/2/19/22	0/1/1/1
4	GLC	D	4	4	-	0/2/19/22	0/1/1/1
5	BGC	E	1	5	-	2/2/22/22	0/1/1/1
5	GLC	E	2	5	1/1/4/5	0/2/19/22	0/1/1/1
5	GLC	E	3	5	-	0/2/19/22	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	E	1	BGC	O5-C5	3.05	1.51	1.44
5	E	1	BGC	O5-C1	2.34	1.48	1.42
4	D	1	BGC	C4-C3	2.16	1.57	1.52
5	E	1	BGC	C1-C2	2.10	1.57	1.52
5	E	2	GLC	C4-C5	-2.09	1.48	1.53
5	E	2	GLC	C1-C2	2.00	1.57	1.52

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	E	1	BGC	C4-C3-C2	-9.44	94.26	110.83
4	D	1	BGC	C1-C2-C3	-4.55	101.07	110.36
2	B	2	G6D	C3-C4-C5	4.41	118.20	111.21
2	B	2	G6D	C6-C5-C4	-3.97	107.26	113.27
5	E	3	GLC	O5-C1-C2	3.32	118.71	110.79
4	D	1	BGC	C6-C5-C4	3.27	121.04	113.02
5	E	1	BGC	C6-C5-C4	3.24	120.98	113.02
5	E	1	BGC	O4-C4-C3	3.21	117.94	110.38
5	E	3	GLC	C1-O5-C5	3.21	116.48	112.19
5	E	2	GLC	C2-C3-C4	-3.07	105.46	110.86
5	E	1	BGC	C1-C2-C3	2.99	116.46	110.36
5	E	1	BGC	C3-C4-C5	-2.88	105.00	110.23
4	D	1	BGC	C1-O5-C5	-2.86	108.12	113.65
2	B	2	G6D	O3-C3-C2	-2.78	104.11	110.20
5	E	2	GLC	C1-O5-C5	2.70	115.81	112.19
2	B	2	G6D	O5-C5-C6	-2.47	104.24	107.72
3	C	1	BGC	C3-C4-C5	2.43	114.64	110.23
3	C	1	BGC	C1-C2-C3	-2.25	105.77	110.36
5	E	1	BGC	O5-C5-C4	2.25	113.75	109.70
3	C	2	GLC	C1-C2-C3	2.25	112.92	109.64
4	D	2	GLC	C2-C3-C4	-2.16	107.06	110.86
5	E	1	BGC	C1-O5-C5	2.09	117.69	113.65
5	E	1	BGC	O5-C1-C2	2.08	113.95	110.30
2	B	2	G6D	O5-C1-C2	-2.05	105.91	110.79
4	D	1	BGC	C3-C4-C5	2.04	113.93	110.23
5	E	2	GLC	O5-C1-C2	2.02	115.61	110.79

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	E	2	GLC	C1

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	E	1	BGC	O5-C5-C6-O6
5	E	1	BGC	C4-C5-C6-O6
3	C	1	BGC	C4-C5-C6-O6
3	C	1	BGC	O5-C5-C6-O6
4	D	1	BGC	C4-C5-C6-O6


There are no ring outliers.


3 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	1	BGC	1	0
5	E	1	BGC	1	0
4	D	2	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 2 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	HSD	A	503	3,2	12,12,12	1.77	2 (16%)	11,17,17	2.87	4 (36%)
9	HSD	A	500	2	12,12,12	1.65	2 (16%)	11,17,17	3.57	7 (63%)
8	SO4	A	499	-	4,4,4	0.38	0	6,6,6	0.09	0
11	GLC	A	512	-	12,12,12	0.51	0	17,17,17	0.47	0
10	PCA	A	511	1	7,8,9	0.81	0	9,10,12	1.95	3 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	HSD	A	503	3,2	-	0/2/22/22	0/1/1/1
9	HSD	A	500	2	-	0/2/22/22	0/1/1/1
11	GLC	A	512	-	-	0/2/22/22	0/1/1/1
10	PCA	A	511	1	-	0/0/11/13	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	A	500	HSD	C2-C3	4.96	1.39	1.32
9	A	503	HSD	C2-C3	4.69	1.39	1.32
9	A	503	HSD	C5-C3	3.40	1.54	1.51
9	A	500	HSD	C5-C3	2.22	1.53	1.51

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	500	HSD	C7-C1-N1	8.08	127.82	111.40
9	A	503	HSD	C7-C1-N1	7.45	126.55	111.40
9	A	500	HSD	O4-C7-C6	5.85	124.16	110.38
9	A	503	HSD	O4-C7-C6	4.06	119.95	110.38
9	A	500	HSD	O2-C5-C3	-3.48	104.12	110.75
10	A	511	PCA	CB-CG-CD	-3.36	99.21	104.41
10	A	511	PCA	O-C-CA	-3.17	116.62	124.77
9	A	500	HSD	O3-C6-C5	3.01	115.61	109.64
9	A	503	HSD	C6-C5-C3	2.57	115.71	111.67
10	A	511	PCA	CB-CA-C	-2.47	109.27	112.66
9	A	500	HSD	O2-C5-C6	2.29	115.30	110.53
9	A	503	HSD	O4-C7-C1	2.13	113.02	109.04
9	A	500	HSD	C5-C3-C2	2.10	125.68	122.23
9	A	500	HSD	O4-C7-C1	2.08	112.92	109.04

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	A	503	HSD	2	0
9	A	500	HSD	2	0
11	A	512	GLC	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	495/496 (99%)	-0.10	10 (2%) 65 66	13, 22, 33, 45	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	484	GLU	6.2
1	A	308	GLY	5.4
1	A	307	ALA	4.5
1	A	459	ASN	2.7
1	A	350	ASN	2.7
1	A	9	GLY	2.3
1	A	454	SER	2.3
1	A	52	TYR	2.2
1	A	351	GLY	2.1
1	A	471	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GLC	E	3	11/12	0.31	0.28	20,20,20,20	0
4	BGC	D	1	12/12	0.51	0.24	20,20,20,20	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GLC	D	4	11/12	0.57	0.18	44,46,48,51	0
5	GLC	E	2	11/12	0.59	0.18	42,44,46,48	0
3	BGC	C	1	12/12	0.70	0.18	36,42,43,43	0
5	BGC	E	1	12/12	0.75	0.17	32,37,38,42	0
4	GLC	D	2	11/12	0.82	0.12	40,41,42,42	0
4	GLC	D	3	11/12	0.86	0.12	36,40,42,45	0
3	GLC	C	2	11/12	0.89	0.11	28,32,33,34	0
3	G6D	C	3	9/11	0.91	0.09	22,24,25,25	0
2	G6D	B	2	9/11	0.91	0.10	28,29,31,33	0
2	GLC	B	1	11/12	0.95	0.06	19,22,24,26	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
9	HSD	A	500	12/12	0.53	0.22	39,47,48,48	0
8	SO4	A	499	5/5	0.70	0.17	78,78,79,80	0
11	GLC	A	512	12/12	0.76	0.27	14,17,21,23	0
9	HSD	A	503	12/12	0.80	0.15	22,23,25,27	0
10	PCA	A	511	8/9	0.94	0.08	24,25,25,25	0
6	CA	A	497	1/1	0.95	0.08	29,29,29,29	0
7	CL	A	498	1/1	0.99	0.02	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.