



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 11:21 PM UTC

PDB ID : 5OLB / pdb_00005olb
Title : crystal structure of autotaxin in complex with PF-8380
Authors : Hoerer, S.; Lammens, A.
Deposited on : 2017-07-27
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	844	4% 88% 5% 7%
2	B	6	67% 33%
3	C	2	100%
4	D	6	17% 83%

2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 7550 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ectonucleotide pyrophosphatase/phosphodiesterase family member 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	785	Total	C	N	O	S	86	21	0
			6520	4128	1132	1209	51			

There are 25 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	32	ASP	-	expression tag	UNP Q9R1E6
A	33	PRO	-	expression tag	UNP Q9R1E6
A	34	SER	-	expression tag	UNP Q9R1E6
A	35	MET	-	expression tag	UNP Q9R1E6
A	?	-	LYS	deletion	UNP Q9R1E6
A	?	-	VAL	deletion	UNP Q9R1E6
A	?	-	GLU	deletion	UNP Q9R1E6
A	?	-	PRO	deletion	UNP Q9R1E6
A	859	SER	-	expression tag	UNP Q9R1E6
A	860	ARG	-	expression tag	UNP Q9R1E6
A	861	GLU	-	expression tag	UNP Q9R1E6
A	862	ASN	-	expression tag	UNP Q9R1E6
A	863	LEU	-	expression tag	UNP Q9R1E6
A	864	TYR	-	expression tag	UNP Q9R1E6
A	865	PHE	-	expression tag	UNP Q9R1E6
A	866	GLN	-	expression tag	UNP Q9R1E6
A	867	GLY	-	expression tag	UNP Q9R1E6
A	868	HIS	-	expression tag	UNP Q9R1E6
A	869	HIS	-	expression tag	UNP Q9R1E6
A	870	HIS	-	expression tag	UNP Q9R1E6
A	871	HIS	-	expression tag	UNP Q9R1E6
A	872	HIS	-	expression tag	UNP Q9R1E6
A	873	HIS	-	expression tag	UNP Q9R1E6
A	874	HIS	-	expression tag	UNP Q9R1E6
A	875	HIS	-	expression tag	UNP Q9R1E6

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)]alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	6	Total	C	N	O	0	0	0
			70	40	2	28			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	6	Total	C	N	O	0	0	0
			72	40	2	30			

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Zn	0	0
			2	2		

- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Ca	0	0
			2	2		

- Molecule 7 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	2	Total	Na	0	0
			2	2		


- Molecule 8 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	K	0	0
			1	1		

- Molecule 9 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	1	Total	Cl	0	0
			1	1		

- Molecule 10 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: C₆H₁₄O₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 11 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	A	1	Total	O	S		0	0
			5	4	1			

- Molecule 12 is (3,5-dichlorophenyl)methyl 4-[3-oxo-3-(2-oxo-2,3-dihydro-1,3-benzoxazol-6-yl)propyl]piperazine-1-carboxylate (CCD ID: 6ZO) (formula: C₂₂H₂₁Cl₂N₃O₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
12	A	1	Total	C	Cl	N	O	0	0
			32	22	2	3	5		


- Molecule 13 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
13	A	805	Total 805	O 805	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ectonucleotide pyrophosphatase/phosphodiesterase family member 2



- Molecule 2: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-3)][alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-alpha-D-mannopyranose-(1-6)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1
MAG2
BMA3
MAN4
MAN5
MAN6

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	124.73Å 100.05Å 78.78Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.65 – 1.82 26.65 – 1.82	Depositor EDS
% Data completeness (in resolution range)	95.2 (26.65-1.82) 95.2 (26.65-1.82)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.20 (at 1.82Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, R_{free}	0.172 , 0.191 0.176 , 0.190	Depositor DCC
R_{free} test set	1690 reflections (2.00%)	wwPDB-VP
Wilson B-factor (Å ²)	19.6	Xtriage
Anisotropy	0.066	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 48.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7550	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, MAN, CL, ZN, 6ZO, NAG, BMA, SO4, K, FUC, CA, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.72	0/6703	1.08	10/9091 (0.1%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	424	MET	N-CA-C	-7.66	98.10	110.20
1	A	237	PHE	CA-CB-CG	6.62	120.42	113.80
1	A	625	LEU	N-CA-C	-6.11	104.90	112.90
1	A	130	CYS	N-CA-C	-5.91	102.16	110.50
1	A	568	CYS	CA-C-N	5.38	128.23	120.38
1	A	568	CYS	C-N-CA	5.38	128.23	120.38
1	A	487	PHE	CA-CB-CG	-5.22	108.58	113.80
1	A	551	ASN	N-CA-C	-5.21	102.04	110.32
1	A	241	PHE	CA-CB-CG	5.12	118.92	113.80
1	A	110	VAL	N-CA-C	-5.03	102.25	108.89

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6520	0	6249	16	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	70	0	61	0	0
3	C	28	0	25	0	0
4	D	72	0	61	0	0
5	A	2	0	0	0	0
6	A	2	0	0	0	0
7	A	2	0	0	0	0
8	A	1	0	0	0	0
9	A	1	0	0	0	0
10	A	10	0	14	1	0
11	A	5	0	0	0	0
12	A	32	0	0	0	0
13	A	805	0	0	2	0
All	All	7550	0	6410	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:222:PRO:HA	1:A:225:HIS:CE1	2.43	0.54
1:A:273:PHE:H	10:A:923:PGE:H22	1.77	0.49
1:A:679:PRO:HD2	13:A:1744:HOH:O	2.13	0.48
1:A:765:SER:HB2	1:A:789:PHE:CZ	2.50	0.47
1:A:776:PRO:HG2	1:A:779:LYS:HB2	1.96	0.47
1:A:593:LEU:HD12	1:A:727:ASN:HB2	1.99	0.44
1:A:68:VAL:CG1	1:A:72:ASP:HB2	2.48	0.44
1:A:379:VAL:HB	1:A:455:ARG:HG2	2.00	0.44
1:A:227:ILE:HA	1:A:232[A]:MET:HE1	1.99	0.44
1:A:539:PHE:O	1:A:541:PRO:HD3	2.19	0.43
1:A:232[A]:MET:HG2	1:A:241:PHE:HB3	1.99	0.42
1:A:408:ILE:HD11	1:A:424:MET:SD	2.60	0.42
1:A:617:TYR:HA	1:A:624:PRO:HA	2.02	0.41
1:A:397:PRO:HA	13:A:1187:HOH:O	2.21	0.41
1:A:700:MET:HA	1:A:793:HIS:NE2	2.35	0.41
1:A:550:PRO:HB2	1:A:607:ILE:HG12	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	800/844 (95%)	776 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	734/768 (96%)	728 (99%)	6 (1%)	73	67

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	ASN
1	A	78	LEU
1	A	398	ASN
1	A	400	LEU
1	A	458	LEU
1	A	588	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	297	ASN
1	A	646	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	750	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

14 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	2,1	14,14,15	0.27	0	17,19,21	0.71	0
2	NAG	B	2	2	14,14,15	0.32	0	17,19,21	0.65	0
2	BMA	B	3	2	11,11,12	0.32	0	15,15,17	0.72	1 (6%)
2	MAN	B	4	2	11,11,12	0.38	0	15,15,17	0.90	1 (6%)
2	FUC	B	5	2	10,10,11	0.38	0	14,14,16	0.51	0
2	FUC	B	6	2	10,10,11	0.43	0	14,14,16	0.60	0
3	NAG	C	1	3,1	14,14,15	0.31	0	17,19,21	0.71	0
3	NAG	C	2	3	14,14,15	0.32	0	17,19,21	0.48	0
4	NAG	D	1	1,4	14,14,15	0.31	0	17,19,21	0.89	1 (5%)
4	NAG	D	2	4	14,14,15	0.31	0	17,19,21	0.60	0
4	BMA	D	3	4	11,11,12	0.53	0	15,15,17	2.25	1 (6%)
4	MAN	D	4	4	11,11,12	0.45	0	15,15,17	0.94	1 (6%)
4	MAN	D	5	4	11,11,12	0.40	0	15,15,17	0.69	1 (6%)
4	MAN	D	6	4	11,11,12	0.41	0	15,15,17	0.83	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	0/2/19/22	0/1/1/1
2	MAN	B	4	2	-	0/2/19/22	0/1/1/1
2	FUC	B	5	2	-	-	0/1/1/1
2	FUC	B	6	2	-	-	0/1/1/1
3	NAG	C	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	C	2	3	-	2/6/23/26	0/1/1/1
4	NAG	D	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	D	2	4	-	0/6/23/26	0/1/1/1
4	BMA	D	3	4	-	0/2/19/22	0/1/1/1
4	MAN	D	4	4	-	0/2/19/22	0/1/1/1
4	MAN	D	5	4	-	0/2/19/22	0/1/1/1
4	MAN	D	6	4	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	3	BMA	C1-O5-C5	8.27	123.28	112.19
4	D	4	MAN	C1-O5-C5	3.10	116.34	112.19
2	B	4	MAN	C1-O5-C5	2.97	116.17	112.19
4	D	6	MAN	C1-O5-C5	2.75	115.88	112.19
4	D	1	NAG	O5-C1-C2	-2.58	107.30	111.29
2	B	3	BMA	C1-O5-C5	2.47	115.50	112.19
4	D	5	MAN	C1-O5-C5	2.13	115.04	112.19

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	2	NAG	C4-C5-C6-O6
3	C	2	NAG	O5-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry

Of 11 ligands modelled in this entry, 8 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
11	SO4	A	924	-	4,4,4	0.38	0	6,6,6	0.10	0
10	PGE	A	923	-	9,9,9	0.15	0	8,8,8	0.16	0
12	6ZO	A	925	5	35,35,35	1.22	1 (2%)	49,49,49	2.20	6 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	PGE	A	923	-	-	5/7/7/7	-
12	6ZO	A	925	5	-	0/18/28/28	0/4/4/4

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	A	925	6ZO	O32-C30	6.22	1.33	1.21

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	A	925	6ZO	C28-N31-C30	-9.85	104.59	109.90
12	A	925	6ZO	O29-C30-N31	8.67	113.33	108.20
12	A	925	6ZO	O29-C30-O32	-3.55	117.14	121.98
12	A	925	6ZO	C26-O29-C30	-3.44	104.94	107.00
12	A	925	6ZO	C26-C28-N31	3.42	108.83	105.61
12	A	925	6ZO	C17-N16-C15	2.54	118.02	111.24

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
10	A	923	PGE	O2-C3-C4-O3
10	A	923	PGE	O3-C5-C6-O4
10	A	923	PGE	C3-C4-O3-C5
10	A	923	PGE	C6-C5-O3-C4
10	A	923	PGE	C4-C3-O2-C2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	A	923	PGE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	785/844 (93%)	-0.20	30 (3%)	44 50	8, 20, 41, 75	51 (6%)

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	69	GLY	6.0
1	A	68	VAL	5.5
1	A	570	ASP	4.7
1	A	71	PRO	4.0
1	A	81	SER	3.8
1	A	397	PRO	3.7
1	A	569	ASP	3.6
1	A	459	ASP	3.5
1	A	77	ASN	3.4
1	A	78	LEU	3.2
1	A	585	GLY	3.1
1	A	264[A]	THR	3.1
1	A	680	TYR	3.0
1	A	855	GLU	3.0
1	A	70	PRO	3.0
1	A	638	SER	2.7
1	A	67	GLU	2.6
1	A	398	ASN	2.6
1	A	51	TRP	2.6
1	A	72	ASP	2.5
1	A	458	LEU	2.5
1	A	586	SER	2.5
1	A	320	PHE	2.4
1	A	278	SER	2.3
1	A	469	PHE	2.3
1	A	584	LYS	2.2
1	A	274	PHE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	396	ILE	2.1
1	A	277[A]	VAL	2.1
1	A	378	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	BMA	D	3	11/12	0.63	0.16	52,57,60,60	0
2	BMA	B	3	11/12	0.69	0.17	49,51,55,56	0
4	MAN	D	5	11/12	0.70	0.18	54,55,59,59	0
4	MAN	D	4	11/12	0.72	0.16	57,60,61,61	0
4	MAN	D	6	11/12	0.75	0.17	46,49,51,51	0
3	NAG	C	2	14/15	0.78	0.14	56,61,63,64	0
2	MAN	B	4	11/12	0.82	0.16	42,47,49,49	0
3	NAG	C	1	14/15	0.86	0.11	34,40,47,51	0
4	NAG	D	2	14/15	0.87	0.10	24,28,35,44	0
2	FUC	B	5	10/11	0.91	0.09	30,33,35,36	0
2	FUC	B	6	10/11	0.92	0.10	37,38,41,41	0
2	NAG	B	2	14/15	0.92	0.10	29,41,45,47	0
2	NAG	B	1	14/15	0.95	0.08	26,29,34,35	0
4	NAG	D	1	14/15	0.97	0.06	14,16,20,20	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain B:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	CA	A	920	1/1	0.81	0.12	61,61,61,61	0
10	PGE	A	923	10/10	0.83	0.14	41,47,48,49	0
7	NA	A	922	1/1	0.85	0.19	37,37,37,37	1
11	SO4	A	924	5/5	0.93	0.10	34,34,36,39	5
12	6ZO	A	925	32/32	0.93	0.10	28,38,43,44	0
7	NA	A	918	1/1	0.97	0.13	24,24,24,24	0
8	K	A	919	1/1	1.00	0.01	13,13,13,13	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
9	CL	A	921	1/1	1.00	0.04	22,22,22,22	0
5	ZN	A	915	1/1	1.00	0.01	18,18,18,18	0
5	ZN	A	916	1/1	1.00	0.02	19,19,19,19	0
6	CA	A	917	1/1	1.00	0.01	16,16,16,16	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.