



## wwPDB EM Validation Summary Report ⓘ

Mar 8, 2026 – 12:11 PM UTC

PDB ID : 8OL1 / pdb\_00008ol1  
EMDB ID : EMD-16936  
Title : cGAS-Nucleosome in complex with SPSB3-ELOBC (composite structure)  
Authors : Xu, P.B.; Ablasser, A.  
Deposited on : 2023-03-29  
Resolution : 3.50 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132  
MolProbity : 4-5-2 with Phenix2.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



| Metric                | Whole archive<br>(#Entries) | EM structures<br>(#Entries) | Similar EM resolution<br>(#Entries, resolution range(Å)) |
|-----------------------|-----------------------------|-----------------------------|----------------------------------------------------------|
| Clashscore            | 229148                      | 23984                       | -                                                        |
| Ramachandran outliers | 224038                      | 23583                       | -                                                        |
| Sidechain outliers    | 223484                      | 23102                       | -                                                        |
| Q-score               | -                           | 25397                       | 13950 ( 3.00 - 4.00 )                                    |

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 1   | A     | 98     | 93% 6% .         |
| 1   | E     | 98     | 88% 10% .        |
| 2   | B     | 81     | 84% 15% .        |
| 2   | F     | 81     | 86% 10% .        |

Continued on next page...

*Continued from previous page...*

| Mol | Chain | Length | Quality of chain |
|-----|-------|--------|------------------|
| 3   | C     | 108    |                  |
| 4   | D     | 95     |                  |
| 5   | G     | 107    |                  |
| 6   | H     | 94     |                  |
| 7   | I     | 145    |                  |
| 8   | J     | 145    |                  |
| 9   | K     | 362    |                  |
| 10  | L     | 244    |                  |
| 11  | M     | 112    |                  |
| 12  | N     | 118    |                  |

## 2 Entry composition

There are 13 unique types of molecules in this entry. The entry contains 18376 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.2.

| Mol | Chain | Residues | Atoms |     |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 1   | A     | 98       | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 810   | 511 | 157 | 140 | 2 |         |       |
| 1   | E     | 98       | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 810   | 511 | 157 | 140 | 2 |         |       |

- Molecule 2 is a protein called Histone H4.

| Mol | Chain | Residues | Atoms |     |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 2   | B     | 80       | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 638   | 401 | 125 | 111 | 1 |         |       |
| 2   | F     | 81       | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 646   | 407 | 126 | 112 | 1 |         |       |

- Molecule 3 is a protein called Histone H2A type 1-H.

| Mol | Chain | Residues | Atoms |     |     |     | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---------|-------|
| 3   | C     | 108      | Total | C   | N   | O   | 0       | 0     |
|     |       |          | 829   | 523 | 162 | 144 |         |       |

- Molecule 4 is a protein called Histone H2B type 1-H.

| Mol | Chain | Residues | Atoms |     |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 4   | D     | 95       | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 746   | 467 | 136 | 141 | 2 |         |       |

- Molecule 5 is a protein called Histone H2A type 1-J.

| Mol | Chain | Residues | Atoms |     |     |     | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---------|-------|
| 5   | G     | 107      | Total | C   | N   | O   | 0       | 0     |
|     |       |          | 822   | 518 | 161 | 143 |         |       |

- Molecule 6 is a protein called Histone H2B type 1-N.

| Mol | Chain | Residues | Atoms |     |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 6   | H     | 94       | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 737   | 461 | 134 | 140 | 2 |         |       |

- Molecule 7 is a DNA chain called DNA (145-MER).

| Mol | Chain | Residues | Atoms |      |     |     |     | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|-----|---------|-------|
| 7   | I     | 145      | Total | C    | N   | O   | P   | 0       | 0     |
|     |       |          | 2954  | 1404 | 537 | 869 | 144 |         |       |

- Molecule 8 is a DNA chain called DNA (145-MER).

| Mol | Chain | Residues | Atoms |      |     |     |     | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|-----|---------|-------|
| 8   | J     | 145      | Total | C    | N   | O   | P   | 0       | 0     |
|     |       |          | 2985  | 1414 | 560 | 867 | 144 |         |       |

- Molecule 9 is a protein called Cyclic GMP-AMP synthase.

| Mol | Chain | Residues | Atoms |      |     |     |    | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 9   | K     | 362      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 2966  | 1895 | 510 | 546 | 15 |         |       |

There are 3 discrepancies between the modelled and reference sequences:

| Chain | Residue | Modelled | Actual | Comment             | Reference  |
|-------|---------|----------|--------|---------------------|------------|
| K     | 285     | ALA      | LYS    | engineered mutation | UNP Q8N884 |
| K     | 300     | ALA      | ARG    | engineered mutation | UNP Q8N884 |
| K     | 428     | ALA      | LYS    | engineered mutation | UNP Q8N884 |

- Molecule 10 is a protein called SPRY domain-containing SOCS box protein 3.

| Mol | Chain | Residues | Atoms |      |     |     |    | AltConf | Trace |
|-----|-------|----------|-------|------|-----|-----|----|---------|-------|
| 10  | L     | 221      | Total | C    | N   | O   | S  | 0       | 0     |
|     |       |          | 1737  | 1093 | 306 | 323 | 15 |         |       |

- Molecule 11 is a protein called Elongin-C.

| Mol | Chain | Residues | Atoms |     |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 11  | M     | 112      | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 873   | 553 | 139 | 173 | 8 |         |       |

- Molecule 12 is a protein called Elongin-B.

| Mol | Chain | Residues | Atoms |     |     |     |   | AltConf | Trace |
|-----|-------|----------|-------|-----|-----|-----|---|---------|-------|
| 12  | N     | 104      | Total | C   | N   | O   | S | 0       | 0     |
|     |       |          | 822   | 520 | 138 | 159 | 5 |         |       |


- Molecule 13 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

| Mol | Chain | Residues | Atoms |    | AltConf |
|-----|-------|----------|-------|----|---------|
| 13  | K     | 1        | Total | Zn | 0       |
|     |       |          | 1     | 1  |         |

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histone H3.2

Chain A:  93% 6% .




- Molecule 1: Histone H3.2

Chain E:  88% 10% .



- Molecule 2: Histone H4

Chain B:  84% 15% .



- Molecule 2: Histone H4

Chain F:  86% 10% .





- Molecule 3: Histone H2A type 1-H


Chain C:  6% 82% 17% .




- Molecule 4: Histone H2B type 1-H




- Molecule 5: Histone H2A type 1-J




- Molecule 6: Histone H2B type 1-N



- Molecule 7: DNA (145-MER)



- Molecule 8: DNA (145-MER)




- Molecule 9: Cyclic GMP-AMP synthase





- Molecule 10: SPRY domain-containing SOCS box protein 3







• Molecule 11: Elongin-C



• Molecule 12: Elongin-B



## 4 Experimental information

| Property                             | Value                     | Source    |
|--------------------------------------|---------------------------|-----------|
| EM reconstruction method             | SINGLE PARTICLE           | Depositor |
| Imposed symmetry                     | POINT, Not provided       |           |
| Number of particles used             | 592494                    | Depositor |
| Resolution determination method      | FSC 0.143 CUT-OFF         | Depositor |
| CTF correction method                | NONE                      | Depositor |
| Microscope                           | FEI TITAN KRIOS           | Depositor |
| Voltage (kV)                         | 300                       | Depositor |
| Electron dose ( $e^-/\text{\AA}^2$ ) | 40                        | Depositor |
| Minimum defocus (nm)                 | 600                       | Depositor |
| Maximum defocus (nm)                 | 2000                      | Depositor |
| Magnification                        | Not provided              |           |
| Image detector                       | FEI FALCON IV (4k x 4k)   | Depositor |
| Maximum map value                    | 37.819                    | Depositor |
| Minimum map value                    | -13.291                   | Depositor |
| Average map value                    | -0.017                    | Depositor |
| Map value standard deviation         | 0.929                     | Depositor |
| Recommended contour level            | 6                         | Depositor |
| Map size ( $\text{\AA}$ )            | 365.184, 365.184, 365.184 | wwPDB     |
| Map dimensions                       | 288, 288, 288             | wwPDB     |
| Map angles ( $^\circ$ )              | 90.0, 90.0, 90.0          | wwPDB     |
| Pixel spacing ( $\text{\AA}$ )       | 1.268, 1.268, 1.268       | Depositor |

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

| Mol | Chain | Bond lengths |             | Bond angles |                 |
|-----|-------|--------------|-------------|-------------|-----------------|
|     |       | RMSZ         | $\# Z  > 5$ | RMSZ        | $\# Z  > 5$     |
| 1   | A     | 0.84         | 0/822       | 1.43        | 4/1101 (0.4%)   |
| 1   | E     | 0.86         | 0/822       | 1.47        | 3/1101 (0.3%)   |
| 2   | B     | 0.90         | 0/645       | 1.51        | 2/862 (0.2%)    |
| 2   | F     | 0.89         | 0/653       | 1.44        | 4/873 (0.5%)    |
| 3   | C     | 0.91         | 0/839       | 1.62        | 4/1131 (0.4%)   |
| 4   | D     | 0.86         | 0/757       | 1.58        | 10/1015 (1.0%)  |
| 5   | G     | 0.83         | 0/831       | 1.48        | 3/1119 (0.3%)   |
| 6   | H     | 0.77         | 0/748       | 1.40        | 1/1004 (0.1%)   |
| 7   | I     | 0.70         | 0/3310      | 1.35        | 2/5103 (0.0%)   |
| 8   | J     | 0.71         | 0/3352      | 1.34        | 7/5176 (0.1%)   |
| 9   | K     | 0.70         | 0/3026      | 1.31        | 6/4060 (0.1%)   |
| 10  | L     | 0.75         | 0/1779      | 1.39        | 5/2401 (0.2%)   |
| 11  | M     | 0.80         | 0/892       | 1.35        | 1/1204 (0.1%)   |
| 12  | N     | 0.83         | 0/838       | 1.42        | 1/1132 (0.1%)   |
| All | All   | 0.77         | 0/19314     | 1.39        | 53/27282 (0.2%) |

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 1   | A     | 0                   | 1                   |
| 1   | E     | 0                   | 2                   |
| 2   | F     | 0                   | 1                   |
| 3   | C     | 0                   | 3                   |
| 4   | D     | 0                   | 2                   |
| 5   | G     | 0                   | 2                   |
| 7   | I     | 0                   | 49                  |
| 8   | J     | 0                   | 52                  |
| 9   | K     | 0                   | 2                   |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | #Chirality outliers | #Planarity outliers |
|-----|-------|---------------------|---------------------|
| 10  | L     | 0                   | 3                   |
| 11  | M     | 0                   | 1                   |
| 12  | N     | 0                   | 3                   |
| All | All   | 0                   | 121                 |

There are no bond length outliers.

The worst 5 of 53 bond angle outliers are listed below:

| Mol | Chain | Res | Type | Atoms     | Z     | Observed(°) | Ideal(°) |
|-----|-------|-----|------|-----------|-------|-------------|----------|
| 2   | F     | 61  | PHE  | CA-CB-CG  | 7.52  | 121.32      | 113.80   |
| 9   | K     | 236 | ARG  | NE-CZ-NH2 | 6.40  | 124.96      | 119.20   |
| 9   | K     | 217 | HIS  | CA-CB-CG  | -6.22 | 107.58      | 113.80   |
| 1   | A     | 116 | ARG  | NE-CZ-NH2 | 6.21  | 124.79      | 119.20   |
| 1   | A     | 134 | ARG  | NE-CZ-NH2 | 6.19  | 124.77      | 119.20   |

There are no chirality outliers.

5 of 121 planarity outliers are listed below:

| Mol | Chain | Res | Type | Group     |
|-----|-------|-----|------|-----------|
| 1   | A     | 40  | ARG  | Sidechain |
| 3   | C     | 32  | ARG  | Sidechain |
| 3   | C     | 35  | ARG  | Sidechain |
| 3   | C     | 57  | TYR  | Sidechain |
| 4   | D     | 37  | TYR  | Sidechain |

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 1   | A     | 810   | 0        | 848      | 0       | 0            |
| 1   | E     | 810   | 0        | 848      | 5       | 0            |
| 2   | B     | 638   | 0        | 676      | 1       | 0            |
| 2   | F     | 646   | 0        | 687      | 4       | 0            |
| 3   | C     | 829   | 0        | 889      | 6       | 0            |
| 4   | D     | 746   | 0        | 769      | 3       | 0            |
| 5   | G     | 822   | 0        | 882      | 0       | 0            |

*Continued on next page...*

*Continued from previous page...*

| Mol | Chain | Non-H | H(model) | H(added) | Clashes | Symm-Clashes |
|-----|-------|-------|----------|----------|---------|--------------|
| 6   | H     | 737   | 0        | 756      | 0       | 0            |
| 7   | I     | 2954  | 0        | 1627     | 0       | 0            |
| 8   | J     | 2985  | 0        | 1628     | 1       | 0            |
| 9   | K     | 2966  | 0        | 2994     | 0       | 0            |
| 10  | L     | 1737  | 0        | 1697     | 1       | 0            |
| 11  | M     | 873   | 0        | 845      | 1       | 0            |
| 12  | N     | 822   | 0        | 824      | 2       | 0            |
| 13  | K     | 1     | 0        | 0        | 0       | 0            |
| All | All   | 18376 | 0        | 15970    | 18      | 0            |

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 18 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

| Atom-1          | Atom-2           | Interatomic distance (Å) | Clash overlap (Å) |
|-----------------|------------------|--------------------------|-------------------|
| 3:C:39:TYR:CG   | 4:D:74:ALA:HB1   | 2.32                     | 0.64              |
| 3:C:63:LEU:O    | 3:C:66:ALA:HB3   | 2.11                     | 0.51              |
| 3:C:108:LEU:H   | 3:C:108:LEU:HD23 | 1.75                     | 0.51              |
| 1:E:99:TYR:CD2  | 2:F:61:PHE:CE2   | 3.00                     | 0.49              |
| 3:C:65:LEU:HD13 | 3:C:93:LEU:HD21  | 1.94                     | 0.49              |

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed    | Favoured | Allowed | Outliers | Percentiles |     |
|-----|-------|-------------|----------|---------|----------|-------------|-----|
| 1   | A     | 96/98 (98%) | 90 (94%) | 6 (6%)  | 0        | 100         | 100 |
| 1   | E     | 96/98 (98%) | 87 (91%) | 9 (9%)  | 0        | 100         | 100 |
| 2   | B     | 78/81 (96%) | 71 (91%) | 5 (6%)  | 2 (3%)   | 4           | 27  |

*Continued on next page...*

Continued from previous page...

| Mol | Chain | Analysed        | Favoured   | Allowed  | Outliers | Percentiles |     |
|-----|-------|-----------------|------------|----------|----------|-------------|-----|
| 2   | F     | 79/81 (98%)     | 77 (98%)   | 2 (2%)   | 0        | 100         | 100 |
| 3   | C     | 106/108 (98%)   | 93 (88%)   | 11 (10%) | 2 (2%)   | 6           | 33  |
| 4   | D     | 93/95 (98%)     | 86 (92%)   | 4 (4%)   | 3 (3%)   | 3           | 24  |
| 5   | G     | 105/107 (98%)   | 101 (96%)  | 4 (4%)   | 0        | 100         | 100 |
| 6   | H     | 92/94 (98%)     | 84 (91%)   | 7 (8%)   | 1 (1%)   | 11          | 43  |
| 9   | K     | 360/362 (99%)   | 331 (92%)  | 24 (7%)  | 5 (1%)   | 9           | 38  |
| 10  | L     | 219/244 (90%)   | 204 (93%)  | 12 (6%)  | 3 (1%)   | 9           | 38  |
| 11  | M     | 110/112 (98%)   | 102 (93%)  | 8 (7%)   | 0        | 100         | 100 |
| 12  | N     | 102/118 (86%)   | 90 (88%)   | 9 (9%)   | 3 (3%)   | 3           | 26  |
| All | All   | 1536/1598 (96%) | 1416 (92%) | 101 (7%) | 19 (1%)  | 13          | 41  |

5 of 19 Ramachandran outliers are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 2   | B     | 24  | ASP  |
| 4   | D     | 91  | SER  |
| 9   | K     | 261 | PRO  |
| 2   | B     | 25  | ASN  |
| 9   | K     | 235 | PRO  |

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

| Mol | Chain | Analysed     | Rotameric | Outliers | Percentiles |    |
|-----|-------|--------------|-----------|----------|-------------|----|
| 1   | A     | 85/86 (99%)  | 82 (96%)  | 3 (4%)   | 32          | 57 |
| 1   | E     | 85/86 (99%)  | 78 (92%)  | 7 (8%)   | 10          | 34 |
| 2   | B     | 65/66 (98%)  | 59 (91%)  | 6 (9%)   | 8           | 32 |
| 2   | F     | 66/66 (100%) | 59 (89%)  | 7 (11%)  | 6           | 27 |
| 3   | C     | 83/83 (100%) | 78 (94%)  | 5 (6%)   | 17          | 44 |
| 4   | D     | 82/82 (100%) | 77 (94%)  | 5 (6%)   | 17          | 43 |
| 5   | G     | 82/82 (100%) | 78 (95%)  | 4 (5%)   | 22          | 48 |

Continued on next page...

*Continued from previous page...*

| Mol | Chain | Analysed        | Rotameric  | Outliers | Percentiles |    |
|-----|-------|-----------------|------------|----------|-------------|----|
| 6   | H     | 81/81 (100%)    | 75 (93%)   | 6 (7%)   | 13          | 37 |
| 9   | K     | 331/331 (100%)  | 316 (96%)  | 15 (4%)  | 24          | 50 |
| 10  | L     | 192/211 (91%)   | 188 (98%)  | 4 (2%)   | 47          | 66 |
| 11  | M     | 96/96 (100%)    | 92 (96%)   | 4 (4%)   | 26          | 52 |
| 12  | N     | 92/103 (89%)    | 88 (96%)   | 4 (4%)   | 26          | 51 |
| All | All   | 1340/1373 (98%) | 1270 (95%) | 70 (5%)  | 22          | 47 |

5 of 70 residues with a non-rotameric sidechain are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 9   | K     | 487 | GLU  |
| 10  | L     | 307 | GLN  |
| 11  | M     | 105 | MET  |
| 1   | E     | 131 | ARG  |
| 1   | E     | 129 | ARG  |

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

| Mol | Chain | Res | Type |
|-----|-------|-----|------|
| 9   | K     | 449 | ASN  |
| 10  | L     | 111 | ASN  |
| 10  | L     | 311 | ASN  |
| 10  | L     | 236 | ASN  |
| 4   | D     | 67  | ASN  |

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.




## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-16936. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map



X Index: 144



Y Index: 144




Z Index: 144


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 144



Y Index: 137




Z Index: 138


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views

This section was not generated.


## 6.6 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 266 nm<sup>3</sup>; this corresponds to an approximate mass of 241 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum ⓘ



\*Reported resolution corresponds to spatial frequency of 0.286 Å<sup>-1</sup>

## 8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-16936 and PDB model 8OL1. Per-residue inclusion information can be found in section 3 on page 7.

### 9.1 Map-model overlay [i](#)


This section was not generated.

### 9.2 Q-score mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


### 9.3 Atom inclusion mapped to coordinate model [i](#)





The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (6).

## 9.4 Atom inclusion [i](#)



At the recommended contour level, 93% of all backbone atoms, 90% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (6) and Q-score for the entire model and for each chain.

| Chain | Atom inclusion                                                                            | Q-score                                                                                   |
|-------|-------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------|
| All   |  0.8980  |  0.4000  |
| A     |  0.9730  |  0.5200  |
| B     |  0.9770  |  0.5110  |
| C     |  0.7960  |  0.3770  |
| D     |  0.7670  |  0.3400  |
| E     |  0.9140  |  0.4890  |
| F     |  0.9400  |  0.4990  |
| G     |  0.9360  |  0.5040  |
| H     |  0.9570  |  0.4970  |
| I     |  0.9620  |  0.4490  |
| J     |  0.9750  |  0.4500  |
| K     |  0.9710  |  0.3650  |
| L     |  0.9280  |  0.2880  |
| M     |  0.4370  |  0.1750  |
| N     |  0.4880 |  0.1610 |

