



wwPDB EM Validation Summary Report ⓘ

Mar 26, 2026 – 07:38 PM UTC

PDB ID : 9OLO / pdb_00009olo
EMDB ID : EMD-70598
Title : 22bin20S complex (NSF-alphaSNAP-2:2 syntaxin-1a:SNAP-25), hydrolyzing, class 19
Authors : White, K.I.; Brunger, A.T.
Deposited on : 2025-05-12
Resolution : 3.56 Å (reported)
Based on initial models : 1JTH, 6MDM

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	12750 (3.06 - 4.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	747	
1	B	747	
1	C	747	
1	D	747	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	747	 64%8%28%
1	F	747	 59%10%31%
2	G	267	 12%15%9%76%
2	H	267	 13%22%7%71%
3	I	222	 13%21%7%72%
3	J	222	 12%21%74%
4	K	296	 82%86%12%.
4	L	296	 77%84%13%.
4	M	296	 75%88%10%.
4	N	296	 83%84%13%.

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 80286 atoms, of which 40215 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vesicle-fusing ATPase.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	A	534	Total	C	H	N	O	S	15	0
			8629	2683	4385	746	792	23		
1	B	728	Total	C	H	N	O	S	12	0
			11687	3645	5919	1014	1078	31		
1	C	536	Total	C	H	N	O	S	11	0
			8621	2681	4379	748	790	23		
1	D	728	Total	C	H	N	O	S	12	0
			11639	3628	5893	1011	1076	31		
1	E	535	Total	C	H	N	O	S	4	0
			8504	2648	4317	733	783	23		
1	F	518	Total	C	H	N	O	S	3	0
			8259	2576	4196	709	757	21		

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P18708
A	-1	ALA	-	expression tag	UNP P18708
A	0	ASN	-	expression tag	UNP P18708
B	-2	GLY	-	expression tag	UNP P18708
B	-1	ALA	-	expression tag	UNP P18708
B	0	ASN	-	expression tag	UNP P18708
C	-2	GLY	-	expression tag	UNP P18708
C	-1	ALA	-	expression tag	UNP P18708
C	0	ASN	-	expression tag	UNP P18708
D	-2	GLY	-	expression tag	UNP P18708
D	-1	ALA	-	expression tag	UNP P18708
D	0	ASN	-	expression tag	UNP P18708
E	-2	GLY	-	expression tag	UNP P18708
E	-1	ALA	-	expression tag	UNP P18708
E	0	ASN	-	expression tag	UNP P18708
F	-2	GLY	-	expression tag	UNP P18708
F	-1	ALA	-	expression tag	UNP P18708
F	0	ASN	-	expression tag	UNP P18708

- Molecule 2 is a protein called Syntaxin-1A.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	G	64	Total	C	H	N	O	S	0	0
			1038	325	512	90	106	5		
2	H	78	Total	C	H	N	O	S	0	0
			1258	392	626	106	128	6		

- Molecule 3 is a protein called Synaptosomal-associated protein 25.

Mol	Chain	Residues	Atoms						AltConf	Trace
3	I	62	Total	C	H	N	O	S	0	0
			981	297	491	85	104	4		
3	J	57	Total	C	H	N	O	S	0	0
			914	274	458	80	98	4		

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	-15	MET	-	expression tag	UNP P60881
I	-14	GLY	-	expression tag	UNP P60881
I	-13	SER	-	expression tag	UNP P60881
I	-12	SER	-	expression tag	UNP P60881
I	-11	HIS	-	expression tag	UNP P60881
I	-10	HIS	-	expression tag	UNP P60881
I	-9	HIS	-	expression tag	UNP P60881
I	-8	HIS	-	expression tag	UNP P60881
I	-7	HIS	-	expression tag	UNP P60881
I	-6	HIS	-	expression tag	UNP P60881
I	-5	SER	-	expression tag	UNP P60881
I	-4	GLN	-	expression tag	UNP P60881
I	-3	ASP	-	expression tag	UNP P60881
I	-2	PRO	-	expression tag	UNP P60881
I	-1	ASN	-	expression tag	UNP P60881
I	0	SER	-	expression tag	UNP P60881
I	85	ALA	CYS	conflict	UNP P60881
I	88	ALA	CYS	conflict	UNP P60881
I	90	ALA	CYS	conflict	UNP P60881
I	92	ALA	CYS	conflict	UNP P60881
J	-15	MET	-	expression tag	UNP P60881
J	-14	GLY	-	expression tag	UNP P60881
J	-13	SER	-	expression tag	UNP P60881
J	-12	SER	-	expression tag	UNP P60881
J	-11	HIS	-	expression tag	UNP P60881

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
J	-10	HIS	-	expression tag	UNP P60881
J	-9	HIS	-	expression tag	UNP P60881
J	-8	HIS	-	expression tag	UNP P60881
J	-7	HIS	-	expression tag	UNP P60881
J	-6	HIS	-	expression tag	UNP P60881
J	-5	SER	-	expression tag	UNP P60881
J	-4	GLN	-	expression tag	UNP P60881
J	-3	ASP	-	expression tag	UNP P60881
J	-2	PRO	-	expression tag	UNP P60881
J	-1	ASN	-	expression tag	UNP P60881
J	0	SER	-	expression tag	UNP P60881
J	85	ALA	CYS	conflict	UNP P60881
J	88	ALA	CYS	conflict	UNP P60881
J	90	ALA	CYS	conflict	UNP P60881
J	92	ALA	CYS	conflict	UNP P60881


- Molecule 4 is a protein called Alpha-soluble NSF attachment protein.

Mol	Chain	Residues	Atoms						AltConf	Trace
4	K	289	Total	C	H	N	O	S	0	0
			4503	1435	2231	378	441	18		
4	L	287	Total	C	H	N	O	S	2	0
			4485	1429	2221	377	440	18		
4	M	289	Total	C	H	N	O	S	0	0
			4503	1435	2231	378	441	18		
4	N	287	Total	C	H	N	O	S	0	0
			4467	1424	2212	375	438	18		

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
K	0	GLY	-	expression tag	UNP P54921
L	0	GLY	-	expression tag	UNP P54921
M	0	GLY	-	expression tag	UNP P54921
N	0	GLY	-	expression tag	UNP P54921

- Molecule 5 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						AltConf
5	A	1	Total 39	C 10	H 12	N 5	O 10	P 2	0
5	B	1	Total 39	C 10	H 12	N 5	O 10	P 2	0
5	C	1	Total 39	C 10	H 12	N 5	O 10	P 2	0
5	D	1	Total 39	C 10	H 12	N 5	O 10	P 2	0
5	F	1	Total 39	C 10	H 12	N 5	O 10	P 2	0

- Molecule 6 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						AltConf
6	A	1	Total 43	C 10	H 12	N 5	O 13	P 3	0
6	B	1	Total 43	C 10	H 12	N 5	O 13	P 3	0
6	C	1	Total 43	C 10	H 12	N 5	O 13	P 3	0
6	D	1	Total 43	C 10	H 12	N 5	O 13	P 3	0
6	E	1	Total 43	C 10	H 12	N 5	O 13	P 3	0
6	E	1	Total 43	C 10	H 12	N 5	O 13	P 3	0
6	F	1	Total 43	C 10	H 12	N 5	O 13	P 3	0

- Molecule 7 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms	AltConf
7	A	1	Total Mg 1 1	0
7	B	2	Total Mg 2 2	0
7	C	2	Total Mg 2 2	0
7	D	2	Total Mg 2 2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
7	E	2	Total	Mg	0
			2	2	
7	F	1	Total	Mg	0
			1	1	

- Molecule 8 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf
8	B	1	Total	O	P	0
			5	4	1	
8	C	1	Total	O	P	0
			5	4	1	
8	D	1	Total	O	P	0
			5	4	1	


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		AltConf
9	A	48	Total	O	0
			48	48	
9	B	67	Total	O	0
			67	67	
9	C	76	Total	O	0
			76	76	
9	D	32	Total	O	0
			32	32	

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
9	E	32	Total 32	O 32	0
9	F	22	Total 22	O 22	0




• Molecule 1: Vesicle-fusing ATPase

Chain C: 66% 6% 28%




• Molecule 1: Vesicle-fusing ATPase

Chain D: 24% 88% 10%




• Molecule 1: Vesicle-fusing ATPase

Chain E:  64% 8% 28%




- Molecule 1: Vesicle-fusing ATPase

Chain F:  59% 10% 31%





- Molecule 2: Syntaxin-1A



Chain G:  12% 15% 9% 76%




• Molecule 4: Alpha-soluble NSF attachment protein





• Molecule 4: Alpha-soluble NSF attachment protein







• Molecule 4: Alpha-soluble NSF attachment protein



• Molecule 4: Alpha-soluble NSF attachment protein



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	104003	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	33.960	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	3000	Depositor
Magnification	22500	Depositor
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	2.994	Depositor
Minimum map value	-1.637	Depositor
Average map value	0.008	Depositor
Map value standard deviation	0.084	Depositor
Recommended contour level	0.3	Depositor
Map size (Å)	322.224, 322.224, 322.224	wwPDB
Map dimensions	294, 294, 294	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.096, 1.096, 1.096	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, MG, ADP, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.13	0/4357	0.34	0/5872
1	B	0.18	0/5876	0.33	0/7913
1	C	0.15	0/4345	0.34	0/5852
1	D	0.19	0/5878	0.37	0/7915
1	E	0.16	0/4271	0.35	0/5756
1	F	0.27	0/4133	0.44	0/5568
2	G	0.15	0/532	0.36	0/712
2	H	0.16	0/638	0.39	0/854
3	I	0.14	0/490	0.25	0/650
3	J	0.14	0/455	0.30	0/603
4	K	0.09	0/2312	0.22	0/3107
4	L	0.09	0/2313	0.22	0/3107
4	M	0.09	0/2312	0.24	0/3107
4	N	0.09	0/2295	0.23	0/3084
All	All	0.17	0/40207	0.34	0/54100

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4244	4385	4315	31	0
1	B	5768	5919	5899	64	0
1	C	4242	4379	4336	33	0
1	D	5746	5893	5851	52	0
1	E	4187	4317	4300	40	0
1	F	4063	4196	4185	58	0
2	G	526	512	511	24	0
2	H	632	626	626	19	0
3	I	490	491	490	18	0
3	J	456	458	457	13	0
4	K	2272	2231	2231	28	0
4	L	2264	2221	2207	27	0
4	M	2272	2231	2231	21	0
4	N	2255	2212	2212	32	0
5	A	27	12	12	0	0
5	B	27	12	12	0	0
5	C	27	12	12	0	0
5	D	27	12	12	0	0
5	F	27	12	12	0	0
6	A	31	12	12	0	0
6	B	31	12	12	0	0
6	C	31	12	12	0	0
6	D	31	12	12	0	0
6	E	62	24	24	0	0
6	F	31	12	12	0	0
7	A	1	0	0	0	0
7	B	2	0	0	0	0
7	C	2	0	0	0	0
7	D	2	0	0	0	0
7	E	2	0	0	0	0
7	F	1	0	0	0	0
8	B	5	0	0	0	0
8	C	5	0	0	0	0
8	D	5	0	0	0	0
9	A	48	0	0	0	0
9	B	67	0	0	1	0
9	C	76	0	0	1	0
9	D	32	0	0	1	0
9	E	32	0	0	0	0
9	F	22	0	0	0	0
All	All	40071	40215	39995	422	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 422 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:414:MET:SD	1:B:449:GLN:NE2	2.53	0.81
4:K:210:ALA:HB1	4:K:244:MET:HE1	1.65	0.79
1:B:46:ILE:HD12	1:B:174:VAL:HG21	1.66	0.78
3:I:69:LYS:NZ	4:K:114:MET:SD	2.52	0.77
4:K:54:MET:SD	4:L:112:THR:OG1	2.41	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	547/747 (73%)	519 (95%)	28 (5%)	0	100	100
1	B	736/747 (98%)	707 (96%)	29 (4%)	0	100	100
1	C	545/747 (73%)	519 (95%)	26 (5%)	0	100	100
1	D	736/747 (98%)	701 (95%)	34 (5%)	1 (0%)	48	79
1	E	537/747 (72%)	503 (94%)	34 (6%)	0	100	100
1	F	515/747 (69%)	486 (94%)	27 (5%)	2 (0%)	30	61
2	G	62/267 (23%)	62 (100%)	0	0	100	100
2	H	76/267 (28%)	73 (96%)	3 (4%)	0	100	100
3	I	60/222 (27%)	60 (100%)	0	0	100	100
3	J	55/222 (25%)	55 (100%)	0	0	100	100
4	K	287/296 (97%)	281 (98%)	6 (2%)	0	100	100
4	L	287/296 (97%)	280 (98%)	7 (2%)	0	100	100
4	M	287/296 (97%)	278 (97%)	9 (3%)	0	100	100
4	N	285/296 (96%)	283 (99%)	2 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	5015/6644 (76%)	4807 (96%)	205 (4%)	3 (0%)	49 79

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	390	GLU
1	D	213	TRP
1	F	399	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	471/638 (74%)	467 (99%)	4 (1%)	73 77
1	B	634/638 (99%)	629 (99%)	5 (1%)	73 77
1	C	467/638 (73%)	466 (100%)	1 (0%)	87 85
1	D	636/638 (100%)	631 (99%)	5 (1%)	73 77
1	E	460/638 (72%)	457 (99%)	3 (1%)	76 78
1	F	446/638 (70%)	446 (100%)	0	100 100
2	G	59/245 (24%)	59 (100%)	0	100 100
2	H	72/245 (29%)	72 (100%)	0	100 100
3	I	54/187 (29%)	54 (100%)	0	100 100
3	J	51/187 (27%)	50 (98%)	1 (2%)	48 66
4	K	237/243 (98%)	237 (100%)	0	100 100
4	L	237/243 (98%)	236 (100%)	1 (0%)	84 80
4	M	237/243 (98%)	237 (100%)	0	100 100
4	N	235/243 (97%)	235 (100%)	0	100 100
All	All	4296/5664 (76%)	4276 (100%)	20 (0%)	81 80

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	408[B]	HIS
1	E	617[B]	ARG
4	L	43	GLU
3	J	49	MET
1	B	434[A]	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	128	GLN
1	F	454	ASN
1	E	353	GLN
4	K	67	GLN
1	B	170	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 25 ligands modelled in this entry, 10 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	ATP	F	801	7	32,33,33	0.32	0	48,52,52	0.69	0
5	ADP	A	801	-	28,29,29	1.39	4 (14%)	43,45,45	1.86	9 (20%)
5	ADP	F	802	-	28,29,29	1.40	4 (14%)	43,45,45	1.91	11 (25%)
8	PO4	C	805	7	4,4,4	0.76	0	6,6,6	0.46	0
8	PO4	B	805	7	4,4,4	0.96	0	6,6,6	0.47	0
6	ATP	C	801	7	32,33,33	0.32	0	48,52,52	0.69	1 (2%)
6	ATP	B	801	7	32,33,33	0.34	0	48,52,52	0.68	0
5	ADP	C	802	-	28,29,29	0.60	0	43,45,45	0.74	2 (4%)
6	ATP	D	801	7	32,33,33	0.36	0	48,52,52	0.69	1 (2%)
6	ATP	A	802	7	32,33,33	0.41	0	48,52,52	0.69	0
6	ATP	E	801	7	32,33,33	0.59	0	48,52,52	0.59	0
8	PO4	D	805	-	4,4,4	0.87	0	6,6,6	0.46	0
5	ADP	D	802	-	28,29,29	0.53	0	43,45,45	0.47	0
6	ATP	E	802	7	32,33,33	0.28	0	48,52,52	0.69	0
5	ADP	B	802	7	28,29,29	1.40	4 (14%)	43,45,45	1.88	11 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ATP	F	801	7	-	4/22/38/38	0/3/3/3
5	ADP	A	801	-	-	2/16/32/32	0/3/3/3
5	ADP	F	802	-	-	5/16/32/32	0/3/3/3
6	ATP	C	801	7	-	5/22/38/38	0/3/3/3
6	ATP	B	801	7	-	9/22/38/38	0/3/3/3
5	ADP	C	802	-	-	6/16/32/32	0/3/3/3
6	ATP	D	801	7	-	8/22/38/38	0/3/3/3
6	ATP	A	802	7	-	7/22/38/38	0/3/3/3
6	ATP	E	801	7	-	4/22/38/38	0/3/3/3
5	ADP	D	802	-	-	2/16/32/32	0/3/3/3
6	ATP	E	802	7	-	6/22/38/38	0/3/3/3
5	ADP	B	802	7	-	4/16/32/32	0/3/3/3

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	F	802	ADP	C5-C4	4.68	1.47	1.39
5	A	801	ADP	C5-C4	4.54	1.47	1.39
5	B	802	ADP	C5-C4	4.52	1.47	1.39
5	F	802	ADP	C5-C6	2.67	1.48	1.41
5	B	802	ADP	C5-C6	2.60	1.48	1.41

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	802	ADP	C5-C4-N3	-5.90	118.59	126.72
5	A	801	ADP	C5-C4-N3	-5.88	118.62	126.72
5	B	802	ADP	C5-C4-N3	-5.79	118.74	126.72
5	F	802	ADP	N3-C4-N9	4.81	135.34	127.17
5	A	801	ADP	N3-C4-N9	4.81	135.34	127.17

There are no chirality outliers.

5 of 62 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	A	801	ADP	C5'-O5'-PA-O1A
5	A	801	ADP	C5'-O5'-PA-O3A
5	B	802	ADP	C5'-O5'-PA-O3A
5	C	802	ADP	C5'-O5'-PA-O1A
5	C	802	ADP	C5'-O5'-PA-O2A

There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand ATP F 801





Ligand ADP A 801


































5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-70598. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 147




Y Index: 147




Z Index: 147


6.2.2 Raw map



X Index: 147



Y Index: 147




Z Index: 147


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 169




Y Index: 150




Z Index: 147


6.3.2 Raw map



X Index: 153



Y Index: 162




Z Index: 143


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.3. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 339 nm^3 ; this corresponds to an approximate mass of 306 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.281 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.281 $\text{\AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.56	-	-
Author-provided FSC curve	3.56	4.06	3.66
Unmasked-calculated*	4.23	6.68	4.31

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.23 differs from the reported value 3.56 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-70598 and PDB model 9OLO. Per-residue inclusion information can be found in section 3 on page 11.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.3 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.3).

9.4 Atom inclusion [i](#)



At the recommended contour level, 67% of all backbone atoms, 66% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.3) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.6650	0.3510
A	0.9240	0.4860
B	0.7250	0.4130
C	0.9440	0.5090
D	0.7340	0.4140
E	0.9170	0.4690
F	0.8490	0.3980
G	0.3790	0.1650
H	0.4590	0.2100
I	0.4230	0.1700
J	0.3640	0.1180
K	0.1960	0.0940
L	0.2230	0.1010
M	0.2240	0.1020
N	0.1730	0.1080

1.0

0.0

<0.0