



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:30 PM UTC

PDB ID : 2OMA / pdb_00002oma
Title : Crystallographic analysis of a chemically modified triosephosphate isomerase from *Trypanosoma cruzi* with dithiobenzylamine (DTBA)
Authors : Rodriguez-Romero, A.; Gomez-Puyou, A.
Deposited on : 2007-01-21
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	250	% 78% 20% .
1	B	250	3% 76% 21% .

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4210 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Triosephosphate isomerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	250	Total	C	N	O	S	0	3	0
			1935	1229	341	357	8			
1	B	250	Total	C	N	O	S	0	3	0
			1932	1228	340	357	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	118	CS1	CYS	modified residue	UNP P52270
B	118	CS1	CYS	modified residue	UNP P52270

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	O S	0	0
			5	4 1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 7 4 3	0	0
3	A	1	Total C O 7 4 3	0	0
3	B	1	Total C O 7 4 3	0	0

- Molecule 4 is TRIETHYLENE GLYCOL (CCD ID: PGE) (formula: $C_6H_{14}O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			10	6	4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	125	Total	O	0	0
			125	125		
5	B	142	Total	O	0	0
			142	142		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Triosephosphate isomerase



• Molecule 1: Triosephosphate isomerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	43.21Å 75.34Å 146.43Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.45 – 2.15 41.45 – 2.15	Depositor EDS
% Data completeness (in resolution range)	91.0 (41.45-2.15) 91.0 (41.45-2.15)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.70 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.196 , 0.250 0.196 , 0.250	Depositor DCC
R_{free} test set	1248 reflections (4.66%)	wwPDB-VP
Wilson B-factor (Å ²)	24.0	Xtriage
Anisotropy	0.054	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 41.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4210	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, PGE, CS1, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.22	2/1971 (0.1%)	1.26	9/2676 (0.3%)
1	B	1.30	9/1968 (0.5%)	1.25	11/2674 (0.4%)
All	All	1.26	11/3939 (0.3%)	1.26	20/5350 (0.4%)

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	49	ILE	CA-CB	8.43	1.59	1.53
1	B	79	VAL	CA-CB	6.34	1.62	1.54
1	B	151	VAL	CA-CB	6.12	1.61	1.54
1	B	114	VAL	CA-CB	5.78	1.61	1.54
1	B	173	ILE	CA-CB	5.72	1.60	1.54
1	B	125	ILE	CA-CB	5.67	1.60	1.53
1	B	203	ILE	CA-CB	5.43	1.60	1.54
1	B	92	VAL	CA-CB	5.24	1.62	1.54
1	B	3	SER	CA-C	5.24	1.59	1.52
1	A	123	HIS	CA-C	-5.18	1.46	1.52
1	B	124	VAL	CA-CB	5.03	1.60	1.54

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	17	GLY	N-CA-C	8.13	120.63	110.96
1	A	18	SER	N-CA-C	-7.61	96.96	108.52
1	B	18	SER	N-CA-C	-6.54	98.78	108.79
1	A	232	PHE	N-CA-C	6.41	119.17	109.23
1	A	180	THR	N-CA-C	-5.88	101.67	109.84
1	B	151	VAL	N-CA-C	5.83	116.01	110.53
1	A	160	TRP	N-CA-C	5.79	119.44	112.38
1	B	168	GLU	CA-C-N	-5.76	112.63	119.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	168	GLU	C-N-CA	-5.76	112.63	119.84
1	B	116	GLN	N-CA-C	5.52	117.30	111.28
1	B	61	PHE	N-CA-C	5.44	118.69	109.76
1	B	151	VAL	N-CA-CB	5.26	117.31	110.57
1	A	76	THR	N-CA-CB	5.21	117.73	109.71
1	B	236	GLY	N-CA-C	5.18	118.94	112.73
1	B	49	ILE	CA-C-N	5.16	124.60	119.24
1	B	49	ILE	C-N-CA	5.16	124.60	119.24
1	B	22	LEU	N-CA-C	5.15	117.80	111.82
1	A	65	ALA	N-CA-C	-5.09	102.52	110.42
1	A	211	TYR	CA-C-N	5.09	125.11	121.65
1	A	211	TYR	C-N-CA	5.09	125.11	121.65

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1935	0	1962	39	0
1	B	1932	0	1955	39	1
2	A	25	0	0	1	0
2	B	20	0	0	0	0
3	A	14	0	20	0	0
3	B	7	0	10	0	0
4	B	10	0	14	1	0
5	A	125	0	0	7	0
5	B	142	0	0	5	1
All	All	4210	0	3961	69	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:245:GLU:HG2	5:B:1114:HOH:O	1.50	1.09
1:A:75:PHE:HB3	1:B:15:CYS:HB3	1.43	0.98
1:A:19:GLU:HG2	1:A:51[A]:MET:HE1	1.59	0.82
1:A:101:LEU:CD1	5:A:1055:HOH:O	2.37	0.72
1:A:51[B]:MET:SD	1:A:55:ARG:HD2	2.31	0.71
1:A:14:LYS:HD3	1:B:74:ALA:HA	1.71	0.70
1:B:23:VAL:HB	1:B:24:PRO:HD3	1.74	0.68
1:A:19:GLU:CG	1:A:51[A]:MET:HE1	2.23	0.68
1:B:23:VAL:O	1:B:27:GLU:HG3	1.94	0.68
1:A:11:ALA:HB1	1:A:238:SER:HB2	1.78	0.66
1:A:221:ARG:HD3	5:A:1108:HOH:O	1.96	0.65
1:A:169:PRO:HG2	1:A:172:ALA:HB3	1.82	0.61
1:A:50:PRO:HG2	1:B:50:PRO:HG2	1.83	0.59
1:A:25:LEU:HD11	5:A:1121:HOH:O	2.02	0.59
1:A:168:GLU:HG2	1:A:212:GLY:HA3	1.86	0.58
1:A:202:ASP:O	1:A:206:GLN:HG2	2.04	0.57
4:B:800:PGE:H6	5:B:1095:HOH:O	2.04	0.57
1:A:141:ALA:HA	1:A:190:LEU:HD21	1.87	0.56
1:A:4:LYS:NZ	1:A:224:TYR:O	2.39	0.56
1:B:12:ASN:HB2	1:B:233[A]:LEU:HD11	1.87	0.55
1:B:14:LYS:O	1:B:66:GLN:NE2	2.33	0.55
1:B:30:ASN:ND2	1:B:57:THR:H	2.05	0.55
1:A:11:ALA:CB	1:A:238:SER:HB2	2.37	0.54
1:A:108:GLU:H	1:A:108:GLU:CD	2.16	0.53
1:B:4:LYS:NZ	1:B:224:TYR:O	2.41	0.53
1:A:75:PHE:CB	1:B:15:CYS:HB3	2.30	0.52
1:A:221:ARG:NH2	1:A:251:LYS:HB3	2.25	0.52
1:A:83:ILE:HD12	1:B:15:CYS:SG	2.50	0.51
1:B:13:TRP:O	1:B:14:LYS:C	2.55	0.50
1:A:75:PHE:HD2	1:B:15:CYS:SG	2.35	0.50
1:A:82[B]:GLN:NE2	5:A:1130:HOH:O	2.44	0.50
1:B:30:ASN:HD21	1:B:57:THR:H	1.60	0.50
1:A:50:PRO:HB3	1:A:87:TYR:OH	2.12	0.49
1:B:14:LYS:HE3	5:B:1138:HOH:O	2.12	0.49
1:A:75:PHE:HD2	1:B:15:CYS:HG	1.61	0.49
1:B:15:CYS:HA	1:B:45:THR:CG2	2.43	0.48
1:B:30:ASN:HD21	1:B:56:LEU:HD12	1.78	0.48
1:A:8:ILE:HA	1:A:39:GLN:O	2.13	0.48
1:B:233[A]:LEU:HD12	5:B:1054:HOH:O	2.14	0.48
1:A:106:THR:O	1:A:110:VAL:HG23	2.15	0.47
1:B:132:ASN:O	1:B:136:GLU:HG2	2.15	0.46
1:B:189:GLU:OE2	1:B:193:ARG:NH2	2.49	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:15:CYS:SG	1:B:69:ILE:HD11	2.56	0.46
1:A:168:GLU:OE1	5:A:1132:HOH:O	2.20	0.45
1:B:202:ASP:OD1	1:B:203:ILE:HG12	2.16	0.45
1:A:83:ILE:HG22	1:B:47:LEU:HD12	1.99	0.45
1:A:17:GLY:N	5:A:1090:HOH:O	2.50	0.44
1:B:49:ILE:N	1:B:50:PRO:CD	2.80	0.44
1:B:51:MET:O	1:B:55:ARG:HB2	2.17	0.44
1:A:14:LYS:HB3	1:A:15:CYS:H	1.71	0.44
1:B:8:ILE:HB	1:B:210:LEU:HD21	2.00	0.43
1:A:74:ALA:HB3	1:B:103:TYR:CE2	2.53	0.43
1:A:221:ARG:HH22	1:A:251:LYS:C	2.26	0.43
1:B:233[A]:LEU:CD1	5:B:1054:HOH:O	2.67	0.43
1:B:15:CYS:HA	1:B:45:THR:HG22	2.00	0.42
1:A:178:VAL:HG23	1:A:213:GLY:O	2.20	0.42
1:B:21:LEU:O	1:B:24:PRO:HD2	2.19	0.42
1:B:42:VAL:HG13	1:B:61:PHE:CE2	2.55	0.41
1:A:8:ILE:HB	1:A:210:LEU:HD21	2.03	0.41
1:A:208:ARG:HD3	1:A:208:ARG:N	2.36	0.41
1:A:41:VAL:HA	1:A:62:GLN:O	2.21	0.41
1:A:193:ARG:NH2	2:A:1002:SO4:O3	2.50	0.41
1:B:11:ALA:CB	1:B:238:SER:HB2	2.50	0.41
1:B:81:LEU:HD13	1:B:117:ALA:HA	2.03	0.41
1:A:222:THR:HG22	5:A:1069:HOH:O	2.20	0.40
1:B:53:LYS:HA	1:B:63:ILE:CD1	2.52	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:225[A]:GLN:OE1	5:B:1044:HOH:O[3_644]	2.17	0.03

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	250/250 (100%)	245 (98%)	4 (2%)	1 (0%)	30	26
1	B	250/250 (100%)	240 (96%)	9 (4%)	1 (0%)	30	26
All	All	500/500 (100%)	485 (97%)	13 (3%)	2 (0%)	30	26

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	14	LYS
1	A	14	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	201/198 (102%)	197 (98%)	4 (2%)	48	54
1	B	200/198 (101%)	195 (98%)	5 (2%)	42	45
All	All	401/396 (101%)	392 (98%)	9 (2%)	44	51

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	LEU
1	A	33	THR
1	A	158	GLU
1	A	221	ARG
1	B	20	SER
1	B	47	LEU
1	B	62	GLN
1	B	128	VAL
1	B	151	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	16	ASN
1	B	30	ASN
1	B	82	GLN
1	B	153	GLN
1	B	206	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	CS1	B	118	1	12,14,15	1.64	1 (8%)	12,17,19	0.84	0
1	CS1	A	118	1	12,14,15	1.44	3 (25%)	12,17,19	0.98	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CS1	B	118	1	-	0/5/8/10	0/1/1/1
1	CS1	A	118	1	-	1/5/8/10	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	118	CS1	CE-SD	-5.07	1.71	1.79
1	A	118	CS1	CE-SD	-3.43	1.74	1.79
1	A	118	CS1	O-C	2.13	1.28	1.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	118	CS1	CA-N	-2.03	1.42	1.48

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	118	CS1	C5-CE-SD-SG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

13 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	1003	-	4,4,4	0.23	0	6,6,6	0.48	0
2	SO4	A	1007	-	4,4,4	0.15	0	6,6,6	0.78	0
2	SO4	B	1006	-	4,4,4	0.31	0	6,6,6	0.47	0
2	SO4	A	1005	-	4,4,4	0.52	0	6,6,6	0.60	0
3	PEG	A	700	-	6,6,6	0.52	0	5,5,5	0.48	0
2	SO4	B	1004	-	4,4,4	0.31	0	6,6,6	1.04	1 (16%)
3	PEG	B	702	-	6,6,6	0.87	0	5,5,5	0.46	0
2	SO4	B	1008	-	4,4,4	0.57	0	6,6,6	0.30	0
2	SO4	B	1009	-	4,4,4	0.40	0	6,6,6	0.46	0
2	SO4	A	1001	-	4,4,4	0.39	0	6,6,6	0.61	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	1002	-	4,4,4	0.41	0	6,6,6	0.34	0
4	PGE	B	800	-	9,9,9	0.63	0	8,8,8	0.32	0
3	PEG	A	701	-	6,6,6	0.56	0	5,5,5	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PEG	A	700	-	-	2/4/4/4	-
3	PEG	B	702	-	-	2/4/4/4	-
4	PGE	B	800	-	-	3/7/7/7	-
3	PEG	A	701	-	-	1/4/4/4	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1004	SO4	O3-S-O2	-2.01	99.05	109.56

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	702	PEG	O2-C3-C4-O4
3	A	700	PEG	O1-C1-C2-O2
3	A	700	PEG	O2-C3-C4-O4
3	B	702	PEG	O1-C1-C2-O2
3	A	701	PEG	O2-C3-C4-O4
4	B	800	PGE	O2-C3-C4-O3
4	B	800	PGE	C1-C2-O2-C3
4	B	800	PGE	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1002	SO4	1	0
4	B	800	PGE	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	249/250 (99%)	-0.06	3 (1%) 76 79	12, 20, 32, 37	5 (2%)
1	B	249/250 (99%)	-0.10	7 (2%) 55 59	10, 17, 30, 46	4 (1%)
All	All	498/500 (99%)	-0.08	10 (2%) 65 69	10, 18, 31, 46	9 (1%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	15	CYS	4.7
1	B	20	SER	3.4
1	A	2	ALA	3.4
1	B	19[A]	GLU	3.0
1	B	17	GLY	2.6
1	B	2	ALA	2.5
1	A	15	CYS	2.2
1	A	174	GLY	2.2
1	B	74	ALA	2.1
1	B	14	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CS1	A	118	14/15	0.95	0.08	18,28,31,31	0
1	CS1	B	118	14/15	0.96	0.06	16,23,27,27	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	SO4	A	1005	5/5	0.74	0.17	53,53,56,56	0
3	PEG	A	700	7/7	0.82	0.13	43,44,48,50	0
3	PEG	A	701	7/7	0.83	0.16	46,47,49,49	0
2	SO4	B	1008	5/5	0.88	0.14	26,27,30,31	5
2	SO4	B	1006	5/5	0.89	0.18	27,27,30,31	5
2	SO4	A	1003	5/5	0.89	0.14	41,44,47,48	0
2	SO4	A	1007	5/5	0.90	0.12	27,28,29,29	5
3	PEG	B	702	7/7	0.90	0.11	25,27,28,28	0
4	PGE	B	800	10/10	0.90	0.09	27,30,36,37	0
2	SO4	A	1001	5/5	0.92	0.12	41,42,44,45	0
2	SO4	B	1009	5/5	0.92	0.15	25,27,28,29	5
2	SO4	A	1002	5/5	0.92	0.15	39,42,46,47	0
2	SO4	B	1004	5/5	0.93	0.09	36,39,46,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.