



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 08:47 AM UTC

PDB ID : 3OMK / pdb_00003omk
Title : Crystal structure of human FXR in complex with (2S)-2-[2-(4-chlorophenyl)-5,6-difluoro-1H-benzimidazol-1-yl]-2-cyclohexyl-N-(2-methylphenyl)ethanamide
Authors : Rudolph, M.G.
Deposited on : 2010-08-27
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	233	3% 83% 14% ..
1	C	233	5% 78% 20% .
2	B	14	7% 79% 7% 14%
2	D	14	7% 64% 14% 21%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4256 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bile acid receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	230	Total	C	N	O	S	0	2	0
			1894	1212	319	351	12			
1	C	230	Total	C	N	O	S	0	1	0
			1888	1208	319	349	12			


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	244	GLY	-	expression tag	UNP Q96RI1
A	245	SER	-	expression tag	UNP Q96RI1
A	246	HIS	-	expression tag	UNP Q96RI1
A	247	MET	-	expression tag	UNP Q96RI1
A	281	ALA	GLU	engineered mutation	UNP Q96RI1
A	354	ALA	GLU	engineered mutation	UNP Q96RI1
C	244	GLY	-	expression tag	UNP Q96RI1
C	245	SER	-	expression tag	UNP Q96RI1
C	246	HIS	-	expression tag	UNP Q96RI1
C	247	MET	-	expression tag	UNP Q96RI1
C	281	ALA	GLU	engineered mutation	UNP Q96RI1
C	354	ALA	GLU	engineered mutation	UNP Q96RI1

- Molecule 2 is a protein called peptide of Nuclear receptor coactivator 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	12	Total	C	N	O	0	0	0
			104	67	19	18			
2	D	11	Total	C	N	O	0	0	0
			96	63	18	15			

- Molecule 3 is (2S)-2-[2-(4-chlorophenyl)-5,6-difluoro-1H-benzimidazol-1-yl]-2-cyclohexyl-N-(2-methylphenyl)ethanamide (CCD ID: OMK) (formula: C₂₈H₂₆ClF₂N₃O).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	Cl	F	N	O	0	0
			35	28	1	2	3	1		
3	C	1	Total	C	Cl	F	N	O	0	0
			35	28	1	2	3	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	122	Total	O	0	0
			122	122		
4	B	4	Total	O	0	0
			4	4		
4	C	76	Total	O	0	0
			76	76		
4	D	2	Total	O	0	0
			2	2		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Bile acid receptor




- Molecule 1: Bile acid receptor



- Molecule 2: peptide of Nuclear receptor coactivator 1



- Molecule 2: peptide of Nuclear receptor coactivator 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	72.32Å 84.71Å 190.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.68 – 1.90 38.68 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.2 (38.68-1.90) 98.4 (38.68-1.90)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 1.89Å)	Xtriage
Refinement program	BUSTER 2.9.5	Depositor
R, R_{free}	0.204 , 0.245 0.217 , 0.262	Depositor DCC
R_{free} test set	2305 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	32.6	Xtriage
Anisotropy	0.255	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 39.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4256	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OMK

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/1941	1.24	3/2622 (0.1%)
1	C	0.79	0/1932	1.26	4/2610 (0.2%)
2	B	0.68	0/105	1.25	1/140 (0.7%)
2	D	0.66	0/97	1.07	0/129
All	All	0.81	0/4075	1.24	8/5501 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	426	HIS	N-CA-C	6.73	118.70	111.36
1	A	290	ILE	N-CA-C	-6.03	104.63	110.72
1	C	349	SER	CA-C-N	5.99	128.30	120.28
1	C	349	SER	C-N-CA	5.99	128.30	120.28
1	A	475	VAL	N-CA-C	-5.95	100.11	108.80
2	B	747	GLN	N-CA-C	5.68	117.47	111.28
1	C	273	ILE	N-CA-C	5.37	116.09	110.62
1	A	472	ILE	N-CA-C	5.13	116.59	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1894	0	1896	24	0
1	C	1888	0	1890	22	0
2	B	104	0	103	0	0
2	D	96	0	99	3	0
3	A	35	0	26	0	0
3	C	35	0	26	1	0
4	A	122	0	0	0	0
4	B	4	0	0	0	0
4	C	76	0	0	1	0
4	D	2	0	0	0	0
All	All	4256	0	4040	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:399:ARG:HB3	1:A:402:ILE:HD13	1.60	0.83
1:C:273:ILE:HD12	1:C:294:MET:HE2	1.66	0.76
1:C:269:MET:SD	1:C:270:PRO:HD2	2.27	0.74
1:A:370:PHE:O	1:A:374:LYS:HG2	1.94	0.68
1:A:462:ASP:HB2	1:A:464:LYS:NZ	2.10	0.67
1:A:273:ILE:HD13	1:A:294:MET:HE1	1.77	0.67
1:A:399:ARG:HB3	1:A:402:ILE:CD1	2.25	0.66
1:C:325:LYS:HD2	2:D:746:HIS:CD2	2.35	0.61
1:C:363:ASP:HA	1:C:366:ILE:HG22	1.83	0.61
1:A:363:ASP:OD1	1:A:459:ARG:NH2	2.38	0.56
1:C:389:LEU:HD11	1:C:441:LEU:HD21	1.87	0.56
1:A:462:ASP:HB2	1:A:464:LYS:HZ1	1.69	0.55
1:C:437:LEU:O	1:C:440:ARG:HB2	2.08	0.54
1:C:376:ILE:HD12	1:C:440:ARG:HG3	1.90	0.53
1:C:388:LEU:HD12	1:C:422:LEU:HD12	1.90	0.53
1:C:270:PRO:HB3	1:C:339:ILE:HG12	1.92	0.51
1:C:302:LEU:HD11	1:C:394:ILE:HD11	1.93	0.51
1:A:401:TYR:CD1	1:A:401:TYR:N	2.80	0.50
1:A:401:TYR:N	1:A:401:TYR:HD1	2.10	0.49
1:A:430:ASN:C	1:A:432:GLN:H	2.19	0.49
1:C:357:ARG:HA	1:C:366:ILE:HG12	1.95	0.48
1:C:451:HIS:HA	1:C:454:MET:HE3	1.94	0.48
1:A:462:ASP:HB2	1:A:464:LYS:HZ3	1.80	0.46
1:C:325:LYS:HD2	2:D:746:HIS:NE2	2.30	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:465:PHE:H	1:A:476:GLN:HE22	1.64	0.46
1:C:369:MET:HE1	3:C:2:OMK:C19	2.46	0.46
1:C:397:PRO:HB2	1:C:405:ARG:HG2	1.97	0.46
1:A:363:ASP:HA	1:A:366:ILE:HD12	1.98	0.45
1:C:323:LEU:HG	1:C:402:ILE:HD11	1.99	0.45
1:C:400:GLN:O	1:C:401:TYR:HB2	2.15	0.45
1:A:458:TRP:HD1	1:A:460:VAL:HG13	1.81	0.45
1:C:445:ARG:NH1	4:C:66:HOH:O	2.47	0.44
1:A:273:ILE:HD13	1:A:294:MET:CE	2.44	0.44
1:A:330:GLU:HB3	1:A:444:LEU:HD13	1.99	0.44
1:C:379:LEU:HD13	1:C:437:LEU:HD23	2.00	0.44
1:A:409:GLU:O	1:A:413:GLU:HG3	2.18	0.43
1:A:469:LEU:O	1:A:473:TRP:HB2	2.18	0.43
1:C:261:MET:O	1:C:265:ASN:ND2	2.52	0.43
1:A:273:ILE:CD1	1:A:294:MET:HE1	2.45	0.42
1:A:472:ILE:HG23	1:A:473:TRP:CD1	2.54	0.42
2:D:749:LEU:HD23	2:D:749:LEU:HA	1.89	0.42
1:A:397:PRO:O	1:A:405:ARG:HG2	2.21	0.41
1:A:469:LEU:HA	1:A:472:ILE:HG22	2.03	0.41
1:A:440:ARG:HD3	1:A:440:ARG:HA	1.83	0.40
1:C:285:GLU:HG3	1:C:286:GLU:N	2.36	0.40
1:C:423:CYS:SG	1:C:433:HIS:HD2	2.44	0.40
1:A:465:PHE:H	1:A:476:GLN:NE2	2.19	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	230/233 (99%)	221 (96%)	9 (4%)	0	100	100
1	C	229/233 (98%)	221 (96%)	8 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	10/14 (71%)	8 (80%)	2 (20%)	0	100	100
2	D	9/14 (64%)	8 (89%)	1 (11%)	0	100	100
All	All	478/494 (97%)	458 (96%)	20 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	213/213 (100%)	202 (95%)	11 (5%)	21	13
1	C	212/213 (100%)	199 (94%)	13 (6%)	17	9
2	B	11/14 (79%)	10 (91%)	1 (9%)	9	3
2	D	10/14 (71%)	10 (100%)	0	100	100
All	All	446/454 (98%)	421 (94%)	25 (6%)	19	11

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	263	SER
1	A	279	LYS
1	A	313	GLN
1	A	321	ILE
1	A	363	ASP
1	A	381	MET
1	A	401	TYR
1	A	422	LEU
1	A	429	GLU
1	A	432	GLN
1	A	444	LEU
2	B	747	GLN
1	C	247	MET
1	C	263	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	273	ILE
1	C	276	LYS
1	C	313	GLN
1	C	318	GLU
1	C	398	ASP
1	C	403	LYS
1	C	440	ARG
1	C	446	THR
1	C	459	ARG
1	C	472	ILE
1	C	475	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	348	HIS
1	A	426	HIS
1	A	450	HIS
1	A	461	ASN
1	C	254	GLN
1	C	298	HIS
1	C	348	HIS
1	C	432	GLN
1	C	433	HIS
1	C	448	ASN
1	C	449	HIS
1	C	450	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	OMK	C	2	-	38,39,39	1.15	1 (2%)	53,56,56	0.99	3 (5%)
3	OMK	A	1	-	38,39,39	1.10	3 (7%)	53,56,56	1.07	3 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OMK	C	2	-	-	8/20/28/28	0/5/5/5
3	OMK	A	1	-	-	8/20/28/28	0/5/5/5

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	2	OMK	C4-C5	3.35	1.45	1.40
3	A	1	OMK	C7-C6	2.53	1.57	1.53
3	A	1	OMK	C2-N1	2.16	1.42	1.38
3	A	1	OMK	C4-C5	2.15	1.43	1.40

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1	OMK	C12-C2-N3	3.07	126.70	122.89
3	C	2	OMK	C29-C17-C7	-3.04	107.99	111.55
3	C	2	OMK	C12-C2-N3	2.75	126.31	122.89
3	A	1	OMK	C5-C4-N1	2.68	107.36	105.12
3	A	1	OMK	F21-C13-C10	2.19	123.02	118.64
3	C	2	OMK	C8-C4-N1	2.19	132.43	128.59

There are no chirality outliers.

All (16) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1	OMK	C19-C12-C2-N3
3	C	2	OMK	C19-C12-C2-N1
3	A	1	OMK	C18-C12-C2-N3
3	A	1	OMK	C19-C12-C2-N1
3	C	2	OMK	C19-C12-C2-N3
3	C	2	OMK	C26-C14-N9-C6
3	C	2	OMK	C18-C12-C2-N3
3	C	2	OMK	C18-C12-C2-N1
3	A	1	OMK	O15-C6-C7-N1
3	A	1	OMK	N9-C6-C7-N1
3	C	2	OMK	N9-C6-C7-N1
3	A	1	OMK	C18-C12-C2-N1
3	A	1	OMK	C26-C14-N9-C6
3	C	2	OMK	C16-C14-N9-C6
3	C	2	OMK	O15-C6-C7-N1
3	A	1	OMK	C16-C14-N9-C6


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	2	OMK	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	230/233 (98%)	0.26	6 (2%) 57 61	18, 39, 65, 83	2 (0%)
1	C	230/233 (98%)	0.36	11 (4%) 35 38	22, 42, 65, 84	1 (0%)
2	B	12/14 (85%)	0.45	1 (8%) 17 18	26, 50, 68, 69	0
2	D	11/14 (78%)	0.98	1 (9%) 15 15	35, 41, 70, 71	0
All	All	483/494 (97%)	0.33	19 (3%) 43 46	18, 41, 67, 84	3 (0%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	475	VAL	4.1
1	C	366	ILE	3.6
1	A	426	HIS	3.5
2	D	756	ASP	3.2
1	A	476	GLN	3.2
1	A	425	ILE	3.1
1	C	271	GLN	2.9
1	C	273	ILE	2.8
1	A	462	ASP	2.6
1	C	247	MET	2.5
1	C	269	MET	2.5
1	C	460	VAL	2.3
1	C	425	ILE	2.2
1	C	317[A]	HIS	2.2
1	A	429	GLU	2.2
2	B	755	LYS	2.1
1	A	442	THR	2.0
1	C	430	ASN	2.0
1	C	476	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	OMK	C	2	35/35	0.96	0.06	21,30,38,43	0
3	OMK	A	1	35/35	0.97	0.06	21,25,32,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around OMK C 2:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.