



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 02:23 AM UTC

PDB ID : 4OMB / pdb\_00004omb  
Title : Phosphate binding protein  
Authors : Neznansky, A.; Opatowsky, Y.  
Deposited on : 2014-01-27  
Resolution : 1.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	347	 4% 82% 14%
1	B	347	 7% 82% 14%
1	C	347	 7% 80% 5% 14%
1	D	347	 12% 80% 6% 14%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9495 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphate binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	298	2256	1437	378	434	7	0	0	0
1	B	298	2256	1437	378	434	7	0	0	0
1	C	298	2256	1437	378	434	7	0	0	0
1	D	298	2256	1437	378	434	7	0	0	0

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	expression tag	UNP Q9Z9M3
A	-9	ASP	-	expression tag	UNP Q9Z9M3
A	-8	ILE	-	expression tag	UNP Q9Z9M3
A	-7	GLY	-	expression tag	UNP Q9Z9M3
A	-6	ILE	-	expression tag	UNP Q9Z9M3
A	-5	ASN	-	expression tag	UNP Q9Z9M3
A	-4	SER	-	expression tag	UNP Q9Z9M3
A	-3	ASP	-	expression tag	UNP Q9Z9M3
A	-2	PRO	-	expression tag	UNP Q9Z9M3
A	-1	ASN	-	expression tag	UNP Q9Z9M3
A	0	SER	-	expression tag	UNP Q9Z9M3
A	324	LYS	-	expression tag	UNP Q9Z9M3
A	325	LEU	-	expression tag	UNP Q9Z9M3
A	326	ALA	-	expression tag	UNP Q9Z9M3
A	327	ALA	-	expression tag	UNP Q9Z9M3
A	328	ALA	-	expression tag	UNP Q9Z9M3
A	329	LEU	-	expression tag	UNP Q9Z9M3
A	330	GLU	-	expression tag	UNP Q9Z9M3
A	331	HIS	-	expression tag	UNP Q9Z9M3
A	332	HIS	-	expression tag	UNP Q9Z9M3
A	333	HIS	-	expression tag	UNP Q9Z9M3

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	334	HIS	-	expression tag	UNP Q9Z9M3
A	335	HIS	-	expression tag	UNP Q9Z9M3
A	336	HIS	-	expression tag	UNP Q9Z9M3
B	-10	MET	-	expression tag	UNP Q9Z9M3
B	-9	ASP	-	expression tag	UNP Q9Z9M3
B	-8	ILE	-	expression tag	UNP Q9Z9M3
B	-7	GLY	-	expression tag	UNP Q9Z9M3
B	-6	ILE	-	expression tag	UNP Q9Z9M3
B	-5	ASN	-	expression tag	UNP Q9Z9M3
B	-4	SER	-	expression tag	UNP Q9Z9M3
B	-3	ASP	-	expression tag	UNP Q9Z9M3
B	-2	PRO	-	expression tag	UNP Q9Z9M3
B	-1	ASN	-	expression tag	UNP Q9Z9M3
B	0	SER	-	expression tag	UNP Q9Z9M3
B	324	LYS	-	expression tag	UNP Q9Z9M3
B	325	LEU	-	expression tag	UNP Q9Z9M3
B	326	ALA	-	expression tag	UNP Q9Z9M3
B	327	ALA	-	expression tag	UNP Q9Z9M3
B	328	ALA	-	expression tag	UNP Q9Z9M3
B	329	LEU	-	expression tag	UNP Q9Z9M3
B	330	GLU	-	expression tag	UNP Q9Z9M3
B	331	HIS	-	expression tag	UNP Q9Z9M3
B	332	HIS	-	expression tag	UNP Q9Z9M3
B	333	HIS	-	expression tag	UNP Q9Z9M3
B	334	HIS	-	expression tag	UNP Q9Z9M3
B	335	HIS	-	expression tag	UNP Q9Z9M3
B	336	HIS	-	expression tag	UNP Q9Z9M3
C	-10	MET	-	expression tag	UNP Q9Z9M3
C	-9	ASP	-	expression tag	UNP Q9Z9M3
C	-8	ILE	-	expression tag	UNP Q9Z9M3
C	-7	GLY	-	expression tag	UNP Q9Z9M3
C	-6	ILE	-	expression tag	UNP Q9Z9M3
C	-5	ASN	-	expression tag	UNP Q9Z9M3
C	-4	SER	-	expression tag	UNP Q9Z9M3
C	-3	ASP	-	expression tag	UNP Q9Z9M3
C	-2	PRO	-	expression tag	UNP Q9Z9M3
C	-1	ASN	-	expression tag	UNP Q9Z9M3
C	0	SER	-	expression tag	UNP Q9Z9M3
C	324	LYS	-	expression tag	UNP Q9Z9M3
C	325	LEU	-	expression tag	UNP Q9Z9M3
C	326	ALA	-	expression tag	UNP Q9Z9M3
C	327	ALA	-	expression tag	UNP Q9Z9M3

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	328	ALA	-	expression tag	UNP Q9Z9M3
C	329	LEU	-	expression tag	UNP Q9Z9M3
C	330	GLU	-	expression tag	UNP Q9Z9M3
C	331	HIS	-	expression tag	UNP Q9Z9M3
C	332	HIS	-	expression tag	UNP Q9Z9M3
C	333	HIS	-	expression tag	UNP Q9Z9M3
C	334	HIS	-	expression tag	UNP Q9Z9M3
C	335	HIS	-	expression tag	UNP Q9Z9M3
C	336	HIS	-	expression tag	UNP Q9Z9M3
D	-10	MET	-	expression tag	UNP Q9Z9M3
D	-9	ASP	-	expression tag	UNP Q9Z9M3
D	-8	ILE	-	expression tag	UNP Q9Z9M3
D	-7	GLY	-	expression tag	UNP Q9Z9M3
D	-6	ILE	-	expression tag	UNP Q9Z9M3
D	-5	ASN	-	expression tag	UNP Q9Z9M3
D	-4	SER	-	expression tag	UNP Q9Z9M3
D	-3	ASP	-	expression tag	UNP Q9Z9M3
D	-2	PRO	-	expression tag	UNP Q9Z9M3
D	-1	ASN	-	expression tag	UNP Q9Z9M3
D	0	SER	-	expression tag	UNP Q9Z9M3
D	324	LYS	-	expression tag	UNP Q9Z9M3
D	325	LEU	-	expression tag	UNP Q9Z9M3
D	326	ALA	-	expression tag	UNP Q9Z9M3
D	327	ALA	-	expression tag	UNP Q9Z9M3
D	328	ALA	-	expression tag	UNP Q9Z9M3
D	329	LEU	-	expression tag	UNP Q9Z9M3
D	330	GLU	-	expression tag	UNP Q9Z9M3
D	331	HIS	-	expression tag	UNP Q9Z9M3
D	332	HIS	-	expression tag	UNP Q9Z9M3
D	333	HIS	-	expression tag	UNP Q9Z9M3
D	334	HIS	-	expression tag	UNP Q9Z9M3
D	335	HIS	-	expression tag	UNP Q9Z9M3
D	336	HIS	-	expression tag	UNP Q9Z9M3

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P 5 4 1	0	0
2	B	1	Total O P 5 4 1	0	0
2	C	1	Total O P 5 4 1	0	0
2	D	1	Total O P 5 4 1	0	0

- Molecule 3 is PLATINUM (II) ION (CCD ID: PT) (formula: Pt).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Pt 1 1	0	0
3	B	1	Total Pt 1 1	0	0
3	C	1	Total Pt 1 1	0	0
3	D	1	Total Pt 1 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	162	Total O 162 162	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	B	122	Total 122	O 122	0	0
4	C	97	Total 97	O 97	0	0
4	D	66	Total 66	O 66	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Phosphate binding protein




- Molecule 1: Phosphate binding protein




- Molecule 1: Phosphate binding protein



- Molecule 1: Phosphate binding protein





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	35.44Å 148.09Å 216.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	122.20 – 1.50 122.20 – 1.50	Depositor EDS
% Data completeness (in resolution range)	98.3 (122.20-1.50) 99.1 (122.20-1.50)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	0.15	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.96 (at 1.50Å)	Xtrriage
Refinement program	REFMAC 5.8.0049	Depositor
R, $R_{free}$	0.190 , 0.216 0.191 , 0.218	Depositor DCC
$R_{free}$ test set	9127 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.1	Xtrriage
Anisotropy	0.605	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 41.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.42$ , $\langle L^2 \rangle = 0.25$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	9495	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.23% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PT, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.99	1/2302 (0.0%)	1.00	2/3123 (0.1%)
1	B	0.92	2/2302 (0.1%)	0.93	0/3123
1	C	0.97	1/2302 (0.0%)	0.92	2/3123 (0.1%)
1	D	0.83	1/2302 (0.0%)	0.89	7/3123 (0.2%)
All	All	0.93	5/9208 (0.1%)	0.94	11/12492 (0.1%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	106	ALA	N-CA	5.99	1.53	1.46
1	C	183	SER	CB-OG	5.99	1.54	1.42
1	A	273	VAL	N-CA	-5.18	1.40	1.46
1	D	183	SER	CB-OG	5.04	1.52	1.42
1	B	183	SER	CB-OG	5.04	1.52	1.42

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	277	PRO	CA-C-N	-7.13	111.20	123.25
1	A	277	PRO	C-N-CA	-7.13	111.20	123.25
1	D	37	SER	N-CA-C	7.08	122.23	112.04
1	D	109	GLN	CB-CA-C	-5.36	101.89	110.79
1	C	182	ASN	CA-C-N	5.31	127.92	120.28

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2256	0	2273	7	0
1	B	2256	0	2273	10	0
1	C	2256	0	2273	12	1
1	D	2256	0	2273	8	1
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	162	0	0	0	1
4	B	122	0	0	0	0
4	C	97	0	0	0	0
4	D	66	0	0	1	0
All	All	9495	0	9092	34	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 34 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:245:LYS:HD2	1:B:246:GLU:H	1.39	0.87
1:C:135:ILE:HD12	1:C:167:LEU:HD11	1.78	0.66
1:C:51:LEU:HD23	1:C:54:LEU:HD12	1.78	0.65
1:D:51:LEU:HD23	1:D:54:LEU:HD12	1.79	0.64
1:B:51:LEU:HD23	1:B:54:LEU:HD12	1.82	0.60

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:306:TYR:O	1:D:312:LYS:NZ[1_655]	2.06	0.14
4:A:621:HOH:O	4:A:633:HOH:O[1_655]	2.14	0.06

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	296/347 (85%)	292 (99%)	4 (1%)	0	100	100
1	B	296/347 (85%)	291 (98%)	5 (2%)	0	100	100
1	C	296/347 (85%)	292 (99%)	4 (1%)	0	100	100
1	D	296/347 (85%)	292 (99%)	4 (1%)	0	100	100
All	All	1184/1388 (85%)	1167 (99%)	17 (1%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/276 (87%)	238 (99%)	3 (1%)	63	38
1	B	241/276 (87%)	238 (99%)	3 (1%)	63	38
1	C	241/276 (87%)	238 (99%)	3 (1%)	63	38
1	D	241/276 (87%)	237 (98%)	4 (2%)	53	26
All	All	964/1104 (87%)	951 (99%)	13 (1%)	61	35

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	37	SER
1	C	71	ILE
1	D	319	LYS
1	D	135	ILE
1	D	245	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	287	GLN
1	D	216	GLN
1	D	42	ASN
1	C	287	GLN
1	D	70	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PO4	A	401	-	4,4,4	1.79	1 (25%)	6,6,6	0.87	0
2	PO4	B	401	-	4,4,4	1.68	2 (50%)	6,6,6	0.92	0
2	PO4	C	401	-	4,4,4	3.24	2 (50%)	6,6,6	1.34	1 (16%)
2	PO4	D	401	-	4,4,4	2.68	1 (25%)	6,6,6	1.21	1 (16%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	401	PO4	P-O1	-5.09	1.38	1.50
2	D	401	PO4	P-O1	-4.91	1.39	1.50
2	C	401	PO4	P-O3	-3.88	1.43	1.54
2	A	401	PO4	P-O2	-3.53	1.44	1.54
2	B	401	PO4	P-O1	-2.59	1.44	1.50

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	401	PO4	O3-P-O1	2.81	120.87	110.95
2	D	401	PO4	O2-P-O1	2.10	118.36	110.95

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	298/347 (85%)	0.17	14 (4%) 36 40	13, 21, 50, 81	0
1	B	298/347 (85%)	0.42	25 (8%) 17 18	15, 26, 67, 87	0
1	C	298/347 (85%)	0.45	23 (7%) 19 21	15, 27, 60, 75	0
1	D	298/347 (85%)	0.80	40 (13%) 7 7	17, 34, 74, 91	0
All	All	1192/1388 (85%)	0.46	102 (8%) 16 17	13, 27, 63, 91	0

The worst 5 of 102 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	322	GLY	5.3
1	C	71	ILE	4.5
1	D	25	ALA	4.5
1	A	322	GLY	4.1
1	A	25	ALA	4.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	PT	B	402	1/1	0.89	0.10	118,118,118,118	0
3	PT	D	402	1/1	0.91	0.09	121,121,121,121	0
3	PT	C	402	1/1	0.92	0.09	110,110,110,110	0
3	PT	A	402	1/1	0.93	0.15	84,84,84,84	0
2	PO4	C	401	5/5	0.98	0.05	15,17,18,19	0
2	PO4	D	401	5/5	0.98	0.05	18,19,21,23	0
2	PO4	A	401	5/5	0.99	0.04	15,15,15,16	0
2	PO4	B	401	5/5	0.99	0.05	17,17,17,18	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.