



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 11:38 AM UTC

PDB ID : 6OMN / pdb_00006omn
Title : Glycosylated BMP2 homodimer
Authors : Juo, Z.S.; Seeherman, H.
Deposited on : 2019-04-19
Resolution : 2.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5070 (2.70-2.66)
Clashscore	190562	5409 (2.70-2.66)
Ramachandran outliers	187476	5324 (2.70-2.66)
Sidechain outliers	187428	5324 (2.70-2.66)
RSRZ outliers	180081	5070 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	E	108	2% 72% 18% 6% 5%
1	F	108	5% 80% 17% ...
1	G	108	5% 77% 16% 7%
1	H	108	2% 72% 21% ...
2	A	5	100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	B	5	 100%
2	C	5	 100%
2	D	5	 100%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3595 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bone morphogenetic protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	E	103	Total	C	N	O	S	0	0	0
			805	510	135	151	9			
1	F	106	Total	C	N	O	S	0	0	0
			833	528	142	154	9			
1	G	108	Total	C	N	O	S	0	0	0
			853	540	148	156	9			
1	H	104	Total	C	N	O	S	0	0	0
			814	516	137	152	9			

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	A	5	Total	C	N	O	0	0	0
			61	34	2	25			
2	B	5	Total	C	N	O	0	0	0
			61	34	2	25			
2	C	5	Total	C	N	O	0	0	0
			61	34	2	25			
2	D	5	Total	C	N	O	0	0	0
			61	34	2	25			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	9	Total	O	0	0
			9	9		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	F	13	Total 13	O 13	0	0
3	G	11	Total 11	O 11	0	0
3	H	13	Total 13	O 13	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Bone morphogenetic protein 2




- Molecule 1: Bone morphogenetic protein 2



- Molecule 1: Bone morphogenetic protein 2



- Molecule 1: Bone morphogenetic protein 2



- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B:  100%

MAG1
MAG2
BMA3
MAN4
MAN5

- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:  100%

MAG1
MAG2
BMA3
MAN4
MAN5

- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  100%

MAG1
MAG2
BMA3
MAN4
MAN5

4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	62.74Å 62.74Å 126.35Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	41.20 – 2.68 41.20 – 2.68	Depositor EDS
% Data completeness (in resolution range)	98.9 (41.20-2.68) 99.1 (41.20-2.68)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 2.69Å)	Xtriage
Refinement program	BUSTER 2.11.1	Depositor
R, R_{free}	0.175 , 0.243 0.194 , 0.250	Depositor DCC
R_{free} test set	776 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	61.2	Xtriage
Anisotropy	0.731	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 47.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.023 for -h,-k,l 0.063 for h,-h-k,-l 0.045 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3595	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MAN, NAG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	E	0.87	0/828	1.53	15/1129 (1.3%)
1	F	0.89	0/856	1.39	6/1165 (0.5%)
1	G	0.86	1/876 (0.1%)	1.44	10/1190 (0.8%)
1	H	0.85	0/837	1.38	6/1140 (0.5%)
All	All	0.87	1/3397 (0.0%)	1.43	37/4624 (0.8%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	89	MET	SD-CE	-5.40	1.66	1.79

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	40	ALA	CA-C-N	-9.58	105.31	123.32
1	E	40	ALA	C-N-CA	-9.58	105.31	123.32
1	E	18	PRO	CB-CA-C	-9.22	98.94	111.21
1	G	40	ALA	CA-C-N	-9.15	106.72	123.01
1	G	40	ALA	C-N-CA	-9.15	106.72	123.01
1	G	41	PHE	N-CA-C	-8.47	97.15	109.18
1	E	28	TRP	CA-C-N	8.23	131.65	120.38
1	E	28	TRP	C-N-CA	8.23	131.65	120.38
1	F	30	ASP	CB-CA-C	7.76	125.58	110.67
1	H	70	VAL	CB-CA-C	7.56	117.71	111.06
1	E	14	CYS	CB-CA-C	6.99	118.39	109.80
1	F	80	VAL	N-CA-C	6.98	116.82	108.45
1	H	56	ASN	N-CA-C	6.87	120.97	111.90
1	F	53	ASP	N-CA-C	-6.20	103.84	111.40
1	G	56	ASN	N-CA-C	6.18	120.06	111.90
1	G	71	ASN	CA-CB-CG	6.14	118.74	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	98	VAL	CB-CA-C	6.11	117.82	111.23
1	H	19	LEU	CB-CA-C	6.09	118.87	110.34
1	F	51	LEU	N-CA-CB	6.07	119.23	110.06
1	G	57	SER	N-CA-C	5.96	118.22	110.53
1	H	23	PHE	N-CA-C	5.93	117.82	111.36
1	H	40	ALA	N-CA-C	-5.93	99.07	108.73
1	E	65	THR	CA-C-N	5.76	128.47	120.29
1	E	65	THR	C-N-CA	5.76	128.47	120.29
1	G	105	ASP	CA-CB-CG	5.68	118.28	112.60
1	G	19	LEU	CB-CA-C	5.68	118.54	110.24
1	E	40	ALA	N-CA-C	-5.54	100.65	109.96
1	F	40	ALA	N-CA-C	-5.26	98.68	107.99
1	H	87	ILE	N-CA-CB	5.21	119.98	111.44
1	G	58	THR	CB-CA-C	5.16	119.00	109.71
1	G	23	PHE	N-CA-C	5.16	117.64	111.71
1	E	99	VAL	N-CA-C	5.15	116.38	108.71
1	E	93	ASP	CA-CB-CG	5.11	117.71	112.60
1	E	22	ASP	CA-CB-CG	5.09	117.69	112.60
1	E	29	ASN	CA-CB-CG	5.05	117.66	112.60
1	F	53	ASP	CA-CB-CG	5.02	117.62	112.60
1	E	41	PHE	N-CA-C	-5.02	102.47	109.69

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	805	0	760	23	0
1	F	833	0	797	16	0
1	G	853	0	823	24	0
1	H	814	0	773	26	0
2	A	61	0	52	0	0
2	B	61	0	52	0	0
2	C	61	0	52	0	0
2	D	61	0	52	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	9	0	0	0	0
3	F	13	0	0	0	0
3	G	11	0	0	0	0
3	H	13	0	0	0	0
All	All	3595	0	3361	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:87:ILE:HD11	1:E:89:MET:HE3	1.44	1.00
1:G:26:VAL:CG1	1:H:66:LEU:CD2	2.60	0.79
1:E:87:ILE:CD1	1:E:89:MET:HE3	2.14	0.77
1:G:26:VAL:CG1	1:H:66:LEU:HD23	2.16	0.76
1:E:92:LEU:HD13	1:E:92:LEU:C	2.11	0.76
1:F:102:ASN:HD21	1:H:102:ASN:HD21	1.35	0.72
1:E:49:PHE:HE1	1:E:66:LEU:HD12	1.55	0.71
1:F:102:ASN:HD21	1:H:102:ASN:ND2	1.89	0.70
1:G:104:GLN:O	1:G:105:ASP:OD1	2.10	0.69
1:E:92:LEU:HD21	1:E:96:GLU:HA	1.77	0.67
1:G:105:ASP:OD2	1:H:58:THR:HG22	1.94	0.67
1:G:26:VAL:HG13	1:H:66:LEU:CD2	2.23	0.67
1:H:84:LEU:HD13	1:H:105:ASP:HB3	1.78	0.66
1:E:92:LEU:HD13	1:E:93:ASP:O	1.96	0.65
1:F:38:TYR:CZ	1:F:40:ALA:HB2	2.32	0.65
1:F:51:LEU:HD11	1:F:62:ILE:HA	1.79	0.62
1:H:38:TYR:CZ	1:H:40:ALA:HB2	2.35	0.61
1:G:26:VAL:HG12	1:H:66:LEU:HD21	1.82	0.60
1:G:26:VAL:CG1	1:H:66:LEU:HD21	2.33	0.58
1:G:26:VAL:HG12	1:H:66:LEU:CD2	2.34	0.57
1:E:92:LEU:C	1:E:92:LEU:CD1	2.77	0.56
1:E:19:LEU:O	1:E:40:ALA:O	2.23	0.56
1:E:22:ASP:O	1:E:26:VAL:HG23	2.07	0.55
1:E:41:PHE:O	1:F:60:HIS:NE2	2.40	0.55
1:H:52:ALA:HB3	1:H:55:LEU:HD23	1.89	0.55
1:H:87:ILE:HD11	1:H:89:MET:HE3	1.89	0.54
1:F:51:LEU:CD1	1:F:62:ILE:HA	2.38	0.53
1:G:19:LEU:O	1:G:40:ALA:O	2.26	0.53
1:E:23:PHE:CZ	1:E:89:MET:HE2	2.45	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:26:VAL:HG13	1:F:66:LEU:HG	1.93	0.51
1:E:26:VAL:HG12	1:F:66:LEU:HD21	1.93	0.50
1:F:38:TYR:CE1	1:F:40:ALA:HB2	2.47	0.49
1:E:26:VAL:CG1	1:F:66:LEU:HD21	2.42	0.49
1:E:28:TRP:CZ2	1:F:66:LEU:HD22	2.48	0.49
1:G:105:ASP:OD2	1:H:58:THR:CB	2.61	0.49
1:E:26:VAL:CG1	1:F:66:LEU:CD2	2.90	0.49
1:G:34:ALA:HB3	1:G:90:LEU:HB2	1.95	0.48
1:G:105:ASP:OD2	1:H:58:THR:CG2	2.59	0.48
1:G:41:PHE:O	1:H:60:HIS:NE2	2.47	0.48
1:E:92:LEU:CD1	1:E:93:ASP:O	2.63	0.47
1:G:26:VAL:HG13	1:H:66:LEU:HG	1.97	0.47
1:G:105:ASP:OD2	1:H:58:THR:HB	2.14	0.47
1:H:55:LEU:O	1:H:112:GLY:HA2	2.15	0.47
1:E:23:PHE:HZ	1:E:89:MET:HE2	1.80	0.46
1:F:84:LEU:HD13	1:F:105:ASP:HB3	1.99	0.45
1:F:68:ASN:HA	1:F:71:ASN:O	2.16	0.45
1:F:24:SER:HA	1:F:29:ASN:OD1	2.17	0.44
1:G:22:ASP:O	1:G:26:VAL:HG23	2.18	0.43
1:H:84:LEU:HA	1:H:107:VAL:O	2.18	0.43
1:G:89:MET:HB2	1:G:103:TYR:CD2	2.54	0.43
1:H:49:PHE:HB2	1:H:65:THR:HG22	2.01	0.43
1:E:92:LEU:HD21	1:E:96:GLU:CA	2.46	0.43
1:H:41:PHE:HB2	1:H:108:VAL:HB	2.01	0.43
1:E:23:PHE:HZ	1:E:89:MET:CE	2.32	0.42
1:E:43:CYS:HB3	1:E:79:CYS:SG	2.60	0.42
1:E:92:LEU:HD13	1:E:93:ASP:N	2.34	0.42
1:G:26:VAL:HG13	1:H:66:LEU:CG	2.49	0.42
1:G:38:TYR:CZ	1:G:40:ALA:HB2	2.54	0.42
1:G:60:HIS:NE2	1:H:41:PHE:O	2.52	0.41
1:H:35:PRO:HD2	1:H:89:MET:HB3	2.01	0.41
1:H:87:ILE:CD1	1:H:89:MET:HE3	2.51	0.41
1:E:106:MET:SD	1:F:59:ASN:HB3	2.60	0.41
1:G:38:TYR:CD1	1:G:87:ILE:HD13	2.56	0.41
1:G:89:MET:HB2	1:G:103:TYR:HD2	1.86	0.40
1:G:55:LEU:O	1:G:112:GLY:HA2	2.22	0.40
1:G:58:THR:HG23	1:G:61:ALA:H	1.84	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	101/108 (94%)	100 (99%)	1 (1%)	0	100	100
1	F	104/108 (96%)	102 (98%)	2 (2%)	0	100	100
1	G	106/108 (98%)	104 (98%)	2 (2%)	0	100	100
1	H	102/108 (94%)	101 (99%)	1 (1%)	0	100	100
All	All	413/432 (96%)	407 (98%)	6 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	92/97 (95%)	89 (97%)	3 (3%)	33	60
1	F	95/97 (98%)	91 (96%)	4 (4%)	26	52
1	G	97/97 (100%)	92 (95%)	5 (5%)	21	43
1	H	93/97 (96%)	89 (96%)	4 (4%)	26	51
All	All	377/388 (97%)	361 (96%)	16 (4%)	26	52

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	74	ILE
1	E	78	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	92	LEU
1	F	51	LEU
1	F	70	VAL
1	F	92	LEU
1	F	104	GLN
1	G	55	LEU
1	G	70	VAL
1	G	74	ILE
1	G	78	CYS
1	G	104	GLN
1	H	57	SER
1	H	70	VAL
1	H	92	LEU
1	H	96	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	102	ASN
1	F	64	GLN
1	G	104	GLN
1	H	17	HIS
1	H	68	ASN
1	H	71	ASN
1	H	102	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

20 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	A	1	2,1	14,14,15	1.54	2 (14%)	17,19,21	1.87	5 (29%)
2	NAG	A	2	2	14,14,15	1.60	2 (14%)	17,19,21	1.92	6 (35%)
2	BMA	A	3	2	11,11,12	1.30	1 (9%)	15,15,17	1.78	2 (13%)
2	MAN	A	4	2	11,11,12	1.40	3 (27%)	15,15,17	1.06	1 (6%)
2	MAN	A	5	2	11,11,12	1.60	2 (18%)	15,15,17	1.27	1 (6%)
2	NAG	B	1	2,1	14,14,15	1.22	1 (7%)	17,19,21	1.57	5 (29%)
2	NAG	B	2	2	14,14,15	1.47	2 (14%)	17,19,21	2.17	6 (35%)
2	BMA	B	3	2	11,11,12	1.37	2 (18%)	15,15,17	1.46	2 (13%)
2	MAN	B	4	2	11,11,12	1.32	1 (9%)	15,15,17	1.45	3 (20%)
2	MAN	B	5	2	11,11,12	1.73	3 (27%)	15,15,17	1.84	3 (20%)
2	NAG	C	1	2,1	14,14,15	0.88	0	17,19,21	1.54	5 (29%)
2	NAG	C	2	2	14,14,15	1.36	2 (14%)	17,19,21	1.30	2 (11%)
2	BMA	C	3	2	11,11,12	1.00	1 (9%)	15,15,17	1.29	3 (20%)
2	MAN	C	4	2	11,11,12	1.27	0	15,15,17	1.58	2 (13%)
2	MAN	C	5	2	11,11,12	1.60	4 (36%)	15,15,17	1.66	3 (20%)
2	NAG	D	1	2,1	14,14,15	1.39	3 (21%)	17,19,21	1.52	2 (11%)
2	NAG	D	2	2	14,14,15	1.02	2 (14%)	17,19,21	2.10	5 (29%)
2	BMA	D	3	2	11,11,12	1.48	3 (27%)	15,15,17	1.85	3 (20%)
2	MAN	D	4	2	11,11,12	1.55	1 (9%)	15,15,17	1.68	5 (33%)
2	MAN	D	5	2	11,11,12	1.66	4 (36%)	15,15,17	1.56	5 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	A	2	2	-	0/6/23/26	0/1/1/1
2	BMA	A	3	2	-	0/2/19/22	0/1/1/1
2	MAN	A	4	2	-	0/2/19/22	0/1/1/1
2	MAN	A	5	2	-	2/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	1/2/19/22	0/1/1/1
2	MAN	B	4	2	-	0/2/19/22	0/1/1/1
2	MAN	B	5	2	-	2/2/19/22	0/1/1/1
2	NAG	C	1	2,1	-	1/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	BMA	C	3	2	-	1/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
2	MAN	C	5	2	-	1/2/19/22	0/1/1/1
2	NAG	D	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1
2	BMA	D	3	2	-	0/2/19/22	0/1/1/1
2	MAN	D	4	2	-	0/2/19/22	0/1/1/1
2	MAN	D	5	2	-	2/2/19/22	0/1/1/1

All (39) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2	NAG	C1-C2	4.28	1.58	1.52
2	A	1	NAG	C1-C2	3.64	1.57	1.52
2	C	2	NAG	C1-C2	3.42	1.57	1.52
2	B	2	NAG	C1-C2	3.40	1.57	1.52
2	B	5	MAN	O5-C5	3.26	1.49	1.43
2	A	3	BMA	C2-C3	3.08	1.57	1.52
2	D	4	MAN	C2-C3	2.98	1.57	1.52
2	D	3	BMA	C4-C5	2.87	1.59	1.53
2	B	4	MAN	C2-C3	2.71	1.56	1.52
2	D	1	NAG	C4-C5	2.56	1.58	1.53
2	A	2	NAG	C3-C2	2.56	1.57	1.52
2	B	3	BMA	C4-C3	2.52	1.58	1.52
2	D	5	MAN	O5-C5	2.50	1.48	1.43
2	D	3	BMA	O2-C2	2.40	1.48	1.43
2	D	5	MAN	O5-C1	2.38	1.47	1.43
2	A	5	MAN	C4-C5	2.37	1.58	1.53
2	A	4	MAN	C2-C3	2.36	1.56	1.52
2	B	1	NAG	C4-C5	2.36	1.58	1.53
2	D	1	NAG	O7-C7	2.32	1.28	1.23
2	D	2	NAG	C4-C5	2.31	1.57	1.53
2	D	3	BMA	O5-C5	2.22	1.47	1.43
2	A	4	MAN	C4-C3	2.19	1.58	1.52
2	D	5	MAN	C1-C2	2.17	1.57	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	NAG	C3-C2	2.15	1.57	1.52
2	B	5	MAN	C4-C5	2.14	1.57	1.53
2	C	3	BMA	O5-C5	2.14	1.47	1.43
2	B	3	BMA	C2-C3	2.13	1.55	1.52
2	A	1	NAG	C7-N2	2.13	1.41	1.34
2	C	5	MAN	C1-C2	2.11	1.57	1.52
2	D	2	NAG	C1-C2	2.08	1.55	1.52
2	D	5	MAN	C4-C5	2.07	1.57	1.53
2	C	5	MAN	C4-C3	2.06	1.57	1.52
2	B	5	MAN	C1-C2	2.06	1.57	1.52
2	C	2	NAG	O5-C5	2.05	1.47	1.43
2	A	4	MAN	O2-C2	2.03	1.47	1.43
2	D	1	NAG	C3-C2	2.03	1.56	1.52
2	A	5	MAN	C2-C3	2.01	1.55	1.52
2	C	5	MAN	C4-C5	2.01	1.57	1.53
2	C	5	MAN	O5-C5	2.00	1.47	1.43

All (69) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	NAG	C1-C2-N2	6.26	120.30	110.43
2	B	2	NAG	C1-C2-N2	5.57	119.21	110.43
2	A	1	NAG	C1-C2-N2	4.38	117.34	110.43
2	A	3	BMA	C2-C3-C4	4.34	118.50	110.86
2	B	2	NAG	C3-C4-C5	4.02	117.53	110.23
2	D	3	BMA	C1-C2-C3	-3.95	103.89	109.64
2	A	2	NAG	C1-C2-N2	3.90	116.58	110.43
2	D	1	NAG	C8-C7-N2	-3.86	109.72	116.12
2	A	3	BMA	C1-O5-C5	-3.62	107.34	112.19
2	B	5	MAN	C1-C2-C3	-3.59	104.41	109.64
2	B	5	MAN	O3-C3-C4	3.52	118.67	110.38
2	C	5	MAN	C3-C4-C5	3.51	116.60	110.23
2	D	3	BMA	O4-C4-C3	-3.48	102.16	110.38
2	C	4	MAN	O4-C4-C5	3.46	117.86	109.32
2	A	1	NAG	C1-O5-C5	-3.34	107.71	112.19
2	D	4	MAN	C2-C3-C4	3.18	116.46	110.86
2	B	4	MAN	C2-C3-C4	3.17	116.43	110.86
2	A	2	NAG	O5-C5-C6	3.09	113.68	107.66
2	D	1	NAG	O3-C3-C2	3.08	115.80	109.40
2	B	1	NAG	O5-C1-C2	-3.08	106.53	111.29
2	D	2	NAG	O4-C4-C3	-3.07	103.14	110.38
2	B	2	NAG	C1-O5-C5	3.01	116.22	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2	NAG	O4-C4-C3	-2.95	103.43	110.38
2	D	5	MAN	O2-C2-C1	2.94	115.94	109.22
2	C	5	MAN	C2-C3-C4	2.93	116.01	110.86
2	B	2	NAG	O5-C1-C2	-2.92	106.77	111.29
2	C	3	BMA	O4-C4-C3	-2.88	103.59	110.38
2	C	4	MAN	O2-C2-C3	2.87	116.11	110.15
2	D	4	MAN	O5-C5-C4	2.81	117.67	110.83
2	A	2	NAG	C2-N2-C7	2.79	126.64	122.90
2	D	2	NAG	O3-C3-C4	-2.79	103.81	110.38
2	C	5	MAN	O5-C5-C4	2.76	117.53	110.83
2	B	4	MAN	C3-C4-C5	2.71	115.14	110.23
2	D	4	MAN	O5-C5-C6	2.68	112.88	107.66
2	B	1	NAG	C3-C4-C5	-2.57	105.57	110.23
2	B	1	NAG	C8-C7-N2	-2.57	111.86	116.12
2	B	3	BMA	C1-C2-C3	-2.55	105.92	109.64
2	A	5	MAN	C2-C3-C4	2.52	115.29	110.86
2	C	2	NAG	C4-C3-C2	2.49	114.67	111.02
2	C	1	NAG	O3-C3-C2	-2.49	104.22	109.40
2	D	5	MAN	O4-C4-C3	-2.49	104.50	110.38
2	B	5	MAN	O5-C5-C6	2.48	112.50	107.66
2	D	2	NAG	C8-C7-N2	-2.45	112.06	116.12
2	C	1	NAG	C8-C7-N2	-2.42	112.11	116.12
2	A	2	NAG	C1-O5-C5	-2.40	108.97	112.19
2	C	1	NAG	O7-C7-C8	2.39	126.30	122.05
2	C	1	NAG	O5-C1-C2	-2.37	107.62	111.29
2	B	4	MAN	C1-O5-C5	2.36	115.35	112.19
2	D	4	MAN	C1-C2-C3	2.35	113.07	109.64
2	C	3	BMA	O5-C5-C6	2.34	112.22	107.66
2	D	2	NAG	O5-C1-C2	-2.32	107.71	111.29
2	B	1	NAG	O3-C3-C2	2.29	114.16	109.40
2	A	1	NAG	C8-C7-N2	-2.28	112.33	116.12
2	A	1	NAG	O4-C4-C5	-2.27	103.73	109.32
2	B	3	BMA	O3-C3-C2	-2.24	105.49	110.05
2	A	1	NAG	C2-N2-C7	-2.23	119.91	122.90
2	A	4	MAN	O5-C5-C6	2.22	111.98	107.66
2	D	3	BMA	C1-O5-C5	2.22	115.16	112.19
2	C	2	NAG	O7-C7-N2	2.20	125.87	121.98
2	D	5	MAN	O5-C5-C4	2.18	116.14	110.83
2	B	1	NAG	O7-C7-N2	2.18	125.83	121.98
2	B	2	NAG	C4-C3-C2	2.16	114.19	111.02
2	C	1	NAG	O3-C3-C4	-2.15	105.30	110.38
2	D	4	MAN	C3-C4-C5	2.13	114.09	110.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	2	NAG	C4-C3-C2	2.11	114.11	111.02
2	D	5	MAN	O5-C5-C6	2.08	111.72	107.66
2	B	2	NAG	C6-C5-C4	-2.06	107.96	113.02
2	D	5	MAN	C2-C3-C4	2.06	114.48	110.86
2	C	3	BMA	O3-C3-C4	-2.02	105.60	110.38

There are no chirality outliers.



All (12) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	5	MAN	C4-C5-C6-O6
2	D	5	MAN	O5-C5-C6-O6
2	B	5	MAN	O5-C5-C6-O6
2	A	5	MAN	O5-C5-C6-O6
2	D	5	MAN	C4-C5-C6-O6
2	C	2	NAG	C4-C5-C6-O6
2	C	2	NAG	O5-C5-C6-O6
2	C	5	MAN	O5-C5-C6-O6
2	B	5	MAN	C4-C5-C6-O6
2	C	3	BMA	C4-C5-C6-O6
2	B	3	BMA	O5-C5-C6-O6
2	C	1	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	E	103/108 (95%)	0.21	2 (1%) 66 63	48, 62, 100, 121	0
1	F	106/108 (98%)	0.25	5 (4%) 36 32	46, 66, 98, 118	0
1	G	108/108 (100%)	0.35	5 (4%) 37 33	48, 66, 105, 134	0
1	H	104/108 (96%)	0.14	2 (1%) 66 63	46, 63, 91, 104	0
All	All	421/432 (97%)	0.24	14 (3%) 49 44	46, 64, 100, 134	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	112	GLY	3.6
1	G	90	LEU	3.5
1	F	30	ASP	3.4
1	F	33	VAL	3.0
1	F	53	ASP	2.7
1	E	51	LEU	2.6
1	G	54	HIS	2.4
1	G	105	ASP	2.4
1	G	10	LEU	2.3
1	H	54	HIS	2.2
1	F	25	ASP	2.1
1	G	93	ASP	2.1
1	H	78	CYS	2.1
1	E	31	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	A	1	14/15	-	-	65,68,71,73	0
2	NAG	A	2	14/15	-	-	72,78,86,87	0
2	BMA	A	3	11/12	-	-	81,85,94,98	0
2	MAN	A	4	11/12	-	-	84,86,89,92	0
2	MAN	A	5	11/12	-	-	99,101,106,108	0
2	NAG	B	1	14/15	-	-	72,75,80,83	0
2	NAG	B	2	14/15	-	-	86,92,98,98	0
2	BMA	B	3	11/12	-	-	97,102,109,114	0
2	MAN	B	4	11/12	-	-	97,99,103,105	0
2	MAN	B	5	11/12	-	-	108,108,112,116	0
2	NAG	C	1	14/15	-	-	61,65,68,72	0
2	NAG	C	2	14/15	-	-	74,81,86,87	0
2	BMA	C	3	11/12	-	-	87,91,99,103	0
2	MAN	C	4	11/12	-	-	90,91,96,97	0
2	MAN	C	5	11/12	-	-	102,104,109,110	0
2	NAG	D	1	14/15	-	-	63,66,70,74	0
2	NAG	D	2	14/15	-	-	75,82,88,89	0
2	BMA	D	3	11/12	-	-	86,90,97,101	0
2	MAN	D	4	11/12	-	-	88,89,94,96	0
2	MAN	D	5	11/12	-	-	102,104,108,108	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain A:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain B:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.