



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:03 AM UTC

PDB ID : 7OM2 / pdb_00007om2
Title : Thosea asigna virus RdRP domain in complex with Mg+2
Authors : Ferrero, D.S.; Falqui, M.; Verdaguer, N.
Deposited on : 2021-05-21
Resolution : 2.07 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3774 (2.08-2.04)
Clashscore	190562	3883 (2.08-2.04)
Ramachandran outliers	187476	3860 (2.08-2.04)
Sidechain outliers	187428	3860 (2.08-2.04)
RSRZ outliers	180081	3775 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	684	5% 88% 9% .
1	B	684	5% 88% 8% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11628 atoms, of which 8 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RNA-dependent RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	659	Total	C	N	O	S	0	0	0
			5221	3314	905	976	26			
1	B	661	Total	C	N	O	S	0	1	0
			5247	3329	912	980	26			

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	MET	-	initiating methionine	UNP Q6A562
A	-11	GLY	-	expression tag	UNP Q6A562
A	-10	SER	-	expression tag	UNP Q6A562
A	-9	SER	-	expression tag	UNP Q6A562
A	-8	HIS	-	expression tag	UNP Q6A562
A	-7	HIS	-	expression tag	UNP Q6A562
A	-6	HIS	-	expression tag	UNP Q6A562
A	-5	HIS	-	expression tag	UNP Q6A562
A	-4	HIS	-	expression tag	UNP Q6A562
A	-3	HIS	-	expression tag	UNP Q6A562
A	-2	SER	-	expression tag	UNP Q6A562
A	-1	GLN	-	expression tag	UNP Q6A562
A	0	ASP	-	expression tag	UNP Q6A562
A	1	LEU	-	expression tag	UNP Q6A562
A	2	GLU	-	expression tag	UNP Q6A562
A	3	ASN	-	expression tag	UNP Q6A562
A	4	LEU	-	expression tag	UNP Q6A562
A	5	TYR	-	expression tag	UNP Q6A562
A	6	PHE	-	expression tag	UNP Q6A562
A	7	GLN	-	expression tag	UNP Q6A562
A	8	GLY	-	expression tag	UNP Q6A562
A	9	GLY	-	expression tag	UNP Q6A562
A	10	SER	-	expression tag	UNP Q6A562
B	-12	MET	-	initiating methionine	UNP Q6A562
B	-11	GLY	-	expression tag	UNP Q6A562

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-10	SER	-	expression tag	UNP Q6A562
B	-9	SER	-	expression tag	UNP Q6A562
B	-8	HIS	-	expression tag	UNP Q6A562
B	-7	HIS	-	expression tag	UNP Q6A562
B	-6	HIS	-	expression tag	UNP Q6A562
B	-5	HIS	-	expression tag	UNP Q6A562
B	-4	HIS	-	expression tag	UNP Q6A562
B	-3	HIS	-	expression tag	UNP Q6A562
B	-2	SER	-	expression tag	UNP Q6A562
B	-1	GLN	-	expression tag	UNP Q6A562
B	0	ASP	-	expression tag	UNP Q6A562
B	1	LEU	-	expression tag	UNP Q6A562
B	2	GLU	-	expression tag	UNP Q6A562
B	3	ASN	-	expression tag	UNP Q6A562
B	4	LEU	-	expression tag	UNP Q6A562
B	5	TYR	-	expression tag	UNP Q6A562
B	6	PHE	-	expression tag	UNP Q6A562
B	7	GLN	-	expression tag	UNP Q6A562
B	8	GLY	-	expression tag	UNP Q6A562
B	9	GLY	-	expression tag	UNP Q6A562
B	10	SER	-	expression tag	UNP Q6A562

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	B	1	Total Mg 1 1	0	0

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			14	3	8	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	604	Total	O	0	0
			604	604		
5	B	505	Total	O	0	0
			505	505		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RNA-dependent RNA polymerase



• Molecule 1: RNA-dependent RNA polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	134.84Å 149.00Å 100.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	80.51 – 2.07 80.51 – 2.07	Depositor EDS
% Data completeness (in resolution range)	94.4 (80.51-2.07) 94.4 (80.51-2.07)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.54 (at 2.07Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158, PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.172 , 0.204 0.176 , 0.208	Depositor DCC
R_{free} test set	5631 reflections (4.55%)	wwPDB-VP
Wilson B-factor (Å ²)	30.9	Xtriage
Anisotropy	0.343	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 44.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11628	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	2/5339 (0.0%)	0.59	0/7229
1	B	0.37	0/5366	0.58	1/7267 (0.0%)
All	All	0.40	2/10705 (0.0%)	0.59	1/14496 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	251	LEU	C-N	6.21	1.41	1.33
1	A	340	GLY	C-N	5.99	1.42	1.33

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	507	TRP	N-CA-C	-5.53	104.95	110.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5221	0	5178	40	0
1	B	5247	0	5206	46	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1	0	0	0	0
3	A	20	0	0	0	0
3	B	15	0	0	0	0
4	A	6	8	8	0	0
5	A	604	0	0	9	0
5	B	505	0	0	9	0
All	All	11620	8	10392	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:553:THR:H	1:B:556:MET:HE3	1.36	0.89
1:B:499:GLU:OE2	5:B:801:HOH:O	2.06	0.71
1:B:246:LYS:NZ	5:B:802:HOH:O	2.23	0.70
1:B:252:ARG:NH2	1:B:413:GLU:OE1	2.20	0.70
1:A:12:ARG:HG2	1:A:13:LEU:HD22	1.75	0.69
1:A:618:GLY:HA2	5:A:914:HOH:O	1.92	0.67
1:A:372:GLN:NE2	5:A:802:HOH:O	2.27	0.67
1:A:473:LYS:HE3	1:A:474:TYR:CE2	2.29	0.67
1:B:617:ASP:OD1	1:B:619:VAL:HG12	1.95	0.67
1:B:137:ARG:HH11	1:B:137:ARG:HG2	1.60	0.66
1:B:580:ARG:NH2	5:B:806:HOH:O	2.27	0.66
1:A:250:THR:OG1	5:A:801:HOH:O	2.12	0.66
1:B:476:ILE:HG12	1:B:487:ILE:HB	1.79	0.65
1:B:30:LEU:HD11	1:B:555:THR:HG21	1.77	0.65
1:B:410:LEU:O	1:B:414:MET:HG3	1.98	0.63
1:A:225:ARG:HD2	5:B:1229:HOH:O	1.98	0.63
1:A:605:ARG:NH1	1:A:623:PRO:HA	2.14	0.62
1:B:280:ARG:HD3	5:B:822:HOH:O	1.98	0.62
1:A:373:MET:HE1	1:A:447:ASP:O	2.00	0.61
1:B:433:LYS:NZ	5:B:805:HOH:O	2.25	0.61
1:B:553:THR:H	1:B:556:MET:CE	2.13	0.61
1:B:553:THR:N	1:B:556:MET:HE3	2.12	0.61
1:B:617:ASP:CG	1:B:619:VAL:HG12	2.27	0.60
1:A:184:LYS:HE3	5:A:941:HOH:O	2.01	0.60
1:A:605:ARG:HG3	5:A:845:HOH:O	2.03	0.59
1:A:391:ASP:HA	1:A:394:VAL:HG22	1.85	0.58
1:A:315:ASN:OD1	1:A:317:TYR:HB3	2.04	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:599:VAL:O	1:B:604:GLY:HA2	2.05	0.57
1:A:599:VAL:O	1:A:604:GLY:HA2	2.06	0.56
1:A:151:THR:HG23	5:A:934:HOH:O	2.06	0.55
1:A:225:ARG:O	1:B:22:ARG:HD2	2.07	0.54
1:A:12:ARG:NH2	5:A:814:HOH:O	2.41	0.53
1:B:472:GLU:HG3	1:B:492:TRP:HB3	1.91	0.52
1:B:137:ARG:NH1	5:B:814:HOH:O	2.42	0.52
1:B:315:ASN:OD1	1:B:317:TYR:HB3	2.10	0.51
1:B:368:PRO:HG2	1:B:455:TYR:OH	2.11	0.51
1:B:532:LEU:HB2	1:B:537:TRP:CE2	2.46	0.50
1:A:605:ARG:HD2	1:A:622:PHE:O	2.12	0.50
1:B:373:MET:HE1	1:B:447:ASP:O	2.10	0.50
1:B:414:MET:HA	1:B:418:SER:OG	2.11	0.50
1:A:139:GLU:HG3	1:A:534:PHE:HZ	1.76	0.49
1:B:228:LEU:HB3	1:B:229:PRO:HD3	1.94	0.49
1:A:27:ARG:HD3	1:A:596:CYS:O	2.13	0.48
1:A:605:ARG:HH11	1:A:623:PRO:HA	1.76	0.48
1:B:619:VAL:HG13	1:B:619:VAL:O	2.12	0.48
1:B:473:LYS:HE2	1:B:474:TYR:CE2	2.49	0.48
1:A:228:LEU:HB3	1:A:229:PRO:HD3	1.96	0.47
1:A:246:LYS:HD2	1:B:671:ILE:HG22	1.96	0.47
1:A:139:GLU:HG3	1:A:534:PHE:CZ	2.49	0.47
1:B:280:ARG:HG2	1:B:280:ARG:NH1	2.30	0.47
1:B:641:ASP:O	1:B:642:HIS:C	2.58	0.47
1:B:553:THR:OG1	1:B:556:MET:HE2	2.16	0.46
1:B:280:ARG:HG2	1:B:280:ARG:HH11	1.81	0.46
1:A:532:LEU:HB2	1:A:537:TRP:CE2	2.50	0.46
1:A:358:ARG:HA	1:A:362:LYS:O	2.15	0.46
1:A:332:TRP:CE2	1:A:345:ARG:HD2	2.51	0.45
1:A:368:PRO:HG2	1:A:455:TYR:OH	2.17	0.45
1:B:648:ARG:HD3	5:B:880:HOH:O	2.17	0.45
1:B:164:ARG:HH21	1:B:372:GLN:HG2	1.80	0.45
1:A:79:GLU:HG2	5:A:1206:HOH:O	2.17	0.44
1:A:30:LEU:HD11	1:A:555:THR:HG21	2.00	0.44
1:A:364:TYR:CE2	1:A:496:LEU:HD13	2.52	0.44
1:B:395:LYS:HE3	5:B:1250:HOH:O	2.18	0.44
1:A:617:ASP:CG	1:A:619:VAL:HG12	2.43	0.44
1:A:121:GLN:HG2	1:A:155:ASP:OD2	2.18	0.44
1:A:332:TRP:NE1	1:A:345:ARG:HD2	2.33	0.44
1:B:61:ALA:O	1:B:65:THR:HG23	2.19	0.43
1:B:552:GLU:HG2	1:B:557:ARG:HG3	1.99	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:137:ARG:HG2	1:B:137:ARG:NH1	2.32	0.43
1:A:13:LEU:HD21	1:B:212:ALA:HB2	2.01	0.43
1:B:661:ARG:HH11	1:B:661:ARG:HB2	1.84	0.43
1:B:139:GLU:HG3	1:B:534:PHE:HZ	1.85	0.42
1:B:169:THR:HG22	1:B:430:LYS:HD3	2.01	0.42
1:B:473:LYS:HE2	1:B:474:TYR:CZ	2.55	0.42
1:A:431:MET:HE3	5:A:954:HOH:O	2.19	0.42
1:A:372:GLN:O	1:A:372:GLN:HG3	2.19	0.41
1:A:414:MET:HA	1:A:418:SER:OG	2.20	0.41
1:B:332:TRP:CE2	1:B:345:ARG:HD2	2.56	0.41
1:A:250:THR:O	1:A:254:LYS:HD3	2.20	0.41
1:A:210:THR:CG2	1:B:18:MET:HE1	2.51	0.40
1:A:521:ARG:HD3	1:A:523:GLU:O	2.21	0.40
1:B:164:ARG:HA	1:B:164:ARG:HD2	1.79	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	655/684 (96%)	644 (98%)	11 (2%)	0	100	100
1	B	660/684 (96%)	649 (98%)	10 (2%)	1 (0%)	43	38
All	All	1315/1368 (96%)	1293 (98%)	21 (2%)	1 (0%)	48	44

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	642	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	559/581 (96%)	554 (99%)	5 (1%)	70	74
1	B	562/581 (97%)	557 (99%)	5 (1%)	70	74
All	All	1121/1162 (96%)	1111 (99%)	10 (1%)	70	74

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	129	SER
1	A	341	LYS
1	A	366	ILE
1	A	430	LYS
1	A	487	ILE
1	B	366	ILE
1	B	371	LYS
1	B	548	TYR
1	B	552	GLU
1	B	641	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	76	GLN
1	B	185	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	B	704	-	4,4,4	0.27	0	6,6,6	0.10	0
3	SO4	A	704	-	4,4,4	0.33	0	6,6,6	0.24	0
3	SO4	B	702	-	4,4,4	0.22	0	6,6,6	0.23	0
3	SO4	A	702	-	4,4,4	0.28	0	6,6,6	0.23	0
3	SO4	B	703	-	4,4,4	0.22	0	6,6,6	0.29	0
3	SO4	A	703	-	4,4,4	0.29	0	6,6,6	0.27	0
4	GOL	A	706	-	5,5,5	0.89	0	5,5,5	1.02	0
3	SO4	A	705	-	4,4,4	0.33	0	6,6,6	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	706	-	-	4/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	706	GOL	O1-C1-C2-C3
4	A	706	GOL	C1-C2-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	706	GOL	O1-C1-C2-O2
4	A	706	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	659/684 (96%)	-0.08	32 (4%) 35 35	19, 29, 56, 131	0
1	B	661/684 (96%)	0.01	32 (4%) 35 35	20, 32, 60, 130	1 (0%)
All	All	1320/1368 (96%)	-0.04	64 (4%) 35 35	19, 31, 59, 131	1 (0%)

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	128	THR	8.0
1	B	126	VAL	7.6
1	B	642	HIS	6.2
1	B	548	TYR	5.8
1	B	128	THR	5.6
1	A	24	MET	5.6
1	A	642	HIS	5.2
1	A	548	TYR	5.0
1	A	125	GLY	5.0
1	B	11	THR	4.9
1	B	145[A]	ARG	4.8
1	A	372	GLN	4.3
1	A	547	LYS	4.3
1	A	11	THR	4.2
1	B	124	ALA	4.1
1	B	643	PRO	3.9
1	A	640	ARG	3.9
1	B	127	ASN	3.8
1	A	551	LYS	3.7
1	B	618	GLY	3.6
1	A	489	GLU	3.5
1	A	546	SER	3.4
1	A	641	ASP	3.4
1	A	124	ALA	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	24	MET	3.1
1	A	618	GLY	2.9
1	B	38	ALA	2.9
1	A	549	ARG	2.8
1	B	547	LYS	2.8
1	B	641	ASP	2.8
1	A	643	PRO	2.8
1	B	546	SER	2.8
1	B	167	MET	2.7
1	B	13	LEU	2.7
1	B	639	SER	2.7
1	B	550	SER	2.6
1	B	549	ARG	2.6
1	B	125	GLY	2.6
1	B	55	GLU	2.6
1	B	524	ASN	2.6
1	B	551	LYS	2.6
1	A	341	LYS	2.6
1	A	129	SER	2.6
1	B	79	GLU	2.5
1	A	617	ASP	2.5
1	A	550	SER	2.5
1	A	619	VAL	2.5
1	B	617	ASP	2.4
1	A	54	ARG	2.3
1	B	12	ARG	2.3
1	A	121	GLN	2.3
1	A	13	LEU	2.3
1	A	600	GLN	2.2
1	A	522	ARG	2.2
1	A	23	ALA	2.2
1	A	523	GLU	2.2
1	B	523	GLU	2.2
1	B	640	ARG	2.1
1	A	524	ASN	2.1
1	B	41	ASP	2.1
1	B	644	CYS	2.1
1	B	603	GLY	2.1
1	A	167	MET	2.0
1	A	280	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	A	706	6/6	0.87	0.14	51,68,80,82	0
3	SO4	A	704	5/5	0.92	0.08	45,54,57,67	0
3	SO4	A	705	5/5	0.94	0.08	44,47,65,71	0
3	SO4	B	703	5/5	0.95	0.09	37,52,63,73	0
3	SO4	B	702	5/5	0.95	0.10	45,47,56,58	0
3	SO4	A	702	5/5	0.96	0.08	46,48,55,57	0
3	SO4	B	704	5/5	0.97	0.10	48,49,56,58	0
3	SO4	A	703	5/5	0.98	0.11	37,43,47,57	0
2	MG	A	701	1/1	1.00	0.02	27,27,27,27	0
2	MG	B	701	1/1	1.00	0.02	31,31,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.