



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 04:00 AM UTC

PDB ID : 8OMG / pdb_00008omg
Title : Crystal structure of mKHK (apo)
Authors : Ebenhoch, E.; Pautsch, P.
Deposited on : 2023-03-31
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	298	28% 87% 7% 5%
1	B	298	4% 93% 6%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4676 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ketohexokinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	282	Total	C	N	O	S	0	1	0
			2181	1365	388	414	14			
1	B	294	Total	C	N	O	S	0	0	0
			2264	1420	403	428	13			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	203	SER	PRO	conflict	UNP P97328
B	203	SER	PRO	conflict	UNP P97328


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	61	Total	O	0	0
			61	61		
2	B	170	Total	O	0	0
			170	170		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Ketohexokinase



• Molecule 1: Ketohexokinase



4 Data and refinement statistics

Property	Value	Source
Space group	P 6	Depositor
Cell constants a, b, c, α , β , γ	154.39Å 154.39Å 50.07Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	66.85 – 1.82 66.85 – 1.82	Depositor EDS
% Data completeness (in resolution range)	64.9 (66.85-1.82) 64.9 (66.85-1.82)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.85 (at 1.82Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.225 , 0.249 (Not available) , 0.234	Depositor DCC
R_{free} test set	2060 reflections (3.36%)	wwPDB-VP
Wilson B-factor (Å ²)	31.3	Xtriage
Anisotropy	0.103	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 43.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.038 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4676	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	0/2216	1.19	4/2990 (0.1%)
1	B	0.78	0/2305	1.16	4/3116 (0.1%)
All	All	0.77	0/4521	1.17	8/6106 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	177	GLU	CA-C-N	6.45	129.21	120.38
1	A	177	GLU	C-N-CA	6.45	129.21	120.38
1	B	176	ARG	CA-C-N	6.19	128.90	120.54
1	B	176	ARG	C-N-CA	6.19	128.90	120.54
1	A	82	ASP	CA-CB-CG	6.03	118.63	112.60
1	B	82	ASP	CA-CB-CG	5.38	117.98	112.60
1	B	76	PHE	CA-CB-CG	5.15	118.95	113.80
1	A	76	PHE	CA-CB-CG	5.08	118.88	113.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2181	0	2159	5	0
1	B	2264	0	2248	6	0
2	A	61	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	170	0	0	0	0
All	All	4676	0	4407	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:220:THR:HG21	1:B:274:MET:HE3	1.81	0.63
1:A:220:THR:HG21	1:A:274:MET:HE3	1.82	0.60
1:A:42:ASN:HB2	1:A:139:GLU:CD	2.31	0.54
1:A:16:ILE:HD13	1:B:16:ILE:HD13	1.94	0.48
1:B:42:ASN:HB2	1:B:139:GLU:CD	2.40	0.47
1:A:39:ARG:HG3	1:A:72:LEU:HD22	1.97	0.46
1:B:13:VAL:HG23	1:B:95:PRO:HB2	1.98	0.46
1:B:39:ARG:HG3	1:B:72:LEU:HD22	1.98	0.46
1:B:64:ALA:HB2	1:B:117:LEU:HD11	2.00	0.43
1:A:225:TRP:HB3	1:A:228:GLU:HB2	2.01	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	200/298 (67%)	194 (97%)	5 (2%)	1 (0%)	24	14
1	B	273/298 (92%)	268 (98%)	5 (2%)	0	100	100
All	All	473/596 (79%)	462 (98%)	10 (2%)	1 (0%)	43	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	210	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/251 (95%)	229 (96%)	9 (4%)	29	11
1	B	247/251 (98%)	244 (99%)	3 (1%)	63	54
All	All	485/502 (97%)	473 (98%)	12 (2%)	42	25

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	79	ARG
1	A	112	LEU
1	A	162	LEU
1	A	173	GLU
1	A	208	LEU
1	A	209	ARG
1	A	211	LEU
1	A	214	ARG
1	A	292	GLN
1	B	79	ARG
1	B	173	GLU
1	B	297	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	198	HIS
1	B	18	ASN
1	B	35	GLN
1	B	151	GLN
1	B	160	GLN
1	B	238	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	282/298 (94%)	1.56	83 (29%) 1 1	20, 63, 101, 123	1 (0%)
1	B	294/298 (98%)	0.36	13 (4%) 39 45	27, 39, 66, 89	0
All	All	576/596 (96%)	0.95	96 (16%) 4 5	20, 48, 91, 123	1 (0%)

All (96) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	112	LEU	6.6
1	A	205	VAL	5.5
1	A	289	CYS	5.3
1	B	297	ILE	5.2
1	A	163	PRO	4.8
1	A	297	ILE	4.5
1	A	254	LEU	4.3
1	A	208	LEU	4.3
1	A	103	ASN	4.3
1	A	162	LEU	4.2
1	B	250	VAL	4.1
1	A	198	HIS	4.0
1	A	241	HIS	4.0
1	A	294	PHE	4.0
1	A	161	PRO	4.0
1	A	166	VAL	3.9
1	A	225	TRP	3.8
1	A	204	ALA	3.6
1	A	226	ALA	3.6
1	A	196	ALA	3.5
1	A	230	ALA	3.5
1	A	295	ASP	3.5
1	A	240	LEU	3.4
1	A	211	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	291	LEU	3.3
1	A	175	PRO	3.2
1	A	217	LYS	3.2
1	A	239	LEU	3.1
1	B	248	PRO	3.1
1	A	218	GLY	3.0
1	A	21	ASP	3.0
1	A	134	LYS	3.0
1	A	233	LEU	3.0
1	A	135	TRP	2.9
1	A	253	THR	2.9
1	A	203	SER	2.9
1	A	215	VAL	2.9
1	A	293	GLY	2.9
1	A	201	PHE	2.9
1	A	174	LYS	2.9
1	A	106	GLY	2.9
1	A	153	ILE	2.8
1	A	165	LYS	2.8
1	A	184	SER	2.8
1	A	224	ALA	2.8
1	A	232	ALA	2.8
1	A	268	LEU	2.8
1	A	91	GLN	2.7
1	A	195	VAL	2.7
1	A	77	ARG	2.6
1	A	36	ARG	2.6
1	A	108	ARG	2.6
1	A	207	ALA	2.6
1	A	282	CYS	2.6
1	A	280	PHE	2.6
1	A	130	LEU	2.6
1	B	162	LEU	2.6
1	A	236	ASP	2.6
1	A	202	GLN	2.6
1	A	271	GLY	2.6
1	B	247	PRO	2.5
1	A	287	LYS	2.5
1	A	221	LEU	2.5
1	A	167	ARG	2.5
1	A	141	ARG	2.4
1	A	229	GLY	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	93	ASP	2.4
1	A	56	ARG	2.4
1	A	279	ARG	2.3
1	A	283	GLN	2.3
1	A	237	GLY	2.3
1	A	252	ASP	2.3
1	A	227	GLU	2.3
1	B	251	VAL	2.3
1	A	219	ALA	2.2
1	A	107	SER	2.2
1	B	29	ASP	2.2
1	B	33	LEU	2.2
1	A	125	PHE	2.2
1	A	11	LEU	2.2
1	A	186	GLY	2.2
1	A	210	GLY	2.2
1	A	288	LYS	2.1
1	A	158	ALA	2.1
1	A	277	ALA	2.1
1	A	164	GLN	2.1
1	A	292	GLN	2.1
1	B	27	ASP	2.1
1	A	104	SER	2.1
1	A	113	TYR	2.1
1	B	104	SER	2.1
1	A	285	ALA	2.1
1	A	132	ARG	2.1
1	B	116	ASN	2.0
1	B	28	THR	2.0
1	B	296	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.