



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:10 PM UTC

PDB ID : 2ONY / pdb_00002ony
Title : Structure of hPNMT with inhibitor 7-(N-4-chlorophenylaminosulfonyl)-THI
Q and AdoHcy
Authors : Drinkwater, N.; Martin, J.L.
Deposited on : 2007-01-24
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	
1	B	289	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4335 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phenylethanolamine N-methyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	2	0	0
			2016	1280	358	369	9			
1	B	268	Total	C	N	O	S	0	0	0
			2083	1320	370	384	9			

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	283	GLU	-	expression tag	UNP P11086
A	284	HIS	-	expression tag	UNP P11086
A	285	HIS	-	expression tag	UNP P11086
A	286	HIS	-	expression tag	UNP P11086
A	287	HIS	-	expression tag	UNP P11086
A	288	HIS	-	expression tag	UNP P11086
A	289	HIS	-	expression tag	UNP P11086
B	783	GLU	-	expression tag	UNP P11086
B	784	HIS	-	expression tag	UNP P11086
B	785	HIS	-	expression tag	UNP P11086
B	786	HIS	-	expression tag	UNP P11086
B	787	HIS	-	expression tag	UNP P11086
B	788	HIS	-	expression tag	UNP P11086
B	789	HIS	-	expression tag	UNP P11086

- Molecule 2 is S-ADENOSYL-L-HOMOCYSTEINE (CCD ID: SAH) (formula: C₁₄H₂₀N₆O₅S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 26	C 14	N 6	O 5	S 1	0	0
2	B	1	Total 26	C 14	N 6	O 5	S 1	0	0

- Molecule 3 is N-(4-CHLOROPHENYL)-1,2,3,4-TETRAHYDROISOQUINOLINE-7-SULFONAMIDE (CCD ID: TMJ) (formula: $C_{15}H_{15}ClN_2O_2S$).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	S	0	0
			21	15	1	2	2	1		

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	74	Total	O	0	0
			74	74		
5	B	84	Total	O	0	0
			84	84		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phenylethanolamine N-methyltransferase



- Molecule 1: Phenylethanolamine N-methyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	93.96Å 93.96Å 188.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.58 – 2.60 45.58 – 2.60	Depositor EDS
% Data completeness (in resolution range)	94.7 (45.58-2.60) 94.8 (45.58-2.60)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 2.61Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.223 , 0.258 0.223 , 0.259	Depositor DCC
R_{free} test set	1288 reflections (4.83%)	wwPDB-VP
Wilson B-factor (Å ²)	68.1	Xtriage
Anisotropy	0.429	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 43.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4335	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, TMJ, SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/2067	0.93	3/2813 (0.1%)
1	B	0.46	0/2136	0.95	9/2908 (0.3%)
All	All	0.44	0/4203	0.94	12/5721 (0.2%)

There are no bond length outliers.

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	163	GLN	CA-C-N	6.26	127.67	119.84
1	A	163	GLN	C-N-CA	6.26	127.67	119.84
1	B	688	SER	CA-C-N	6.08	125.30	118.97
1	B	688	SER	C-N-CA	6.08	125.30	118.97
1	A	103	LEU	N-CA-C	5.62	119.07	110.20
1	B	539	ASN	N-CA-C	5.52	120.67	113.88
1	B	663	GLN	CA-C-N	5.38	126.57	119.84
1	B	663	GLN	C-N-CA	5.38	126.57	119.84
1	B	574	THR	N-CA-C	5.27	118.41	109.76
1	B	726	GLU	N-CA-C	-5.24	106.40	112.89
1	B	764	THR	N-CA-C	5.23	119.30	113.02
1	B	541	ALA	N-CA-C	-5.07	102.68	110.58

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2016	0	1983	43	0
1	B	2083	0	2045	37	0
2	A	26	0	19	2	0
2	B	26	0	19	0	0
3	A	21	0	15	0	0
4	B	5	0	0	0	0
5	A	74	0	0	1	0
5	B	84	0	0	3	0
All	All	4335	0	4081	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124:SER:O	1:A:128:GLN:HG3	1.85	0.75
1:A:59:ARG:O	1:A:63:GLN:HG3	1.87	0.73
1:B:695:GLN:HE22	1:B:742:ALA:HA	1.58	0.69
1:B:569:GLU:OE1	1:B:751:ARG:NH2	2.26	0.68
1:A:74:THR:HG22	1:A:96:ASP:HB3	1.77	0.67
1:A:237:GLU:O	1:A:241:GLU:HG3	1.95	0.66
1:B:537:ARG:NH1	5:B:4005:HOH:O	2.28	0.65
1:B:685:GLU:O	1:B:731:VAL:HG13	1.97	0.65
1:A:91:CYS:HA	1:A:97:ILE:HD11	1.80	0.63
1:B:646:GLN:O	1:B:650:ARG:HG3	2.00	0.61
1:A:94:PHE:O	1:A:150:ARG:HD3	2.03	0.59
1:A:28:GLN:NE2	1:A:28:GLN:HA	2.19	0.57
1:B:638:GLU:OE1	1:B:643:LYS:HE3	2.05	0.57
1:A:205:LEU:HD11	1:A:209:GLY:HA3	1.87	0.57
1:B:517:ALA:HB3	1:B:518:PRO:HD3	1.87	0.56
1:A:31:GLU:OE2	1:A:226:GLU:HG2	2.06	0.56
1:B:575:LEU:HD11	1:B:679:VAL:HG23	1.87	0.55
1:A:96:ASP:OD1	1:A:152:LYS:HG3	2.07	0.55
1:A:158:ASP:OD1	1:A:160:HIS:HB2	2.07	0.55
1:A:49:ASN:O	1:A:55:PRO:HG2	2.08	0.54
1:A:185:GLU:O	1:A:231:VAL:HG13	2.08	0.54
1:B:721:TRP:HA	1:B:729:LEU:O	2.08	0.53
1:B:618:PRO:C	1:B:620:ALA:H	2.17	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:128:GLN:HB3	1:B:628:GLN:OE1	2.09	0.52
1:B:559:ARG:O	1:B:563:GLN:HG3	2.09	0.51
1:A:36:LEU:HD21	1:A:82:PRO:HG2	1.92	0.51
1:B:583:THR:HA	1:B:623:TRP:CZ2	2.46	0.50
1:A:32:PRO:HG3	1:A:106:ASN:ND2	2.27	0.50
1:A:57:LYS:HD3	1:A:214:ILE:HG21	1.94	0.50
1:A:125:MET:HB2	5:A:3009:HOH:O	2.12	0.49
1:A:100:THR:O	1:A:101:ASP:HB2	2.12	0.49
1:B:723:LEU:HD23	1:B:728:ARG:HA	1.95	0.49
1:A:175:ALA:O	1:A:206:ARG:HB3	2.12	0.48
1:A:91:CYS:O	1:A:150:ARG:HD2	2.12	0.48
1:B:520:GLN:HG3	5:B:4052:HOH:O	2.12	0.48
1:B:740:ARG:HD3	5:B:4067:HOH:O	2.14	0.47
1:A:174:PRO:HG2	1:A:206:ARG:HB2	1.95	0.47
1:A:88:LEU:HD13	1:A:127:SER:HA	1.96	0.46
1:B:594:PHE:O	1:B:650:ARG:HD3	2.15	0.46
1:B:516:SER:HB2	1:B:520:GLN:NE2	2.30	0.46
1:A:128:GLN:HG2	1:A:140:TRP:CD1	2.51	0.46
1:A:83:THR:HA	1:A:123:TRP:CZ2	2.51	0.46
1:B:549:ASN:O	1:B:555:PRO:HG2	2.16	0.46
1:B:670:PRO:O	1:B:671:ALA:C	2.58	0.45
1:B:576:ILE:HG12	1:B:598:THR:HB	1.98	0.45
1:B:671:ALA:HA	1:B:672:PRO:HD3	1.83	0.45
1:B:640:TRP:O	1:B:644:GLU:HG3	2.16	0.45
1:A:129:HIS:CD2	1:A:133:ILE:HD11	2.52	0.45
1:B:574:THR:HG21	1:B:672:PRO:HB2	1.99	0.45
1:A:229:LEU:HD22	1:A:229:LEU:N	2.32	0.44
1:B:759:PRO:HD2	1:B:762:LEU:HD12	2.00	0.44
1:A:189:PRO:HD2	1:A:193:SER:CB	2.47	0.44
1:B:596:ASP:C	1:B:596:ASP:OD2	2.61	0.44
1:A:243:LEU:O	1:A:248:TYR:HB2	2.18	0.44
1:A:121:PHE:HD2	1:A:123:TRP:CE2	2.37	0.43
1:A:91:CYS:HB3	1:A:147:LEU:HD13	2.00	0.43
1:B:732:VAL:O	1:B:732:VAL:HG13	2.19	0.43
1:B:556:TRP:CE2	1:B:756:TYR:HB2	2.54	0.43
1:B:565:PHE:CZ	1:B:590:ALA:HB2	2.53	0.43
1:B:559:ARG:HG2	1:B:563:GLN:NE2	2.34	0.42
1:A:182:PHE:HA	1:A:185:GLU:OE2	2.20	0.42
1:B:576:ILE:HB	1:B:678:LEU:HD23	2.02	0.41
1:B:720:SER:HA	1:B:731:VAL:O	2.21	0.41
1:A:253:LEU:HD13	1:A:275:ALA:HB2	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:112:ARG:HG3	1:A:112:ARG:HH11	1.85	0.41
1:A:195:GLN:HE22	1:A:245:ARG:HB2	1.86	0.41
1:B:767:ASP:OD2	1:B:769:VAL:HG12	2.20	0.41
1:A:85:TYR:CG	1:A:86:GLN:N	2.88	0.41
1:A:112:ARG:HG3	1:A:112:ARG:NH1	2.35	0.41
1:A:159:VAL:HG22	2:A:3001:SAH:N1	2.36	0.41
1:A:181:ALA:O	2:A:3001:SAH:N	2.54	0.41
1:A:98:THR:OG1	1:A:153:ARG:NH1	2.54	0.41
1:A:28:GLN:HA	1:A:28:GLN:HE21	1.85	0.41
1:A:41:ALA:O	1:A:42:PRO:C	2.63	0.41
1:A:188:SER:HB3	1:A:193:SER:HB2	2.02	0.41
1:B:598:THR:HG23	1:B:653:ARG:HG3	2.03	0.41
1:B:606:ASN:HD22	1:B:606:ASN:HA	1.58	0.41
1:B:753:LEU:HD13	1:B:775:ALA:HB2	2.03	0.41
1:B:628:GLN:HE21	1:B:628:GLN:HA	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/289 (88%)	241 (94%)	14 (6%)	0	100	100
1	B	266/289 (92%)	256 (96%)	10 (4%)	0	100	100
All	All	521/578 (90%)	497 (95%)	24 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/233 (91%)	205 (97%)	7 (3%)	33	61
1	B	218/233 (94%)	215 (99%)	3 (1%)	59	81
All	All	430/466 (92%)	420 (98%)	10 (2%)	44	71

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	LYS
1	A	82	PRO
1	A	106	ASN
1	A	191	LEU
1	A	206	ARG
1	A	230	THR
1	A	232	VAL
1	B	582	PRO
1	B	606	ASN
1	B	628	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	28	GLN
1	A	106	ASN
1	A	108	GLN
1	A	129	HIS
1	A	163	GLN
1	A	195	GLN
1	A	210	HIS
1	A	263	GLN
1	A	278	GLN
1	B	520	GLN
1	B	593	HIS
1	B	606	ASN
1	B	608	GLN
1	B	628	GLN
1	B	629	HIS
1	B	695	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	PO4	B	4001	-	4,4,4	1.78	2 (50%)	6,6,6	0.49	0
2	SAH	B	3002	-	27,28,28	0.90	1 (3%)	36,40,40	0.71	0
2	SAH	A	3001	-	27,28,28	0.90	1 (3%)	36,40,40	0.69	0
3	TMJ	A	2001	-	23,23,23	1.52	2 (8%)	32,33,33	1.73	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAH	B	3002	-	-	4/15/31/31	0/3/3/3
2	SAH	A	3001	-	-	7/15/31/31	0/3/3/3
3	TMJ	A	2001	-	-	7/11/18/18	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2001	TMJ	O1-S1	5.98	1.50	1.43
2	A	3001	SAH	C8-N9	3.46	1.43	1.37
2	B	3002	SAH	C8-N9	3.29	1.43	1.37
3	A	2001	TMJ	C10-N2	2.88	1.47	1.43
4	B	4001	PO4	P-O2	-2.09	1.48	1.54
4	B	4001	PO4	P-O4	-2.03	1.48	1.54

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	2001	TMJ	O1-S1-O2	-8.43	109.28	119.52
3	A	2001	TMJ	C8-S1-N2	2.43	109.89	106.88

There are no chirality outliers.

All (18) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	2001	TMJ	C10-N2-S1-O1
3	A	2001	TMJ	C10-N2-S1-O2
3	A	2001	TMJ	C10-N2-S1-C8
3	A	2001	TMJ	C9-C8-S1-O2
3	A	2001	TMJ	C7-C8-S1-O2
2	A	3001	SAH	C2'-C1'-N9-C8
2	B	3002	SAH	C2'-C1'-N9-C8
3	A	2001	TMJ	C7-C8-S1-N2
2	B	3002	SAH	CB-CG-SD-C5'
2	A	3001	SAH	OXT-C-CA-CB
2	A	3001	SAH	O-C-CA-CB
2	A	3001	SAH	C2'-C1'-N9-C4
3	A	2001	TMJ	C9-C8-S1-N2
2	B	3002	SAH	C2'-C1'-N9-C4
2	A	3001	SAH	CB-CG-SD-C5'
2	A	3001	SAH	O4'-C1'-N9-C8
2	B	3002	SAH	O4'-C1'-N9-C8
2	A	3001	SAH	OXT-C-CA-N


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	3001	SAH	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	257/289 (88%)	0.28	7 (2%) 56 50	41, 71, 96, 112	9 (3%)
1	B	268/289 (92%)	0.08	9 (3%) 48 42	45, 63, 88, 114	5 (1%)
All	All	525/578 (90%)	0.18	16 (3%) 52 47	41, 66, 92, 114	14 (2%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	24	ALA	4.8
1	B	781	GLY	4.1
1	A	163	GLN	3.1
1	A	26	ALA	2.9
1	B	670	PRO	2.3
1	B	514	PRO	2.2
1	B	731	VAL	2.2
1	B	529	ARG	2.2
1	A	280	VAL	2.2
1	B	730	THR	2.2
1	B	527	TYR	2.2
1	A	245	ARG	2.2
1	A	237	GLU	2.1
1	B	727	ALA	2.1
1	B	766	VAL	2.0
1	A	29	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	TMJ	A	2001	21/21	0.85	0.19	75,77,78,78	0
4	PO4	B	4001	5/5	0.92	0.15	98,99,100,100	0
2	SAH	A	3001	26/26	0.95	0.09	64,72,75,78	0
2	SAH	B	3002	26/26	0.95	0.09	52,58,62,62	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around SAH A 3001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around SAH B 3002:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.