



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:51 AM UTC

PDB ID : 2OOR / pdb_00002oor
Title : Structure of transhydrogenase (dI.NAD⁺)₂(dIII.H₂NADPH)₁ asymmetric complex
Authors : Bhakta, T.; Jackson, J.B.
Deposited on : 2007-01-26
Resolution : 2.32 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7754 (2.34-2.30)
Clashscore	190562	8383 (2.34-2.30)
Ramachandran outliers	187476	8303 (2.34-2.30)
Sidechain outliers	187428	8303 (2.34-2.30)
RSRZ outliers	180081	7760 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	384	41% 95% ...
1	B	384	13% 89% 6% 5%
2	C	174	40% 93% 6% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	TXP	C	400	X	-	-	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6981 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NAD(P) transhydrogenase subunit alpha part 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	378	Total	C	N	O	S	0	0	0
			2779	1753	479	529	18			
1	B	364	Total	C	N	O	S	0	0	0
			2675	1691	463	505	16			

- Molecule 2 is a protein called NAD(P) transhydrogenase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	173	Total	C	N	O	S	0	0	0
			1302	826	215	250	11			

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (CCD ID: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	B	1	Total	C	N	O	P	0	0
			36	15	6	13	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is 1,4,5,6-TETRAHYDRONICOTINAMIDE ADENINE DINUCLEOTIDE PHOSPHATE (CCD ID: TXP) (formula: $C_{21}H_{32}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	35	Total O 35 35	0	0
6	B	45	Total O 45 45	0	0
6	C	11	Total O 11 11	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NAD(P) transhydrogenase subunit alpha part 1




- Molecule 1: NAD(P) transhydrogenase subunit alpha part 1



- Molecule 2: NAD(P) transhydrogenase subunit beta





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	72.19Å 74.45Å 204.84Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	102.60 – 2.32 102.42 – 2.32	Depositor EDS
% Data completeness (in resolution range)	91.0 (102.60-2.32) 90.9 (102.42-2.32)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.73 (at 2.32Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.247 , 0.275 0.253 , 0.281	Depositor DCC
R_{free} test set	2530 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	43.1	Xtriage
Anisotropy	0.345	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 39.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.017 for k,h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6981	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, NAD, TXP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	2/2816 (0.1%)	0.83	0/3816
1	B	0.65	0/2710	0.86	0/3674
2	C	0.65	0/1325	0.91	0/1792
All	All	0.65	2/6851 (0.0%)	0.86	0/9282

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	170	VAL	CA-CB	6.54	1.60	1.53
1	A	281	VAL	CA-CB	5.41	1.61	1.54

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2779	0	2904	2	0
1	B	2675	0	2809	5	0
2	C	1302	0	1297	3	0
3	A	44	0	26	0	0
3	B	36	0	20	0	0
4	B	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	C	48	0	27	0	0
6	A	35	0	0	0	0
6	B	45	0	0	0	0
6	C	11	0	0	0	0
All	All	6981	0	7091	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:142:ASN:HD21	1:B:186:GLN:HE21	1.19	0.89
1:A:70:VAL:HG23	1:A:95:VAL:HG23	1.87	0.57
1:B:142:ASN:ND2	1:B:186:GLN:HE21	1.98	0.56
2:C:53:PRO:HB2	2:C:95:MET:HE2	1.89	0.53
2:C:63:GLN:HG2	2:C:98:LEU:HB3	1.96	0.47
1:B:121:ALA:O	1:B:366:THR:OG1	2.30	0.45
1:B:255:LEU:HD11	1:B:284:MET:HE2	2.01	0.43
1:B:142:ASN:HD21	1:B:186:GLN:NE2	2.01	0.42
1:A:156:PHE:CE2	1:A:158:ARG:HB2	2.54	0.42
2:C:95:MET:HA	2:C:95:MET:HE3	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	376/384 (98%)	355 (94%)	21 (6%)	0	100	100
1	B	360/384 (94%)	345 (96%)	15 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	C	171/174 (98%)	163 (95%)	7 (4%)	1 (1%)	21	25
All	All	907/942 (96%)	863 (95%)	43 (5%)	1 (0%)	48	59

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	182	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/296 (99%)	282 (96%)	11 (4%)	29	43
1	B	284/296 (96%)	265 (93%)	19 (7%)	15	20
2	C	137/138 (99%)	130 (95%)	7 (5%)	21	31
All	All	714/730 (98%)	677 (95%)	37 (5%)	21	30

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	LEU
1	A	96	LEU
1	A	170	VAL
1	A	177	VAL
1	A	221	VAL
1	A	228	THR
1	A	237	LYS
1	A	253	LYS
1	A	281	VAL
1	A	327	VAL
1	A	337	LYS
1	B	9	ARG
1	B	35	VAL
1	B	50	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	74	VAL
1	B	89	LEU
1	B	147	ARG
1	B	176	LEU
1	B	220	THR
1	B	251	VAL
1	B	252	LEU
1	B	281	VAL
1	B	289	VAL
1	B	327	VAL
1	B	339	LEU
1	B	349	LYS
1	B	352	LYS
1	B	362	THR
1	B	366	THR
1	B	375	VAL
2	C	45	ASN
2	C	63	GLN
2	C	73	LEU
2	C	75	LYS
2	C	104	VAL
2	C	187	LEU
2	C	198	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	142	ASN
1	A	186	GLN
1	A	247	GLN
1	A	320	HIS
1	A	338	ASN
1	B	37	GLN
1	B	105	ASN
1	B	186	GLN
1	B	320	HIS
1	B	338	ASN
1	B	376	HIS
2	C	63	GLN
2	C	120	GLN
2	C	182	ASN
2	C	183	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	TXP	C	400	-	51,52,52	3.25	9 (17%)	73,80,80	2.47	17 (23%)
4	GOL	B	700	-	5,5,5	0.40	0	5,5,5	0.42	0
3	NAD	A	500	-	46,48,48	1.82	6 (13%)	64,73,73	1.58	9 (14%)
3	NAD	B	600	-	38,39,48	1.09	3 (7%)	55,60,73	1.68	9 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	TXP	C	400	-	1/1/14/16	13/35/77/77	0/5/5/5
4	GOL	B	700	-	-	2/4/4/4	-
3	NAD	A	500	-	-	4/30/62/62	0/5/5/5
3	NAD	B	600	-	-	6/22/54/62	0/4/4/5

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	400	TXP	C2N-C3N	-14.68	1.35	1.53
5	C	400	TXP	C2N-N1N	-12.25	1.27	1.47
3	A	500	NAD	O7N-C7N	9.52	1.41	1.24
5	C	400	TXP	C4N-C5N	8.42	1.52	1.32
5	C	400	TXP	O7N-C7N	5.03	1.32	1.23
5	C	400	TXP	C3N-C7N	-4.85	1.47	1.53
5	C	400	TXP	PN-O3	3.76	1.63	1.59
5	C	400	TXP	PA-O3	3.08	1.62	1.59
3	A	500	NAD	C2A-N1A	2.78	1.38	1.33
3	B	600	NAD	C2A-N1A	2.77	1.38	1.33
3	A	500	NAD	C2A-N3A	2.75	1.38	1.33
3	B	600	NAD	C2A-N3A	2.71	1.38	1.33
3	A	500	NAD	PA-O3	2.70	1.62	1.59
3	A	500	NAD	C2N-N1N	2.27	1.37	1.35
3	A	500	NAD	PN-O3	2.11	1.61	1.59
3	B	600	NAD	C8A-N7A	2.09	1.35	1.31
5	C	400	TXP	C5A-N7A	-2.09	1.35	1.39
5	C	400	TXP	C8A-N7A	2.06	1.35	1.31

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	400	TXP	C6N-N1N-C2N	9.77	124.46	109.43
5	C	400	TXP	C3N-C4N-C5N	-8.18	109.42	123.33
5	C	400	TXP	C2N-C3N-C4N	8.13	120.23	108.63
5	C	400	TXP	C6N-C5N-C4N	-5.91	111.20	123.06
3	B	600	NAD	N3A-C2A-N1A	-5.76	119.86	128.58
3	A	500	NAD	N3A-C2A-N1A	-5.74	119.89	128.58
5	C	400	TXP	C5A-C4A-N3A	-5.27	119.46	126.72
5	C	400	TXP	N1A-C2A-N3A	-4.88	121.20	128.58
3	A	500	NAD	C5A-C4A-N3A	-4.63	120.33	126.72
3	B	600	NAD	C5A-C4A-N3A	-4.54	120.46	126.72
3	B	600	NAD	N9A-C8A-N7A	-3.86	108.47	113.94
3	A	500	NAD	N9A-C8A-N7A	-3.73	108.64	113.94
5	C	400	TXP	C3N-C7N-N7N	3.68	120.62	116.80
5	C	400	TXP	N9A-C8A-N7A	-3.57	108.87	113.94
3	B	600	NAD	C2A-N3A-C4A	3.54	120.47	111.83
3	A	500	NAD	C2A-N3A-C4A	3.52	120.44	111.83
5	C	400	TXP	C2A-N3A-C4A	3.51	120.39	111.83
3	B	600	NAD	C5A-N7A-C8A	3.49	108.93	103.45
3	A	500	NAD	C5A-N7A-C8A	3.43	108.85	103.45
5	C	400	TXP	N3A-C4A-N9A	3.16	132.54	127.17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	500	NAD	N3A-C4A-N9A	3.14	132.51	127.17
5	C	400	TXP	C4N-C3N-C7N	3.09	118.46	109.80
3	B	600	NAD	N3A-C4A-N9A	3.00	132.27	127.17
5	C	400	TXP	C4A-C5A-N7A	-2.90	107.27	110.58
5	C	400	TXP	C5A-N7A-C8A	2.78	107.81	103.45
3	A	500	NAD	C3N-C7N-N7N	2.45	120.75	117.74
3	A	500	NAD	O4B-C1B-N9A	2.42	112.74	108.09
3	B	600	NAD	C4A-C5A-N7A	-2.39	107.84	110.58
3	B	600	NAD	O4B-C1B-N9A	2.36	112.62	108.09
3	A	500	NAD	C4A-C5A-N7A	-2.33	107.92	110.58
3	B	600	NAD	C3D-C2D-C1D	2.32	105.84	101.46
5	C	400	TXP	OP2-P2-OP3	2.26	116.28	107.80
5	C	400	TXP	C4A-N9A-C8A	2.20	108.05	105.74
5	C	400	TXP	O2A-PA-O3	2.01	112.71	107.27
5	C	400	TXP	C5A-C4A-N9A	2.00	108.00	105.81

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	C	400	TXP	C3N

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	500	NAD	C5D-O5D-PN-O1N
3	A	500	NAD	C5D-O5D-PN-O2N
3	B	600	NAD	PA-O3-PN-O5D
3	B	600	NAD	O4D-C4D-C5D-O5D
5	C	400	TXP	C2'A-C1'A-N1N-C6N
5	C	400	TXP	O4'A-C1'A-N1N-C6N
3	B	600	NAD	C3D-C4D-C5D-O5D
5	C	400	TXP	O4'-C4'-C5'-O5'
5	C	400	TXP	C1'-C2'-O2'-P2
4	B	700	GOL	C1-C2-C3-O3
5	C	400	TXP	C2N-C3N-C7N-N7N
5	C	400	TXP	C2N-C3N-C7N-O7N
5	C	400	TXP	C3'-C4'-C5'-O5'
3	B	600	NAD	PN-O3-PA-O5B
3	A	500	NAD	C5B-O5B-PA-O1A
3	A	500	NAD	C5D-O5D-PN-O3
5	C	400	TXP	C5'-O5'-PA-O3
5	C	400	TXP	C5'-O5'-PA-O1A

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	C	400	TXP	C5'-O5'-PA-O2A
3	B	600	NAD	C4D-C5D-O5D-PN
4	B	700	GOL	O2-C2-C3-O3
5	C	400	TXP	C3'-C2'-O2'-P2
3	B	600	NAD	PN-O3-PA-O1A
5	C	400	TXP	PA-O3-PN-O5'A
5	C	400	TXP	O4'A-C4'A-C5'A-O5'A


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	378/384 (98%)	1.84	157 (41%) 0 0	29, 63, 96, 108	0
1	B	364/384 (94%)	0.89	51 (14%) 6 7	26, 45, 74, 121	0
2	C	173/174 (99%)	1.82	69 (39%) 1 0	34, 66, 91, 99	0
All	All	915/942 (97%)	1.46	277 (30%) 1 1	26, 54, 93, 121	0

All (277) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	33	ALA	7.2
1	B	248	ALA	7.1
1	A	102	ALA	6.3
2	C	41	PHE	6.3
1	B	246	LYS	6.1
1	B	243	PHE	5.6
1	A	232	ALA	5.6
1	B	351	THR	5.5
1	A	207	THR	5.4
1	A	375	VAL	5.4
1	A	374	ILE	5.4
1	A	112	LEU	5.3
1	A	365	GLY	5.1
1	A	233	GLY	5.1
1	B	240	GLY	5.1
1	A	117	ILE	5.0
2	C	202	MET	5.0
1	A	58	ALA	5.0
1	A	368	VAL	4.9
1	A	366	THR	4.9
1	A	108	VAL	4.9
1	B	268	PRO	4.8
1	B	221	VAL	4.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	159	THR	4.7
1	B	247	GLN	4.7
1	A	64	ALA	4.7
1	A	74	VAL	4.6
1	A	111	ALA	4.5
1	A	119	ALA	4.5
2	C	162	PHE	4.5
1	A	235	TYR	4.5
1	A	27	VAL	4.5
1	A	120	TYR	4.4
1	A	100	LEU	4.4
1	A	351	THR	4.3
1	A	358	LEU	4.3
1	A	353	THR	4.3
2	C	199	VAL	4.3
1	B	242	GLU	4.3
1	A	378	ALA	4.3
2	C	39	ALA	4.3
1	A	126	PRO	4.3
1	A	226	MET	4.2
1	A	363	VAL	4.2
1	A	128	ILE	4.1
1	B	245	LYS	4.1
1	A	62	ALA	4.1
1	A	34	ILE	4.1
1	A	75	GLN	4.0
2	C	40	ALA	4.0
1	A	372	GLY	4.0
1	A	107	PRO	4.0
2	C	185	MET	4.0
1	A	65	LEU	4.0
1	B	381	GLY	3.9
1	A	225	ALA	3.9
1	A	229	ALA	3.9
1	A	362	THR	3.9
1	A	95	VAL	3.8
2	C	158	GLY	3.8
2	C	172	ALA	3.8
1	A	349	LYS	3.8
1	A	355	VAL	3.8
1	A	80	ALA	3.8
1	B	244	ARG	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	C	198	ILE	3.7
1	A	89	LEU	3.7
2	C	34	GLY	3.7
1	B	220	THR	3.6
2	C	32	LYS	3.6
1	A	109	VAL	3.6
1	A	281	VAL	3.6
2	C	187	LEU	3.6
2	C	188	PHE	3.5
1	A	376	HIS	3.5
1	A	347	VAL	3.5
1	A	29	LEU	3.5
1	B	358	LEU	3.5
2	C	42	ILE	3.5
1	A	124	LEU	3.5
1	A	354	LEU	3.5
1	A	1	MET	3.4
1	A	68	ALA	3.4
1	A	57	ILE	3.4
1	A	227	LYS	3.4
1	A	84	THR	3.4
2	C	49	VAL	3.4
1	A	83	GLY	3.4
1	A	78	MET	3.3
1	B	241	GLU	3.3
1	A	35	VAL	3.3
1	A	367	CYS	3.3
2	C	38	ASP	3.3
1	A	369	THR	3.3
1	A	377	PRO	3.3
1	B	12	GLY	3.3
2	C	180	PHE	3.3
2	C	46	ALA	3.2
1	B	205	ALA	3.2
1	A	116	LYS	3.2
1	A	51	THR	3.2
1	A	4	ALA	3.2
2	C	153	ASP	3.2
1	B	379	LEU	3.2
1	A	66	SER	3.2
1	B	349	LYS	3.2
2	C	44	LYS	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	16	VAL	3.2
1	A	350	ASP	3.2
1	A	223	ASP	3.1
2	C	186	MET	3.1
2	C	200	GLN	3.1
1	A	71	VAL	3.1
1	A	356	MET	3.1
2	C	177	GLU	3.1
1	A	221	VAL	3.1
2	C	154	VAL	3.1
1	B	1	MET	3.1
1	B	371	ASP	3.0
2	C	43	MET	3.0
1	A	361	GLU	3.0
2	C	163	ILE	3.0
2	C	103	ASN	3.0
2	C	201	ALA	3.0
1	A	87	VAL	3.0
1	B	355	VAL	3.0
1	A	96	LEU	3.0
1	B	251	VAL	3.0
2	C	31	VAL	3.0
2	C	72	VAL	3.0
1	A	99	HIS	3.0
1	A	60	THR	3.0
1	B	350	ASP	3.0
2	C	45	ASN	3.0
1	A	335	PHE	2.9
1	B	281	VAL	2.9
1	A	3	ILE	2.9
1	A	56	THR	2.9
1	A	371	ASP	2.9
1	A	251	VAL	2.9
1	A	352	LYS	2.9
1	A	50	LEU	2.9
2	C	179	PHE	2.9
2	C	35	SER	2.9
1	B	303	LEU	2.9
1	A	269	GLY	2.8
2	C	124	VAL	2.8
1	A	79	THR	2.8
1	B	267	ILE	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	105	ASN	2.8
1	A	33	VAL	2.8
1	B	310	VAL	2.8
2	C	183	ASN	2.8
1	A	54	GLY	2.8
2	C	47	SER	2.8
1	A	26	LEU	2.8
2	C	73	LEU	2.8
1	A	121	ALA	2.8
1	B	353	THR	2.8
1	A	94	ALA	2.8
2	C	36	ALA	2.8
2	C	189	GLY	2.7
1	B	250	ALA	2.7
2	C	171	TYR	2.7
1	A	276	ILE	2.7
1	B	249	GLU	2.7
1	A	88	ALA	2.7
2	C	140	THR	2.7
1	A	345	PRO	2.7
2	C	142	PRO	2.7
1	A	343	LEU	2.7
1	A	113	THR	2.7
1	A	90	ILE	2.7
1	A	373	ALA	2.6
1	A	370	ARG	2.6
1	A	348	ASP	2.6
1	B	9	ARG	2.6
1	A	18	ILE	2.6
2	C	144	SER	2.6
2	C	102	ALA	2.6
1	A	6	PRO	2.6
1	A	70	VAL	2.6
1	A	103	LEU	2.6
1	A	252	LEU	2.6
1	B	348	ASP	2.6
2	C	78	VAL	2.6
1	A	32	GLU	2.5
1	A	55	ALA	2.5
1	A	206	ALA	2.5
1	B	222	ASP	2.5
1	A	268	PRO	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	342	PHE	2.5
1	B	127	ARG	2.5
2	C	167	MET	2.5
1	A	114	LYS	2.5
1	B	129	SER	2.5
1	A	39	ALA	2.5
1	A	222	ASP	2.5
2	C	77	GLY	2.5
2	C	197	GLN	2.5
1	A	22	VAL	2.5
1	B	357	LYS	2.5
2	C	160	VAL	2.4
2	C	178	LEU	2.4
1	A	359	GLU	2.4
2	C	155	GLU	2.4
1	A	72	TRP	2.4
1	A	106	ARG	2.4
1	A	59	SER	2.4
1	B	138	SER	2.4
2	C	143	SER	2.4
1	A	73	LYS	2.4
2	C	75	LYS	2.4
1	A	77	PRO	2.3
1	A	61	ALA	2.3
1	A	364	SER	2.3
2	C	146	ILE	2.3
1	A	115	ARG	2.3
1	A	85	ASP	2.3
1	A	228	THR	2.3
1	A	23	VAL	2.3
1	A	91	LYS	2.3
1	A	17	ALA	2.3
1	B	356	MET	2.3
2	C	194	MET	2.3
2	C	175	GLU	2.3
1	A	30	GLY	2.3
1	B	362	THR	2.3
2	C	121	THR	2.3
1	A	304	SER	2.3
1	A	5	ILE	2.3
1	A	253	LYS	2.3
1	A	357	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	352	LYS	2.3
1	A	10	ARG	2.2
1	A	97	MET	2.2
1	A	104	THR	2.2
1	A	224	GLU	2.2
1	A	360	ASP	2.2
1	A	234	GLY	2.2
1	B	375	VAL	2.2
1	B	307	GLY	2.2
1	B	206	ALA	2.2
2	C	65	ALA	2.2
1	A	309	ILE	2.2
1	A	9	ARG	2.2
1	B	27	VAL	2.2
2	C	71	ASP	2.2
1	A	271	PRO	2.2
1	A	313	HIS	2.2
1	A	129	SER	2.2
1	A	275	LEU	2.1
1	A	69	ASP	2.1
2	C	107	ASP	2.1
1	A	92	GLU	2.1
1	A	209	GLU	2.1
1	B	279	GLU	2.1
2	C	145	PRO	2.1
2	C	184	THR	2.1
2	C	126	PHE	2.1
1	A	98	CYS	2.1
1	A	125	MET	2.1
2	C	161	LEU	2.1
1	A	37	GLN	2.1
1	B	269	GLY	2.1
2	C	30	SER	2.1
1	B	92	GLU	2.1
1	B	266	LEU	2.1
1	B	311	VAL	2.1
2	C	133	VAL	2.1
1	A	12	GLY	2.1
1	A	76	ARG	2.1
1	A	283	LYS	2.0
1	A	43	ALA	2.0
1	A	131	ALA	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	239	MET	2.0
1	B	343	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	B	700	6/6	0.80	0.17	48,49,49,50	0
3	NAD	B	600	36/44	0.81	0.16	76,78,81,81	0
3	NAD	A	500	44/44	0.89	0.12	48,53,65,65	0
5	TXP	C	400	48/48	0.89	0.12	42,48,51,53	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around NAD B 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAD A 500:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.