



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:54 AM UTC

PDB ID : 5OPK / pdb_00005opk
Title : Crystal structure of D52N/R367Q cN-II mutant bound to dATP and free phosphate
Authors : Hnizda, A.; Pachl, P.; Rezacova, P.
Deposited on : 2017-08-10
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1187 (1.74-1.74)
Clashscore	190562	1207 (1.74-1.74)
Ramachandran outliers	187476	1200 (1.74-1.74)
Sidechain outliers	187428	1200 (1.74-1.74)
RSRZ outliers	180081	1188 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	478	8% 96% ..

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4390 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytosolic purine 5'-nucleotidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	472	3988	2572	666	727	23	0	17	0

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	52	ASN	ASP	conflict	UNP P49902
A	367	GLN	ARG	conflict	UNP P49902
A	?	-	SER	deletion	UNP P49902
A	?	-	SER	deletion	UNP P49902
A	?	-	ASN	deletion	UNP P49902
A	?	-	GLU	deletion	UNP P49902
A	?	-	ARG	deletion	UNP P49902
A	?	-	PRO	deletion	UNP P49902
A	?	-	ASP	deletion	UNP P49902
A	?	-	ILE	deletion	UNP P49902

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Mg	0	0
			2	2		

- Molecule 5 is 2'-DEOXYADENOSINE 5'-TRIPHOSPHATE (CCD ID: DTP) (formula: C₁₀H₁₆N₅O₁₂P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	N	O	P	0	0
			30	10	5	12	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	297	Total	O	0	15
			311	311		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cytosolic purine 5'-nucleotidase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	91.58Å 127.10Å 130.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.99 – 1.74 48.99 – 1.74	Depositor EDS
% Data completeness (in resolution range)	98.8 (48.99-1.74) 99.2 (48.99-1.74)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.28 (at 1.74Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
R, R_{free}	0.178 , 0.203 0.189 , 0.213	Depositor DCC
R_{free} test set	3881 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	23.9	Xtriage
Anisotropy	0.237	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.012 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4390	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DTP, GOL, PO4, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	1/4097 (0.0%)	0.84	0/5528

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	277	ARG	CA-C	5.41	1.57	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3988	0	3927	9	0
2	A	54	0	72	1	0
3	A	5	0	0	0	0
4	A	2	0	0	0	0
5	A	30	0	12	2	0
6	A	311	0	0	3	0
All	All	4390	0	4011	12	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:456[B]:ARG:NH2	6:A:602:HOH:O	2.14	0.80
1:A:223[A]:GLU:H	1:A:223[A]:GLU:CD	1.91	0.78
5:A:514:DTP:O2G	5:A:514:DTP:H5'2	1.96	0.65
2:A:503:GOL:H11	6:A:823:HOH:O	1.96	0.64
5:A:514:DTP:O2G	5:A:514:DTP:C5'	2.48	0.61
1:A:145[A]:ASP:OD1	1:A:145[A]:ASP:N	2.30	0.60
1:A:456[B]:ARG:CZ	6:A:602:HOH:O	2.53	0.54
1:A:322:GLN:O	1:A:325:ILE:HG22	2.11	0.49
1:A:232[A]:LEU:HD12	1:A:236:LEU:HD13	1.98	0.45
1:A:43:MET:HE1	1:A:49:PHE:HE2	1.82	0.43
1:A:358:LEU:C	1:A:358:LEU:HD23	2.46	0.41
1:A:28:ARG:HD3	1:A:470:TYR:O	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	486/478 (102%)	479 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	436/424 (103%)	434 (100%)	2 (0%)	81 75

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	202	ARG
1	A	208	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	174	ASN
1	A	322	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 2 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	503	-	5,5,5	0.15	0	5,5,5	0.55	0
2	GOL	A	502	-	5,5,5	0.54	0	5,5,5	1.01	0
2	GOL	A	506	-	5,5,5	0.17	0	5,5,5	0.56	0
3	PO4	A	510	4	4,4,4	1.36	0	6,6,6	1.27	1 (16%)
2	GOL	A	507	-	5,5,5	0.49	0	5,5,5	1.22	0
2	GOL	A	501	-	5,5,5	0.60	0	5,5,5	0.97	0
2	GOL	A	508	-	5,5,5	0.39	0	5,5,5	0.85	0
2	GOL	A	505	-	5,5,5	0.28	0	5,5,5	0.50	0
5	DTP	A	514	4	31,32,32	1.44	7 (22%)	46,50,50	1.72	11 (23%)
2	GOL	A	504	-	5,5,5	0.31	0	5,5,5	0.40	0
2	GOL	A	509	-	5,5,5	0.57	0	5,5,5	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	503	-	-	2/4/4/4	-
2	GOL	A	502	-	-	2/4/4/4	-
2	GOL	A	506	-	-	0/4/4/4	-
2	GOL	A	507	-	-	1/4/4/4	-
2	GOL	A	501	-	-	2/4/4/4	-
2	GOL	A	508	-	-	3/4/4/4	-
2	GOL	A	505	-	-	2/4/4/4	-
5	DTP	A	514	4	-	7/22/34/34	0/3/3/3
2	GOL	A	504	-	-	2/4/4/4	-
2	GOL	A	509	-	-	1/4/4/4	-

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	514	DTP	C5-C4	4.31	1.46	1.39
5	A	514	DTP	PA-O3A	2.65	1.62	1.59
5	A	514	DTP	C8-N7	2.58	1.36	1.31
5	A	514	DTP	C4-N9	-2.58	1.32	1.37
5	A	514	DTP	PB-O3B	2.24	1.61	1.59
5	A	514	DTP	C5-N7	-2.22	1.35	1.39
5	A	514	DTP	C5-C6	2.06	1.46	1.41

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	514	DTP	C5-C4-N3	-4.19	120.95	126.72
5	A	514	DTP	N3-C2-N1	-3.88	122.71	128.58
5	A	514	DTP	N3-C4-N9	3.42	132.98	127.17
5	A	514	DTP	C2-N3-C4	3.18	119.60	111.83
5	A	514	DTP	O2A-PA-O3A	2.86	115.02	107.27
5	A	514	DTP	N6-C6-N1	2.71	124.42	118.38
5	A	514	DTP	C2-N1-C6	2.48	122.81	118.73
5	A	514	DTP	C4-N9-C8	2.35	108.21	105.74
5	A	514	DTP	O3A-PA-O1A	-2.34	103.65	110.70
5	A	514	DTP	C5-C6-N6	-2.23	117.76	123.29
5	A	514	DTP	O2B-PB-O3B	2.09	112.92	107.27
3	A	510	PO4	O4-P-O2	2.07	114.34	107.91

There are no chirality outliers.

All (22) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	504	GOL	O1-C1-C2-O2
2	A	504	GOL	O1-C1-C2-C3
5	A	514	DTP	C5'-O5'-PA-O2A
5	A	514	DTP	C5'-O5'-PA-O3A
5	A	514	DTP	O4'-C4'-C5'-O5'
5	A	514	DTP	C3'-C4'-C5'-O5'
2	A	501	GOL	O1-C1-C2-C3
2	A	502	GOL	O1-C1-C2-C3
2	A	503	GOL	O1-C1-C2-C3
2	A	508	GOL	C1-C2-C3-O3
2	A	501	GOL	O1-C1-C2-O2
2	A	509	GOL	O2-C2-C3-O3
2	A	503	GOL	O1-C1-C2-O2
5	A	514	DTP	PB-O3A-PA-O2A
5	A	514	DTP	C5'-O5'-PA-O1A
2	A	502	GOL	O1-C1-C2-O2
2	A	505	GOL	O2-C2-C3-O3
2	A	505	GOL	C1-C2-C3-O3
2	A	507	GOL	O1-C1-C2-C3
2	A	508	GOL	O1-C1-C2-C3
2	A	508	GOL	O2-C2-C3-O3
5	A	514	DTP	PB-O3A-PA-O1A

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	503	GOL	1	0
5	A	514	DTP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	472/478 (98%)	0.28	38 (8%) 18 23	6, 24, 57, 102	17 (3%)

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	473[A]	PHE	12.1
1	A	402	LEU	11.4
1	A	398	PHE	4.6
1	A	419	ILE	4.5
1	A	400	ALA	4.3
1	A	399	LEU	3.9
1	A	488	SER	3.7
1	A	397	ILE	3.6
1	A	15	MET	3.5
1	A	145[A]	ASP	3.5
1	A	3	THR	3.4
1	A	401	GLU	3.2
1	A	418	SER	3.1
1	A	417	SER	3.0
1	A	29	ARG	3.0
1	A	395	LEU	3.0
1	A	131	PRO	2.8
1	A	423	ILE	2.7
1	A	210	TYR	2.7
1	A	383	THR	2.7
1	A	394	SER	2.6
1	A	482	VAL	2.5
1	A	309	THR	2.5
1	A	308	LYS	2.3
1	A	129	ARG	2.3
1	A	481	HIS	2.2
1	A	388	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	390	GLU	2.2
1	A	26	LYS	2.2
1	A	27	TYR	2.2
1	A	480	ALA	2.1
1	A	44	GLU	2.1
1	A	135	GLU	2.1
1	A	396	ASP	2.1
1	A	128	ILE	2.1
1	A	188	GLY	2.0
1	A	387	SER	2.0
1	A	325	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GOL	A	504	6/6	0.74	0.21	42,49,52,54	0
2	GOL	A	505	6/6	0.76	0.25	34,44,47,50	0
2	GOL	A	509	6/6	0.79	0.20	39,46,54,58	0
2	GOL	A	506	6/6	0.81	0.23	50,54,55,59	0
2	GOL	A	508	6/6	0.85	0.18	34,45,50,52	0
2	GOL	A	501	6/6	0.86	0.17	24,34,38,45	0
2	GOL	A	507	6/6	0.87	0.17	29,36,41,43	0
2	GOL	A	502	6/6	0.89	0.15	30,36,38,47	0
2	GOL	A	503	6/6	0.89	0.16	40,44,46,54	0
5	DTP	A	514	30/30	0.91	0.12	18,25,35,37	30
4	MG	A	513	1/1	0.96	0.07	18,18,18,18	1
3	PO4	A	510	5/5	0.99	0.04	15,16,16,16	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MG	A	512	1/1	0.99	0.03	18,18,18,18	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.