



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 26, 2026 – 01:32 PM UTC

PDB ID : 1OQV / pdb_00001oqv
Title : Structure of TcpA, the Type IV pilin subunit from the toxin co-regulated pilus of *Vibrio cholerae* classical biotype
Authors : Craig, L.; Tainer, J.A.
Deposited on : 2003-03-11
Resolution : 1.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1553 (1.30-1.30)
Clashscore	190562	1595 (1.30-1.30)
Ramachandran outliers	187476	1551 (1.30-1.30)
Sidechain outliers	187428	1551 (1.30-1.30)
RSRZ outliers	180081	1549 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	192	8% 73% 15% •• 11%
1	B	192	10% 74% 13% • 11%
1	C	192	12% 78% 10% • 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	600	-	-	X	-

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4577 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called toxin-coregulated pilus subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	171	Total	C	N	O	S	0	15	0
			1270	795	213	255	7			
1	B	171	Total	C	N	O	S	0	12	0
			1259	787	211	254	7			
1	C	171	Total	C	N	O	S	0	14	0
			1265	790	211	257	7			

There are 63 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	8	MET	-	expression tag	UNP P23024
A	9	GLY	-	expression tag	UNP P23024
A	10	SER	-	expression tag	UNP P23024
A	11	SER	-	expression tag	UNP P23024
A	12	HIS	-	expression tag	UNP P23024
A	13	HIS	-	expression tag	UNP P23024
A	14	HIS	-	expression tag	UNP P23024
A	15	HIS	-	expression tag	UNP P23024
A	16	HIS	-	expression tag	UNP P23024
A	17	HIS	-	expression tag	UNP P23024
A	18	SER	-	expression tag	UNP P23024
A	19	SER	-	expression tag	UNP P23024
A	20	GLY	-	expression tag	UNP P23024
A	21	LEU	-	expression tag	UNP P23024
A	22	VAL	-	expression tag	UNP P23024
A	23	PRO	-	expression tag	UNP P23024
A	24	ARG	-	expression tag	UNP P23024
A	25	GLY	-	expression tag	UNP P23024
A	26	SER	-	expression tag	UNP P23024
A	27	HIS	-	expression tag	UNP P23024
A	28	MET	-	expression tag	UNP P23024
B	208	MET	-	expression tag	UNP P23024
B	209	GLY	-	expression tag	UNP P23024

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	210	SER	-	expression tag	UNP P23024
B	211	SER	-	expression tag	UNP P23024
B	212	HIS	-	expression tag	UNP P23024
B	213	HIS	-	expression tag	UNP P23024
B	214	HIS	-	expression tag	UNP P23024
B	215	HIS	-	expression tag	UNP P23024
B	216	HIS	-	expression tag	UNP P23024
B	217	HIS	-	expression tag	UNP P23024
B	218	SER	-	expression tag	UNP P23024
B	219	SER	-	expression tag	UNP P23024
B	220	GLY	-	expression tag	UNP P23024
B	221	LEU	-	expression tag	UNP P23024
B	222	VAL	-	expression tag	UNP P23024
B	223	PRO	-	expression tag	UNP P23024
B	224	ARG	-	expression tag	UNP P23024
B	225	GLY	-	expression tag	UNP P23024
B	226	SER	-	expression tag	UNP P23024
B	227	HIS	-	expression tag	UNP P23024
B	228	MET	-	expression tag	UNP P23024
C	408	MET	-	expression tag	UNP P23024
C	409	GLY	-	expression tag	UNP P23024
C	410	SER	-	expression tag	UNP P23024
C	411	SER	-	expression tag	UNP P23024
C	412	HIS	-	expression tag	UNP P23024
C	413	HIS	-	expression tag	UNP P23024
C	414	HIS	-	expression tag	UNP P23024
C	415	HIS	-	expression tag	UNP P23024
C	416	HIS	-	expression tag	UNP P23024
C	417	HIS	-	expression tag	UNP P23024
C	418	SER	-	expression tag	UNP P23024
C	419	SER	-	expression tag	UNP P23024
C	420	GLY	-	expression tag	UNP P23024
C	421	LEU	-	expression tag	UNP P23024
C	422	VAL	-	expression tag	UNP P23024
C	423	PRO	-	expression tag	UNP P23024
C	424	ARG	-	expression tag	UNP P23024
C	425	GLY	-	expression tag	UNP P23024
C	426	SER	-	expression tag	UNP P23024
C	427	HIS	-	expression tag	UNP P23024
C	428	MET	-	expression tag	UNP P23024

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		
2	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	265	Total	O	0	0
			265	265		
3	B	269	Total	O	0	0
			269	269		
3	C	231	Total	O	0	0
			231	231		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: toxin-coregulated pilus subunit

Chain A: 




- Molecule 1: toxin-coregulated pilus subunit

Chain B: 



- Molecule 1: toxin-coregulated pilus subunit

Chain C: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	157.32Å 157.32Å 35.72Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 1.30 30.00 – 1.30	Depositor EDS
% Data completeness (in resolution range)	5.0 (30.00-1.30) 98.3 (30.00-1.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 1.10Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.116 , 0.174 0.225 , 0.235	Depositor DCC
R_{free} test set	9432 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	11.7	Xtriage
Anisotropy	0.223	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 31.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.032 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4577	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 33.37 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.0868e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	1/1346 (0.1%)	1.49	22/1820 (1.2%)
1	B	0.69	1/1321 (0.1%)	1.44	16/1788 (0.9%)
1	C	0.61	0/1349	1.25	11/1825 (0.6%)
All	All	0.68	2/4016 (0.0%)	1.40	49/5433 (0.9%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	297	SER	N-CA	-5.23	1.39	1.46
1	A	30	SER	CB-OG	5.14	1.52	1.42

The worst 5 of 49 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	41	ASN	CA-CB-CG	-13.68	98.92	112.60
1	B	296	PHE	CA-C-N	12.78	140.21	122.19
1	B	296	PHE	C-N-CA	12.78	140.21	122.19
1	A	96	PHE	CA-C-N	12.19	140.12	122.09
1	A	96	PHE	C-N-CA	12.19	140.12	122.09

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1270	0	1273	24	0
1	B	1259	0	1256	27	0
1	C	1265	0	1256	16	0
2	A	6	0	8	4	0
2	B	12	0	16	3	0
3	A	265	0	0	9	0
3	B	269	0	0	15	1
3	C	231	0	0	11	0
All	All	4577	0	3809	67	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 67 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:285:LYS:HE3	3:B:973:HOH:O	1.57	1.05
1:A:49[B]:GLN:HG2	3:A:1370:HOH:O	1.72	0.89
1:C:572:LYS:HE3	3:C:1353:HOH:O	1.80	0.81
1:B:235:LYS:HE2	3:B:704:HOH:O	1.81	0.79
1:B:229:ASP:HA	3:B:1332:HOH:O	1.84	0.78

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:1213:HOH:O	3:B:1235:HOH:O[4_565]	1.92	0.28

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	183/192 (95%)	181 (99%)	2 (1%)	0	100	100
1	B	180/192 (94%)	178 (99%)	2 (1%)	0	100	100
1	C	184/192 (96%)	182 (99%)	2 (1%)	0	100	100
All	All	547/576 (95%)	541 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/146 (97%)	138 (97%)	4 (3%)	38	6
1	B	139/146 (95%)	138 (99%)	1 (1%)	76	50
1	C	143/146 (98%)	140 (98%)	3 (2%)	47	11
All	All	424/438 (97%)	416 (98%)	8 (2%)	55	14

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	530[B]	MET
1	C	530[A]	MET
1	B	229	ASP
1	A	80[B]	SER
1	C	527	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	232	ASN
1	B	238	GLN
1	C	449	GLN
1	A	105	ASN
1	A	41	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	601	-	5,5,5	0.72	0	5,5,5	0.71	0
2	GOL	A	600	-	5,5,5	0.90	0	5,5,5	0.82	0
2	GOL	B	602	-	5,5,5	0.66	0	5,5,5	0.86	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	601	-	-	2/4/4/4	-
2	GOL	A	600	-	-	4/4/4/4	-
2	GOL	B	602	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	600	GOL	O2-C2-C3-O3
2	A	600	GOL	O1-C1-C2-C3
2	A	600	GOL	C1-C2-C3-O3
2	B	601	GOL	O1-C1-C2-C3
2	B	601	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	600	GOL	4	0
2	B	602	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	171/192 (89%)	0.84	15 (8%)	15	17	6, 11, 26, 92	15 (8%)
1	B	171/192 (89%)	0.79	19 (11%)	10	11	6, 11, 28, 55	12 (7%)
1	C	171/192 (89%)	0.97	24 (14%)	6	7	7, 13, 30, 87	14 (8%)
All	All	513/576 (89%)	0.87	58 (11%)	10	11	6, 12, 29, 92	41 (7%)

The worst 5 of 58 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	229	ASP	6.6
1	A	29	ASP	6.3
1	C	569	PRO	6.0
1	B	330[A]	MET	5.5
1	A	30	SER	5.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	A	600	6/6	0.63	0.22	41,45,47,48	0
2	GOL	B	601	6/6	0.70	0.20	27,48,49,55	0
2	GOL	B	602	6/6	0.70	0.21	54,61,64,65	0

6.5 Other polymers [i](#)

There are no such residues in this entry.