



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 06:43 AM UTC

PDB ID : 2OQL / pdb_00002oql
Title : Structure of Phosphotriesterase mutant H254Q/H257F
Authors : Kim, J.; Tsai, P.; Raushel, F.M.; Almo, S.C.
Deposited on : 2007-01-31
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	 90% 9% .
1	B	331	 91% 7% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5906 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Parathion hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	330	Total	C	N	O	S	0	7	0
			2580	1629	459	485	7			
1	B	330	Total	C	N	O	S	0	8	0
			2594	1636	465	486	7			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	254	GLN	HIS	engineered mutation	UNP P0A434
A	257	PHE	HIS	engineered mutation	UNP P0A434
B	254	GLN	HIS	engineered mutation	UNP P0A434
B	257	PHE	HIS	engineered mutation	UNP P0A434

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (CCD ID: BTB) (formula: C₈H₁₉NO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	348	Total	O	0	0
			348	348		
5	B	334	Total	O	0	0
			334	334		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Parathion hydrolase



- Molecule 1: Parathion hydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	43.53Å 45.37Å 79.11Å 104.75° 93.13° 97.97°	Depositor
Resolution (Å)	50.00 – 1.80 50.00 – 1.80	Depositor EDS
% Data completeness (in resolution range)	91.8 (50.00-1.80) 91.8 (50.00-1.80)	Depositor EDS
R_{merge}	0.01	Depositor
R_{sym}	0.01	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.146 , 0.189 0.146 , 0.189	Depositor DCC
R_{free} test set	2483 reflections (4.64%)	wwPDB-VP
Wilson B-factor (Å ²)	15.1	Xtriage
Anisotropy	0.029	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 41.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5906	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, BTB, ZN, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/2614	0.86	2/3549 (0.1%)
1	B	0.79	1/2628 (0.0%)	0.87	1/3564 (0.0%)
All	All	0.77	1/5242 (0.0%)	0.86	3/7113 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	333	ILE	CA-CB	5.19	1.57	1.54

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	361	THR	CB-CA-C	-7.45	94.18	111.09
1	B	363	ARG	N-CA-CB	6.67	121.76	110.49
1	A	274	ILE	CB-CA-C	5.70	117.19	110.93

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	362	LEU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2580	0	2598	20	0
1	B	2594	0	2608	16	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	14	0	19	1	0
3	B	14	0	19	0	0
4	A	6	0	8	0	0
4	B	12	0	16	0	0
5	A	348	0	0	3	0
5	B	334	0	0	2	0
All	All	5906	0	5268	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:PHE:CZ	1:A:173[A]:THR:HG23	2.05	0.92
1:B:132:PHE:CZ	1:B:173:THR:HG23	2.13	0.84
1:A:263:GLU:HG3	1:A:274:ILE:HD11	1.70	0.72
1:A:361:THR:HG21	5:A:929:HOH:O	1.92	0.69
1:B:57:HIS:HB2	1:B:303:LEU:HB3	1.79	0.65
1:A:361:THR:CG2	5:A:929:HOH:O	2.46	0.64
1:B:41[B]:ARG:NE	1:B:118[B]:ARG:HG2	2.15	0.60
1:A:131:TRP:CG	1:A:132:PHE:H	2.22	0.56
1:B:337[A]:ARG:NH1	1:B:343:GLN:H	2.04	0.55
1:A:57:HIS:HB2	1:A:303:LEU:HB3	1.89	0.55
1:B:131:TRP:CG	1:B:132:PHE:H	2.25	0.54
1:B:35:ASP:N	5:B:1026:HOH:O	2.41	0.53
1:A:173[A]:THR:HG22	5:A:1026:HOH:O	2.06	0.53
1:B:57:HIS:O	1:B:303:LEU:HA	2.08	0.53
1:A:57:HIS:O	1:A:303:LEU:HA	2.09	0.53
1:A:232:ASP:HA	1:A:283:LEU:HD23	1.92	0.52
1:A:46:ILE:HG23	1:A:355:ALA:HB1	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:333:ILE:HB	1:A:334:PRO:HD3	1.92	0.50
1:A:319:ARG:HD3	1:A:319:ARG:C	2.38	0.49
1:B:333:ILE:HB	1:B:334:PRO:HD3	1.96	0.48
1:A:302:TRP:CH2	1:A:321:ASN:HB3	2.48	0.48
1:A:106:ILE:HG22	1:A:106:ILE:O	2.14	0.48
1:A:314:MET:HE3	1:A:314:MET:HA	1.96	0.47
1:B:41[B]:ARG:CZ	1:B:118[B]:ARG:HG2	2.45	0.47
1:A:364:ALA:O	1:A:365:SER:HB3	2.16	0.46
1:B:106:ILE:HG22	1:B:106:ILE:O	2.14	0.46
3:A:701:BTB:H31	3:A:701:BTB:H71	1.83	0.46
1:A:202:THR:HB	1:A:208:ASP:HB2	1.98	0.45
1:A:194:THR:OG1	1:A:196[B]:VAL:HG22	2.17	0.44
1:B:314:MET:HE3	1:B:314:MET:HA	1.99	0.44
1:A:132:PHE:CE2	1:A:173[A]:THR:HG23	2.49	0.44
1:B:302:TRP:CH2	1:B:321:ASN:HB3	2.52	0.43
1:A:85:ARG:O	1:A:89:ARG:HG3	2.19	0.42
1:B:225:ARG:HD3	1:B:363:ARG:H	1.85	0.42
1:B:249:LEU:HD13	1:B:295[B]:GLN:HA	2.02	0.41
1:B:135:PRO:O	1:B:139:ARG:HG3	2.21	0.41
1:B:264:ASP:CG	5:B:978:HOH:O	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	335/331 (101%)	321 (96%)	14 (4%)	0	100	100
1	B	335/331 (101%)	324 (97%)	10 (3%)	1 (0%)	36	25
All	All	670/662 (101%)	645 (96%)	24 (4%)	1 (0%)	43	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	363	ARG

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	272/265 (103%)	264 (97%)	8 (3%)	37	25
1	B	272/265 (103%)	264 (97%)	8 (3%)	37	25
All	All	544/530 (103%)	528 (97%)	16 (3%)	38	25

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	184	LEU
1	A	206	GLN
1	A	237	LEU
1	A	262	LEU
1	A	274	ILE
1	A	299	SER
1	A	303	LEU
1	A	361	THR
1	B	164	ARG
1	B	173	THR
1	B	184	LEU
1	B	264	ASP
1	B	299	SER
1	B	303	LEU
1	B	337[A]	ARG
1	B	337[B]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	212	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	KCX	B	169	1,2	10,11,12	0.98	1 (10%)	6,12,14	1.86	1 (16%)
1	KCX	A	169	1,2	10,11,12	1.05	1 (10%)	6,12,14	1.62	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	B	169	1,2	-	1/9/10/12	-
1	KCX	A	169	1,2	-	0/9/10/12	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	169	KCX	OQ1-CX	2.77	1.26	1.21
1	B	169	KCX	OQ1-CX	2.41	1.26	1.21

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	169	KCX	OQ1-CX-NZ	-4.33	118.35	124.92
1	A	169	KCX	OQ1-CX-NZ	-3.26	119.97	124.92
1	A	169	KCX	CE-NZ-CX	2.06	125.47	121.98

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	169	KCX	C-CA-CB-CG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	BTB	B	701	-	13,13,13	0.68	0	7,16,16	0.85	0
4	GOL	A	802	2	5,5,5	0.36	0	5,5,5	0.43	0
4	GOL	B	801	-	5,5,5	0.44	0	5,5,5	0.58	0
3	BTB	A	701	-	13,13,13	1.01	1 (7%)	7,16,16	0.73	0
4	GOL	B	803	-	5,5,5	0.41	0	5,5,5	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BTB	B	701	-	-	1/21/21/21	-
4	GOL	A	802	2	-	1/4/4/4	-
4	GOL	B	801	-	-	2/4/4/4	-
3	BTB	A	701	-	-	0/21/21/21	-
4	GOL	B	803	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	701	BTB	C5-N	2.22	1.51	1.48

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	801	GOL	O1-C1-C2-C3
4	B	803	GOL	C1-C2-C3-O3
4	B	803	GOL	O2-C2-C3-O3
4	B	801	GOL	O1-C1-C2-O2
3	B	701	BTB	O1-C1-C2-C3
4	A	802	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	701	BTB	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	329/331 (99%)	-0.68	0 100 100	4, 12, 19, 29	7 (2%)
1	B	329/331 (99%)	-0.65	0 100 100	5, 12, 20, 30	8 (2%)
All	All	658/662 (99%)	-0.67	0 100 100	4, 12, 20, 30	15 (2%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	KCX	A	169	12/13	0.96	0.04	8,9,12,12	0
1	KCX	B	169	12/13	0.96	0.05	10,11,13,13	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	B	803	6/6	0.76	0.24	54,54,54,54	0
4	GOL	A	802	6/6	0.87	0.15	32,33,33,34	0
4	GOL	B	801	6/6	0.88	0.11	28,29,29,30	0
3	BTB	B	701	14/14	0.95	0.06	12,15,16,17	0
3	BTB	A	701	14/14	0.96	0.06	11,13,16,16	0
2	ZN	A	402	1/1	1.00	0.01	17,17,17,17	1
2	ZN	B	401	1/1	1.00	0.02	17,17,17,17	0
2	ZN	B	402	1/1	1.00	0.01	19,19,19,19	1
2	ZN	A	401	1/1	1.00	0.01	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.