



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 02:14 AM UTC

PDB ID : 3OQ8 / pdb_00003oq8
Title : Crystal structure of isocitrate lyase from Brucella melitensis, bound to the product mimic malonate
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2010-09-02
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1898 (2.26-2.26)
Clashscore	190562	2005 (2.26-2.26)
Ramachandran outliers	187476	1965 (2.26-2.26)
Sidechain outliers	187428	1966 (2.26-2.26)
RSRZ outliers	180081	1898 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	433	 90% 8% .
1	B	433	 92% 5% .
1	C	433	 92% 6% .
1	D	433	 92% 6% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 14231 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Isocitrate lyase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	424	Total	C	N	O	S	0	4	0
			3251	2054	563	619	15			
1	B	422	Total	C	N	O	S	0	1	0
			3232	2040	558	619	15			
1	C	423	Total	C	N	O	S	0	3	0
			3246	2051	564	616	15			
1	D	423	Total	C	N	O	S	0	3	0
			3244	2050	563	616	15			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP Q2YQA0
A	-2	PRO	-	expression tag	UNP Q2YQA0
A	-1	GLY	-	expression tag	UNP Q2YQA0
A	0	SER	-	expression tag	UNP Q2YQA0
B	-3	GLY	-	expression tag	UNP Q2YQA0
B	-2	PRO	-	expression tag	UNP Q2YQA0
B	-1	GLY	-	expression tag	UNP Q2YQA0
B	0	SER	-	expression tag	UNP Q2YQA0
C	-3	GLY	-	expression tag	UNP Q2YQA0
C	-2	PRO	-	expression tag	UNP Q2YQA0
C	-1	GLY	-	expression tag	UNP Q2YQA0
C	0	SER	-	expression tag	UNP Q2YQA0
D	-3	GLY	-	expression tag	UNP Q2YQA0
D	-2	PRO	-	expression tag	UNP Q2YQA0
D	-1	GLY	-	expression tag	UNP Q2YQA0
D	0	SER	-	expression tag	UNP Q2YQA0

- Molecule 2 is MALONIC ACID (CCD ID: MLA) (formula: C₃H₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			7	3	4		
2	B	1	Total	C	O	0	0
			7	3	4		
2	C	1	Total	C	O	0	0
			7	3	4		
2	D	1	Total	C	O	0	0
			7	3	4		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0

- Molecule 4 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Na 2 2	0	0
4	B	2	Total Na 2 2	0	0
4	C	3	Total Na 3 3	0	0
4	D	2	Total Na 2 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	292	Total O 292 292	0	0
5	B	286	Total O 286 286	0	0
5	C	313	Total O 313 313	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	286	Total 286	O 286	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Isocitrate lyase




• Molecule 1: Isocitrate lyase



• Molecule 1: Isocitrate lyase



• Molecule 1: Isocitrate lyase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	77.50Å 136.03Å 181.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.20 – 2.25 39.20 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.0 (39.20-2.25) 98.9 (39.20-2.25)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.04 (at 2.24Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.149 , 0.199 (Not available) , 0.164	Depositor DCC
R_{free} test set	4530 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	17.7	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 33.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	14231	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MLA, NA, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.97	2/3333 (0.1%)	0.92	1/4515 (0.0%)
1	B	0.96	0/3304	0.93	0/4474
1	C	1.00	1/3324 (0.0%)	0.93	2/4500 (0.0%)
1	D	0.95	0/3322	0.93	3/4497 (0.1%)
All	All	0.97	3/13283 (0.0%)	0.93	6/17986 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	272	ALA	CA-CB	5.87	1.60	1.53
1	A	326	ALA	CA-CB	5.07	1.61	1.53
1	C	37	ILE	CA-CB	-5.03	1.48	1.54

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	399	ASP	N-CA-C	-5.36	105.33	111.07
1	D	8	ILE	CA-C-N	5.28	125.59	119.47
1	D	8	ILE	C-N-CA	5.28	125.59	119.47
1	D	311	SER	N-CA-C	5.12	116.11	109.65
1	C	215	THR	CA-C-N	5.04	126.14	119.84
1	C	215	THR	C-N-CA	5.04	126.14	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3251	0	3145	19	0
1	B	3232	0	3125	17	0
1	C	3246	0	3153	14	0
1	D	3244	0	3147	14	0
2	A	7	0	2	0	0
2	B	7	0	2	0	0
2	C	7	0	2	0	0
2	D	7	0	2	0	0
3	A	12	0	18	0	0
3	B	8	0	12	2	0
3	C	16	0	24	0	0
3	D	8	0	12	0	0
4	A	2	0	0	0	0
4	B	2	0	0	0	0
4	C	3	0	0	0	0
4	D	2	0	0	0	0
5	A	292	0	0	3	0
5	B	286	0	0	1	0
5	C	313	0	0	1	0
5	D	286	0	0	1	0
All	All	14231	0	12644	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:349:GLN:HG2	5:C:472:HOH:O	1.84	0.77
1:D:269:ILE:HD12	1:D:300:ALA:HB3	1.72	0.71
1:A:28:VAL:O	1:A:32[B]:ARG:HG2	1.95	0.66
1:B:53:ILE:HD12	1:B:305:LEU:HD13	1.78	0.64
1:A:280:MET:HE2	1:A:282:THR:HG22	1.82	0.62
1:D:269:ILE:HD11	1:D:297:VAL:HA	1.83	0.60
1:C:115:LYS:O	1:C:119:ARG:HG3	2.02	0.60

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:32:ARG:HD3	5:D:438:HOH:O	2.05	0.57
1:C:350:LEU:C	1:C:350:LEU:HD23	2.35	0.52
1:D:363:ASP:OD2	1:D:364:ARG:HG2	2.11	0.50
1:D:70:ALA:CB	1:D:117:ILE:HG23	2.43	0.49
1:B:86:TRP:CD1	1:B:101:ASP:HB2	2.48	0.48
1:C:28:VAL:O	1:C:32:ARG:HB2	2.12	0.48
1:D:328:PHE:CE1	1:D:332:LEU:HD11	2.50	0.47
1:A:28:VAL:O	1:A:32[B]:ARG:CG	2.63	0.47
1:B:79:LYS:O	1:B:143:PRO:HD2	2.15	0.46
1:A:30[A]:ARG:NH2	5:A:576:HOH:O	2.47	0.45
1:C:316:TRP:CZ2	1:D:366:MET:HE2	2.52	0.45
1:C:38:LYS:HA	1:C:38:LYS:HD2	1.88	0.45
1:B:384:TYR:CZ	1:B:386:ALA:HB3	2.52	0.45
1:D:185:CYS:HB3	1:D:188:LEU:HG	1.98	0.45
1:B:174:HIS:HB3	1:B:220:VAL:HB	1.98	0.45
1:A:86:TRP:CD1	1:A:101:ASP:HB2	2.51	0.45
1:C:83:LEU:C	1:C:83:LEU:HD23	2.42	0.44
1:B:350:LEU:C	1:B:350:LEU:HD23	2.42	0.43
1:A:366:MET:HE2	1:B:316:TRP:CZ2	2.53	0.43
1:C:86:TRP:HB3	1:C:344:LEU:HD21	2.01	0.43
1:C:147:ASP:CG	1:C:174:HIS:CE1	2.96	0.43
1:A:350:LEU:C	1:A:350:LEU:HD23	2.43	0.43
1:A:384:TYR:CZ	1:A:386:ALA:HB3	2.54	0.43
1:B:115:LYS:O	1:B:119:ARG:HG2	2.19	0.42
1:D:65:LEU:C	1:D:65:LEU:HD12	2.43	0.42
1:D:86:TRP:CD1	1:D:101:ASP:HB2	2.54	0.42
1:A:174:HIS:HB3	1:A:220:VAL:HB	2.00	0.42
1:D:174:HIS:HB3	1:D:220:VAL:HB	2.00	0.42
1:B:86:TRP:HB3	1:B:344:LEU:HD21	2.01	0.42
1:B:83:LEU:HD23	1:B:83:LEU:C	2.44	0.42
1:A:240:GLN:N	1:A:241:PRO:CD	2.83	0.42
1:C:240:GLN:N	1:C:241:PRO:CD	2.83	0.42
1:A:53:ILE:HD12	1:A:305:LEU:HD13	2.01	0.42
1:A:32[B]:ARG:HA	1:A:212:VAL:HG13	2.01	0.42
1:B:60:ASN:HA	1:B:340:GLN:O	2.19	0.42
1:B:65:LEU:HD12	1:B:65:LEU:C	2.45	0.42
1:C:366:MET:HE2	1:D:316:TRP:CZ2	2.55	0.41
1:D:115:LYS:O	1:D:119:ARG:CG	2.68	0.41
1:C:185:CYS:HB3	1:C:188:LEU:HG	2.01	0.41
1:A:27:ASP:OD1	1:A:30[B]:ARG:NH2	2.47	0.41
1:A:389:HIS:ND1	1:B:348:HIS:NE2	2.63	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:60:ASN:HA	1:A:340:GLN:O	2.21	0.41
1:A:65:LEU:HD13	5:A:605:HOH:O	2.21	0.41
1:A:207:ARG:NE	5:A:526:HOH:O	2.42	0.41
1:C:366:MET:HG2	1:D:329:GLN:HG3	2.03	0.41
1:A:420:GLU:O	1:A:424:PHE:HB2	2.20	0.41
1:B:115:LYS:NZ	5:B:533:HOH:O	2.47	0.41
1:B:127:ILE:CD1	3:B:475:EDO:H12	2.52	0.40
1:B:127:ILE:HD12	3:B:475:EDO:H12	2.02	0.40
1:C:317:LYS:NZ	1:C:322:ASP:OD1	2.48	0.40
1:A:389:HIS:HD1	1:B:348:HIS:CE1	2.39	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	426/433 (98%)	415 (97%)	11 (3%)	0	100	100
1	B	421/433 (97%)	414 (98%)	7 (2%)	0	100	100
1	C	424/433 (98%)	414 (98%)	10 (2%)	0	100	100
1	D	424/433 (98%)	414 (98%)	10 (2%)	0	100	100
All	All	1695/1732 (98%)	1657 (98%)	38 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	318/327 (97%)	311 (98%)	7 (2%)	45	56
1	B	317/327 (97%)	315 (99%)	2 (1%)	78	84
1	C	318/327 (97%)	316 (99%)	2 (1%)	78	84
1	D	317/327 (97%)	314 (99%)	3 (1%)	70	79
All	All	1270/1308 (97%)	1256 (99%)	14 (1%)	68	75

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	SER
1	A	32[A]	ARG
1	A	32[B]	ARG
1	A	226	GLU
1	A	364	ARG
1	A	387	THR
1	A	388	LYS
1	B	212	VAL
1	B	387	THR
1	C	212	VAL
1	C	387	THR
1	D	32	ARG
1	D	212	VAL
1	D	387	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	319	ASN
1	A	349	GLN
1	A	365	GLN
1	B	240	GLN
1	B	374	GLN
1	C	374	GLN
1	D	199	HIS
1	D	202	ASN
1	D	256	GLN
1	D	319	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 24 ligands modelled in this entry, 9 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MLA	A	440	-	6,6,6	1.20	0	7,7,7	1.06	0
3	EDO	A	465	-	3,3,3	0.50	0	2,2,2	0.05	0
3	EDO	C	470	-	3,3,3	0.97	0	2,2,2	0.48	0
2	MLA	D	440	-	6,6,6	1.14	1 (16%)	7,7,7	1.28	0
3	EDO	A	450	-	3,3,3	0.63	0	2,2,2	0.25	0
2	MLA	C	440	-	6,6,6	1.06	0	7,7,7	1.08	0
3	EDO	D	475	-	3,3,3	0.65	0	2,2,2	0.16	0
3	EDO	A	475	-	3,3,3	0.65	0	2,2,2	0.22	0
3	EDO	B	475	-	3,3,3	0.75	0	2,2,2	0.22	0
3	EDO	B	465	-	3,3,3	0.66	0	2,2,2	0.13	0
2	MLA	B	440	-	6,6,6	1.21	0	7,7,7	1.34	1 (14%)
3	EDO	C	465	-	3,3,3	0.80	0	2,2,2	0.34	0
3	EDO	D	465	-	3,3,3	0.77	0	2,2,2	0.22	0
3	EDO	C	475	-	3,3,3	0.73	0	2,2,2	0.31	0
3	EDO	C	450	-	3,3,3	0.66	0	2,2,2	0.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MLA	A	440	-	-	0/4/4/4	-
3	EDO	A	465	-	-	1/1/1/1	-
3	EDO	C	470	-	-	1/1/1/1	-
2	MLA	D	440	-	-	2/4/4/4	-
3	EDO	A	450	-	-	0/1/1/1	-
2	MLA	C	440	-	-	1/4/4/4	-
3	EDO	D	475	-	-	1/1/1/1	-
3	EDO	A	475	-	-	1/1/1/1	-
3	EDO	B	475	-	-	1/1/1/1	-
3	EDO	B	465	-	-	0/1/1/1	-
2	MLA	B	440	-	-	2/4/4/4	-
3	EDO	C	465	-	-	0/1/1/1	-
3	EDO	D	465	-	-	0/1/1/1	-
3	EDO	C	475	-	-	1/1/1/1	-
3	EDO	C	450	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	440	MLA	O3B-C3	-2.23	1.23	1.30

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	440	MLA	O1B-C1-C2	-2.19	115.87	122.11

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	475	EDO	O1-C1-C2-O2
3	B	475	EDO	O1-C1-C2-O2
3	C	470	EDO	O1-C1-C2-O2
3	A	475	EDO	O1-C1-C2-O2
3	C	475	EDO	O1-C1-C2-O2
2	D	440	MLA	C1-C2-C3-O3A
2	B	440	MLA	O1B-C1-C2-C3
2	B	440	MLA	O1A-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	C	440	MLA	C1-C2-C3-O3A
2	D	440	MLA	C1-C2-C3-O3B
3	A	465	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	475	EDO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	424/433 (97%)	-0.79	2 (0%) 87 88	7, 15, 25, 42	4 (0%)
1	B	422/433 (97%)	-0.74	0 100 100	8, 16, 28, 36	1 (0%)
1	C	423/433 (97%)	-0.80	1 (0%) 91 92	7, 14, 24, 33	3 (0%)
1	D	423/433 (97%)	-0.74	1 (0%) 91 92	8, 15, 26, 33	3 (0%)
All	All	1692/1732 (97%)	-0.76	4 (0%) 91 92	7, 15, 26, 42	11 (0%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	3	ASP	2.5
1	A	426	PRO	2.3
1	A	3	ASP	2.1
1	C	3	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	EDO	C	470	4/4	0.82	0.14	25,27,30,30	0
3	EDO	C	475	4/4	0.91	0.14	23,24,27,28	0
3	EDO	B	465	4/4	0.92	0.09	21,26,26,28	0
3	EDO	C	465	4/4	0.92	0.10	18,22,24,24	0
3	EDO	D	465	4/4	0.92	0.11	20,25,26,28	0
4	NA	C	430	1/1	0.92	0.05	29,29,29,29	0
3	EDO	B	475	4/4	0.94	0.09	19,25,26,27	0
3	EDO	A	475	4/4	0.94	0.08	22,24,25,26	0
4	NA	C	460	1/1	0.95	0.05	18,18,18,18	0
3	EDO	D	475	4/4	0.95	0.09	22,25,26,26	0
2	MLA	B	440	7/7	0.96	0.06	14,17,19,20	0
3	EDO	A	465	4/4	0.96	0.06	19,24,24,24	0
3	EDO	A	450	4/4	0.97	0.06	20,21,23,26	0
3	EDO	C	450	4/4	0.97	0.05	19,19,19,20	0
4	NA	A	501	1/1	0.98	0.02	14,14,14,14	0
4	NA	B	460	1/1	0.98	0.04	17,17,17,17	0
2	MLA	D	440	7/7	0.98	0.04	17,17,18,19	0
4	NA	C	501	1/1	0.98	0.03	12,12,12,12	0
2	MLA	A	440	7/7	0.98	0.05	15,17,19,19	0
4	NA	D	501	1/1	0.98	0.03	15,15,15,15	0
4	NA	A	460	1/1	0.99	0.04	15,15,15,15	0
4	NA	B	501	1/1	0.99	0.03	13,13,13,13	0
4	NA	D	460	1/1	0.99	0.03	20,20,20,20	0
2	MLA	C	440	7/7	0.99	0.04	14,16,17,19	0

6.5 Other polymers [i](#)

There are no such residues in this entry.