



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:51 AM UTC

PDB ID : 3OQJ / pdb_00003oqj
Title : Crystal structure of B. licheniformis CDPS yvmC-BLIC in complex with CAPSO
Authors : Bonnefond, L.; Arai, T.; Suzuki, T.; Ishitani, R.; Nureki, O.
Deposited on : 2010-09-03
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	257	
1	B	257	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	3CX	A	258	X	-	-	-
2	3CX	B	258	X	-	-	-

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3628 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Putative uncharacterized protein yvmC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	213	Total	C	N	O	S	0	0	0
			1692	1073	305	307	7			
1	B	220	Total	C	N	O	S	0	0	0
			1733	1097	312	316	8			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	250	LEU	-	expression tag	UNP Q65EX3
A	251	GLU	-	expression tag	UNP Q65EX3
A	252	HIS	-	expression tag	UNP Q65EX3
A	253	HIS	-	expression tag	UNP Q65EX3
A	254	HIS	-	expression tag	UNP Q65EX3
A	255	HIS	-	expression tag	UNP Q65EX3
A	256	HIS	-	expression tag	UNP Q65EX3
A	257	HIS	-	expression tag	UNP Q65EX3
B	250	LEU	-	expression tag	UNP Q65EX3
B	251	GLU	-	expression tag	UNP Q65EX3
B	252	HIS	-	expression tag	UNP Q65EX3
B	253	HIS	-	expression tag	UNP Q65EX3
B	254	HIS	-	expression tag	UNP Q65EX3
B	255	HIS	-	expression tag	UNP Q65EX3
B	256	HIS	-	expression tag	UNP Q65EX3
B	257	HIS	-	expression tag	UNP Q65EX3

- Molecule 2 is (2S)-3-(cyclohexylamino)-2-hydroxypropane-1-sulfonic acid (CCD ID: 3CX) (formula: C₉H₁₉NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			15	9	1	4	1		
2	B	1	Total	C	N	O	S	0	0
			15	9	1	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	71	Total	O	0	0
			71	71		
3	B	102	Total	O	0	0
			102	102		

- Molecule 1: Putative uncharacterized protein yvmC



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	150.86Å 54.71Å 101.52Å 90.00° 130.44° 90.00°	Depositor
Resolution (Å)	38.63 – 2.40 38.63 – 2.40	Depositor EDS
% Data completeness (in resolution range)	96.5 (38.63-2.40) 95.8 (38.63-2.40)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.40 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.6.4_486	Depositor
R, R_{free}	0.178 , 0.225 0.173 , 0.218	Depositor DCC
R_{free} test set	1998 reflections (8.02%)	wwPDB-VP
Wilson B-factor (Å ²)	29.6	Xtriage
Anisotropy	0.178	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 40.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.024 for -h-2*1,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3628	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 3CX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/1727	0.80	0/2334
1	B	0.52	0/1768	0.79	2/2389 (0.1%)
All	All	0.52	0/3495	0.79	2/4723 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	110	ASN	CA-C-N	6.77	126.45	119.82
1	B	110	ASN	C-N-CA	6.77	126.45	119.82

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1692	0	1655	29	0
1	B	1733	0	1681	36	0
2	A	15	0	17	2	0
2	B	15	0	17	0	0
3	A	71	0	0	0	0
3	B	102	0	0	3	0
All	All	3628	0	3370	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:116:THR:HG22	1:B:118:SER:H	1.23	1.04
1:A:39:PHE:CE2	1:A:92:GLU:HG3	2.07	0.89
1:A:17:THR:HG23	1:A:22:CYS:HB2	1.55	0.87
1:A:15:THR:HB	1:A:230:LEU:HD11	1.67	0.76
1:A:17:THR:CG2	1:A:22:CYS:HB2	2.16	0.75
1:B:208:TRP:CE3	1:B:211:GLY:HA3	2.24	0.73
1:B:123:GLN:NE2	1:B:125:ALA:HB3	2.09	0.67
1:B:18:LEU:HD23	1:B:223:MET:HE1	1.76	0.67
1:B:140:GLN:HB3	1:B:142:HIS:CE1	2.32	0.65
1:A:39:PHE:CZ	1:A:92:GLU:HG3	2.33	0.62
1:B:68:GLY:HA2	1:B:116:THR:HG23	1.82	0.62
1:B:15:THR:HB	1:B:230:LEU:HD11	1.83	0.61
1:B:123:GLN:HE21	1:B:126:TYR:H	1.51	0.58
1:A:39:PHE:HE2	1:A:88:LYS:HE3	1.69	0.57
1:A:208:TRP:HZ3	1:A:223:MET:HE1	1.69	0.57
1:A:171:ALA:O	1:A:175:GLU:OE1	2.24	0.56
1:B:120:PHE:HB3	1:B:123:GLN:HG2	1.87	0.56
1:B:233:SER:HA	3:B:352:HOH:O	2.06	0.55
1:B:23:ASN:HD21	1:B:27:LYS:HE3	1.70	0.55
1:A:184:GLU:OE1	2:A:258:3CX:HAJ	2.07	0.54
1:A:208:TRP:CE3	1:A:211:GLY:HA3	2.43	0.54
1:A:208:TRP:CZ3	1:A:223:MET:HE1	2.42	0.53
1:B:42:ARG:HG3	1:B:42:ARG:HH11	1.73	0.53
1:B:23:ASN:ND2	1:B:27:LYS:HE3	2.23	0.52
1:B:116:THR:HG21	3:B:281:HOH:O	2.08	0.52
1:B:17:THR:OG1	1:B:22:CYS:HB3	2.09	0.51
1:B:144:ARG:NE	3:B:295:HOH:O	2.42	0.51
1:B:123:GLN:HE21	1:B:126:TYR:N	2.08	0.51
1:B:123:GLN:HE22	1:B:194:ILE:HA	1.76	0.51
1:B:42:ARG:HD3	1:B:47:TYR:CZ	2.46	0.51
1:B:170:SER:OG	1:B:173:MET:HG3	2.11	0.51
1:A:137:PHE:CE2	1:A:144:ARG:HG3	2.46	0.50
1:A:212:GLU:O	1:A:215:SER:HB2	2.11	0.50
1:B:18:LEU:HB3	1:B:22:CYS:SG	2.52	0.50
1:A:110:ASN:OD1	1:A:112:GLU:OE1	2.29	0.50
1:B:127:ARG:CZ	1:B:127:ARG:HB3	2.43	0.48
1:B:116:THR:HG22	1:B:118:SER:N	2.08	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:208:TRP:CD2	1:B:211:GLY:HA3	2.48	0.48
1:B:42:ARG:HG3	1:B:42:ARG:NH1	2.28	0.48
1:B:68:GLY:HA2	1:B:116:THR:CG2	2.44	0.48
1:B:130:ARG:NH2	1:B:179:GLU:OE2	2.47	0.47
1:A:39:PHE:CE2	1:A:88:LYS:HE3	2.50	0.46
1:B:95:ARG:O	1:B:98:ARG:HG2	2.16	0.46
1:A:39:PHE:HZ	1:A:88:LYS:HG3	1.81	0.46
1:A:60:GLN:HG2	1:B:161:GLY:HA3	1.98	0.45
1:A:223:MET:HE2	1:A:223:MET:HB2	1.70	0.45
1:B:123:GLN:HE22	1:B:125:ALA:HB3	1.81	0.44
1:A:191:ALA:N	1:A:192:PRO:CD	2.81	0.43
1:A:101:GLU:HG3	1:A:114:ILE:HD12	2.00	0.43
1:B:50:ARG:NH1	1:B:234:GLU:OE1	2.37	0.42
1:A:94:SER:O	1:A:98:ARG:HG3	2.19	0.42
1:B:38:PRO:HD3	1:B:67:ALA:HB2	2.01	0.42
1:A:146:ALA:HB1	1:A:210:LEU:HD13	2.01	0.42
1:A:144:ARG:O	1:A:148:LEU:HG	2.19	0.42
1:A:188:PHE:HB3	1:A:208:TRP:CH2	2.55	0.42
1:A:37:SER:OG	1:A:38:PRO:HD2	2.20	0.42
1:B:19:THR:H	1:B:22:CYS:HB2	1.85	0.41
1:A:210:LEU:O	1:A:210:LEU:HD12	2.21	0.41
1:B:217:ASN:ND2	1:B:217:ASN:O	2.54	0.41
1:B:116:THR:HB	1:B:119:ASP:OD1	2.20	0.41
2:A:258:3CX:HNA1	2:A:258:3CX:HAK	1.81	0.41
1:A:33:LEU:HD11	1:A:65:LEU:HB2	2.02	0.40
1:A:17:THR:HG23	1:A:18:LEU:N	2.37	0.40
1:B:110:ASN:HA	1:B:111:PRO:HD2	1.84	0.40
1:A:210:LEU:O	1:A:214:ILE:HG13	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	209/257 (81%)	204 (98%)	5 (2%)	0	100	100
1	B	216/257 (84%)	209 (97%)	6 (3%)	1 (0%)	24	37
All	All	425/514 (83%)	413 (97%)	11 (3%)	1 (0%)	43	58

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	162	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	174/220 (79%)	171 (98%)	3 (2%)	53	74
1	B	176/220 (80%)	172 (98%)	4 (2%)	44	66
All	All	350/440 (80%)	343 (98%)	7 (2%)	48	70

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	THR
1	A	42	ARG
1	A	57	ARG
1	B	116	THR
1	B	210	LEU
1	B	227	GLN
1	B	233	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	23	ASN
1	B	123	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	3CX	B	258	-	14,15,15	2.19	2 (14%)	16,20,20	1.18	2 (12%)
2	3CX	A	258	-	14,15,15	2.17	4 (28%)	16,20,20	1.42	3 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3CX	B	258	-	1/1/3/4	3/10/18/18	0/1/1/1
2	3CX	A	258	-	1/1/3/4	4/10/18/18	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	258	3CX	OAC-CAM	-6.80	1.23	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	258	3CX	OAC-CAM	-6.46	1.24	1.43
2	A	258	3CX	CAJ-NAL	-2.34	1.41	1.47
2	B	258	3CX	CAK-CAM	-2.24	1.47	1.51
2	A	258	3CX	CAK-CAM	-2.10	1.48	1.51
2	A	258	3CX	OAA-SAO	2.02	1.50	1.45

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	258	3CX	OAC-CAM-CAJ	3.21	119.85	109.29
2	A	258	3CX	OAD-SAO-CAK	2.77	111.31	105.97
2	B	258	3CX	OAD-SAO-CAK	2.71	111.20	105.97
2	B	258	3CX	OAC-CAM-CAJ	2.42	117.24	109.29
2	A	258	3CX	OAA-SAO-CAK	2.07	109.85	106.76

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	258	3CX	CAM
2	B	258	3CX	CAM

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	258	3CX	NAL-CAJ-CAM-CAK
2	A	258	3CX	SAO-CAK-CAM-CAJ
2	A	258	3CX	CAH-CAN-NAL-CAJ
2	B	258	3CX	NAL-CAJ-CAM-OAC
2	B	258	3CX	SAO-CAK-CAM-OAC
2	B	258	3CX	SAO-CAK-CAM-CAJ
2	A	258	3CX	CAI-CAN-NAL-CAJ

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	258	3CX	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	213/257 (82%)	-0.08	7 (3%) 49 45	16, 31, 54, 61	0
1	B	220/257 (85%)	-0.15	9 (4%) 41 37	17, 28, 56, 75	0
All	All	433/514 (84%)	-0.12	16 (3%) 45 41	16, 30, 56, 75	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	165	ASP	3.7
1	A	167	VAL	3.6
1	B	18	LEU	3.5
1	B	166	VAL	3.5
1	B	167	VAL	3.2
1	A	39	PHE	3.2
1	A	157	GLY	3.0
1	B	161	GLY	2.9
1	B	164	MET	2.8
1	B	19	THR	2.8
1	A	168	GLU	2.4
1	B	158	ARG	2.3
1	B	39	PHE	2.2
1	A	18	LEU	2.2
1	A	16	GLU	2.2
1	A	171	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	3CX	A	258	15/15	0.89	0.12	28,38,52,66	0
2	3CX	B	258	15/15	0.95	0.09	26,32,35,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.