



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 07:19 PM UTC

PDB ID : 3OQL / pdb_00003oql
Title : Crystal structure of a TenA homolog (PSPTO1738) from *Pseudomonas syringae* pv. tomato str. DC3000 at 2.54 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2010-09-03
Resolution : 2.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	262	82%11%7%
1	B	262	82%11%7%
1	C	262	80%12%8%
1	D	262	78%14%7%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7812 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TenA homolog.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	243	Total	C	N	O	S	Se	0	2	0
			1949	1237	346	350	6	10			
1	B	244	Total	C	N	O	S	Se	0	2	0
			1951	1238	345	352	6	10			
1	C	242	Total	C	N	O	S	Se	0	1	0
			1932	1227	343	346	6	10			
1	D	244	Total	C	N	O	S	Se	0	1	0
			1923	1220	341	346	6	10			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q885U3
B	0	GLY	-	expression tag	UNP Q885U3
C	0	GLY	-	expression tag	UNP Q885U3
D	0	GLY	-	expression tag	UNP Q885U3


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	9	Total	O	0	0
			9	9		
2	B	7	Total	O	0	0
			7	7		
2	C	18	Total	O	0	0
			18	18		
2	D	23	Total	O	0	0
			23	23		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TenA homolog




- Molecule 1: TenA homolog




- Molecule 1: TenA homolog



- Molecule 1: TenA homolog





4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	67.34Å 67.34Å 479.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.88 – 2.54 29.88 – 2.54	Depositor EDS
% Data completeness (in resolution range)	(Not available) (29.88-2.54) 99.6 (29.88-2.54)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 2.54Å)	Xtriage
Refinement program	BUSTER-TNT BUSTER 2.8.0, BUSTER 2.8.0	Depositor
R, R_{free}	0.176 , 0.202 (Not available) , 0.209	Depositor DCC
R_{free} test set	1887 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	65.0	Xtriage
Anisotropy	0.011	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.41$, $\langle L^2 \rangle = 0.23$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7812	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/1996	1.32	7/2696 (0.3%)
1	B	0.83	1/1999 (0.1%)	1.32	6/2703 (0.2%)
1	C	0.87	2/1973 (0.1%)	1.32	7/2666 (0.3%)
1	D	0.88	2/1968 (0.1%)	1.33	8/2664 (0.3%)
All	All	0.86	5/7936 (0.1%)	1.32	28/10729 (0.3%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	54	MSE	SE-CE	-6.87	1.74	1.95
1	D	72	MSE	SE-CE	-6.49	1.75	1.95
1	D	54	MSE	SE-CE	-5.83	1.77	1.95
1	C	201	HIS	CA-C	5.55	1.58	1.52
1	B	72	MSE	SE-CE	-5.11	1.80	1.95

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	158	ILE	N-CA-CB	6.86	118.10	110.62
1	B	38	HIS	CA-C-N	6.51	129.30	120.38
1	B	38	HIS	C-N-CA	6.51	129.30	120.38
1	D	38	HIS	CA-C-N	6.42	129.18	120.38
1	D	38	HIS	C-N-CA	6.42	129.18	120.38
1	A	38	HIS	CA-C-N	6.27	128.97	120.38
1	A	38	HIS	C-N-CA	6.27	128.97	120.38
1	C	38	HIS	CA-C-N	6.21	128.88	120.38
1	C	38	HIS	C-N-CA	6.21	128.88	120.38
1	A	27	ASP	CA-CB-CG	5.72	118.32	112.60
1	D	13	GLU	CB-CG-CD	5.57	122.08	112.60
1	C	181	GLU	CA-C-N	5.57	127.74	120.28
1	C	181	GLU	C-N-CA	5.57	127.74	120.28
1	B	181	GLU	CA-C-N	5.56	127.73	120.28
1	B	181	GLU	C-N-CA	5.56	127.73	120.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	27	ASP	CA-CB-CG	5.55	118.15	112.60
1	A	181	GLU	CA-C-N	5.51	127.66	120.28
1	A	181	GLU	C-N-CA	5.51	127.66	120.28
1	C	44	MSE	CA-C-N	5.51	127.93	120.38
1	C	44	MSE	C-N-CA	5.51	127.93	120.38
1	B	199	ASP	CA-CB-CG	5.43	118.03	112.60
1	D	181	GLU	CA-C-N	5.39	127.51	120.28
1	D	181	GLU	C-N-CA	5.39	127.51	120.28
1	A	44	MSE	CA-C-N	5.29	127.62	120.38
1	A	44	MSE	C-N-CA	5.29	127.62	120.38
1	B	27	ASP	CA-CB-CG	5.28	117.88	112.60
1	C	27	ASP	CA-CB-CG	5.19	117.79	112.60
1	D	199	ASP	CA-CB-CG	5.06	117.66	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1949	0	1871	13	0
1	B	1951	0	1863	16	0
1	C	1932	0	1849	16	0
1	D	1923	0	1821	23	0
2	A	9	0	0	0	0
2	B	7	0	0	0	0
2	C	18	0	0	0	0
2	D	23	0	0	1	0
All	All	7812	0	7404	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:72:MSE:HE1	1:D:101:GLU:HG3	1.48	0.93
1:C:72:MSE:HE1	1:C:101:GLU:HG3	1.60	0.82
1:C:69:ALA:HA	1:C:72:MSE:HE3	1.69	0.74
1:A:66:GLU:O	1:B:91[B]:ARG:NH2	2.26	0.67
1:C:134:LEU:HD11	1:C:158:ILE:HD11	1.79	0.62
1:C:91[B]:ARG:NH2	1:D:66:GLU:O	2.34	0.60
1:A:72:MSE:HE1	1:A:101:GLU:HG3	1.85	0.58
1:D:69:ALA:HA	1:D:72:MSE:HE3	1.86	0.58
1:D:20:TRP:CZ2	1:D:213:VAL:HG21	2.39	0.57
1:D:91[A]:ARG:HG2	1:D:95:MSE:HE2	1.87	0.56
1:C:91[A]:ARG:HG2	1:C:95:MSE:HE2	1.88	0.56
1:B:75:ASN:HA	1:B:78:LYS:HD2	1.88	0.55
1:B:91[A]:ARG:HG2	1:B:95:MSE:HE2	1.86	0.55
1:C:66:GLU:O	1:D:91[B]:ARG:NH2	2.40	0.54
1:C:20:TRP:CZ2	1:C:213:VAL:HG21	2.43	0.53
1:D:89:MSE:HE2	1:D:208:ILE:HG13	1.90	0.53
1:C:102:LEU:HD12	1:D:95:MSE:HE3	1.92	0.52
1:B:20:TRP:CZ2	1:B:213:VAL:HG21	2.44	0.51
1:D:189:LYS:HD2	1:D:193:MSE:HE3	1.92	0.51
1:B:69:ALA:HA	1:B:72:MSE:HE2	1.93	0.51
1:A:72:MSE:CE	1:A:101:GLU:HG3	2.41	0.50
1:A:91[A]:ARG:HG2	1:A:95:MSE:HE2	1.94	0.50
1:D:20:TRP:CZ2	1:D:213:VAL:CG2	2.95	0.50
1:D:97:ASN:OD1	1:D:201:HIS:HE1	1.94	0.49
1:A:134:LEU:HD11	1:A:158:ILE:HD11	1.95	0.49
1:C:95:MSE:HE3	1:D:102:LEU:HD12	1.94	0.49
1:D:213:VAL:HG11	1:D:225:LEU:HD11	1.95	0.49
1:B:69:ALA:HA	1:B:72:MSE:CE	2.43	0.48
1:A:102:LEU:HD12	1:B:95:MSE:HE3	1.96	0.47
1:A:12:MSE:HG2	1:A:203:TRP:CD1	2.48	0.47
1:C:12:MSE:HG2	1:C:203:TRP:CD1	2.50	0.46
1:B:198:ASP:C	1:B:200:ALA:H	2.23	0.46
1:B:72:MSE:HE1	1:B:101:GLU:HG3	1.96	0.46
1:D:68:PHE:CD2	1:D:72:MSE:HE2	2.51	0.46
1:A:60:GLY:O	1:A:63:PRO:HD2	2.16	0.46
1:B:12:MSE:HG2	1:B:203:TRP:CD1	2.51	0.45
1:C:60:GLY:O	1:C:63:PRO:HD2	2.17	0.45
1:B:50:SER:HB3	1:B:53:ILE:HD12	1.98	0.45
1:B:60:GLY:O	1:B:63:PRO:HD2	2.16	0.45
1:D:60:GLY:O	1:D:63:PRO:HD2	2.16	0.44
1:D:43:GLN:NE2	2:D:266:HOH:O	2.50	0.44
1:D:154:THR:HA	1:D:158:ILE:HG12	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:50:SER:HB3	1:D:53:ILE:HD12	1.99	0.43
1:B:20:TRP:CZ2	1:B:213:VAL:CG2	3.01	0.43
1:A:124:ASP:HB3	1:A:126[B]:ARG:HE	1.84	0.43
1:C:20:TRP:CZ2	1:C:213:VAL:CG2	3.02	0.42
1:C:162:THR:HA	1:C:165:TRP:NE1	2.34	0.42
1:C:213:VAL:HG12	1:C:221:LEU:HD21	2.01	0.42
1:A:12:MSE:HE1	1:A:199:ASP:HB2	2.02	0.41
1:B:129:PRO:HB3	1:D:231:LYS:HG3	2.02	0.41
1:D:20:TRP:CH2	1:D:213:VAL:HG21	2.56	0.41
1:A:50:SER:HB3	1:A:53:ILE:HD12	2.02	0.41
1:D:127:VAL:HB	1:D:131:LEU:HD12	2.02	0.41
1:A:129:PRO:HB3	1:C:231:LYS:HG3	2.02	0.41
1:C:50:SER:HB3	1:C:53:ILE:HD12	2.02	0.41
1:D:162:THR:HA	1:D:165:TRP:NE1	2.35	0.41
1:A:162:THR:HA	1:A:165:TRP:NE1	2.36	0.40
1:B:162:THR:HA	1:B:165:TRP:NE1	2.36	0.40
1:D:242:ARG:HD2	1:D:242:ARG:HA	1.93	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	241/262 (92%)	236 (98%)	5 (2%)	0	100	100
1	B	244/262 (93%)	237 (97%)	7 (3%)	0	100	100
1	C	239/262 (91%)	235 (98%)	4 (2%)	0	100	100
1	D	243/262 (93%)	235 (97%)	8 (3%)	0	100	100
All	All	967/1048 (92%)	943 (98%)	24 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/208 (95%)	193 (98%)	5 (2%)	42	60
1	B	197/208 (95%)	191 (97%)	6 (3%)	36	53
1	C	195/208 (94%)	190 (97%)	5 (3%)	40	59
1	D	192/208 (92%)	189 (98%)	3 (2%)	55	73
All	All	782/832 (94%)	763 (98%)	19 (2%)	43	61

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	THR
1	A	92	ARG
1	A	186	LYS
1	A	220	GLN
1	A	244	ILE
1	B	92	ARG
1	B	120	GLU
1	B	182	GLU
1	B	186	LYS
1	B	220	GLN
1	B	244	ILE
1	C	92	ARG
1	C	159	GLU
1	C	186	LYS
1	C	220	GLN
1	C	244	ILE
1	D	92	ARG
1	D	186	LYS
1	D	244	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	GLN
1	A	215	ASN
1	B	42	GLN
1	B	104	HIS
1	B	215	ASN
1	C	42	GLN
1	C	67	GLN
1	C	215	ASN
1	D	42	GLN
1	D	43	GLN
1	D	67	GLN
1	D	104	HIS
1	D	201	HIS
1	D	215	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	233/262 (88%)	-0.21	2 (0%) 81 83	40, 70, 97, 118	2 (0%)
1	B	234/262 (89%)	-0.14	1 (0%) 88 90	38, 75, 104, 137	2 (0%)
1	C	232/262 (88%)	-0.31	3 (1%) 75 75	30, 64, 83, 100	1 (0%)
1	D	234/262 (89%)	-0.31	1 (0%) 88 90	36, 65, 88, 106	1 (0%)
All	All	933/1048 (89%)	-0.24	7 (0%) 82 84	30, 68, 95, 137	6 (0%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	246	LEU	3.6
1	C	199	ASP	3.4
1	D	246	LEU	3.1
1	A	197	TYR	2.7
1	C	91[A]	ARG	2.5
1	C	246	LEU	2.2
1	A	196	GLN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.