



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:00 AM UTC

PDB ID : 4OQB / pdb_00004oqb
Title : Structure of Human PARP-1 bound to a DNA double strand break in complex with (2Z)-2-{4-[2-(morpholin-4-yl)ethoxy]benzylidene}-3-oxo-2,3-dihydro-1-benzofuran-7-carboxamide
Authors : Pascal, J.M.; Steffen, J.D.
Deposited on : 2014-02-07
Resolution : 3.36 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.36 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1099 (3.40-3.32)
Clashscore	190562	1116 (3.40-3.32)
Ramachandran outliers	187476	1101 (3.40-3.32)
Sidechain outliers	187428	1101 (3.40-3.32)
RSRZ outliers	180081	1099 (3.40-3.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	 78% 5% 17%
1	D	267	 78% 5% 17%
2	C	505	 81% 9% 10%
2	F	505	 82% 9% 10%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	M	26	 88%12%
3	N	26	 92%8%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11775 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Poly [ADP-ribose] polymerase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	222	Total	C	N	O	S	0	1	0
			1755	1110	305	327	13			
1	D	222	Total	C	N	O	S	0	1	0
			1755	1110	305	327	13			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	205	ASP	-	linker	UNP P09874
A	206	ILE	-	linker	UNP P09874
A	367	LEU	-	expression tag	UNP P09874
A	368	GLU	-	expression tag	UNP P09874
A	369	HIS	-	expression tag	UNP P09874
A	370	HIS	-	expression tag	UNP P09874
A	371	HIS	-	expression tag	UNP P09874
A	372	HIS	-	expression tag	UNP P09874
A	373	HIS	-	expression tag	UNP P09874
A	374	HIS	-	expression tag	UNP P09874
D	205	ASP	-	linker	UNP P09874
D	206	ILE	-	linker	UNP P09874
D	367	LEU	-	expression tag	UNP P09874
D	368	GLU	-	expression tag	UNP P09874
D	369	HIS	-	expression tag	UNP P09874
D	370	HIS	-	expression tag	UNP P09874
D	371	HIS	-	expression tag	UNP P09874
D	372	HIS	-	expression tag	UNP P09874
D	373	HIS	-	expression tag	UNP P09874
D	374	HIS	-	expression tag	UNP P09874

- Molecule 2 is a protein called Poly [ADP-ribose] polymerase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	456	Total	C	N	O	S	0	0	0
			3586	2296	605	672	13			
2	F	456	Total	C	N	O	S	0	0	0
			3586	2296	605	672	13			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	1015	LEU	-	expression tag	UNP P09874
C	1016	GLU	-	expression tag	UNP P09874
C	1017	HIS	-	expression tag	UNP P09874
C	1018	HIS	-	expression tag	UNP P09874
C	1019	HIS	-	expression tag	UNP P09874
C	1020	HIS	-	expression tag	UNP P09874
C	1021	HIS	-	expression tag	UNP P09874
C	1022	HIS	-	expression tag	UNP P09874
F	1015	LEU	-	expression tag	UNP P09874
F	1016	GLU	-	expression tag	UNP P09874
F	1017	HIS	-	expression tag	UNP P09874
F	1018	HIS	-	expression tag	UNP P09874
F	1019	HIS	-	expression tag	UNP P09874
F	1020	HIS	-	expression tag	UNP P09874
F	1021	HIS	-	expression tag	UNP P09874
F	1022	HIS	-	expression tag	UNP P09874


- Molecule 3 is a DNA chain called DNA (26-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	M	26	Total	C	N	O	P	0	0	0
			530	251	100	154	25			
3	N	26	Total	C	N	O	P	0	0	0
			530	251	100	154	25			

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Zn	0	0
			2	2		
4	D	2	Total	Zn	0	0
			2	2		

- Molecule 5 is (2Z)-2-{4-[2-(morpholin-4-yl)ethoxy]benzylidene}-3-oxo-2,3-dihydro-1-benzofuran-7-carboxamide (CCD ID: 2UT) (formula: C₂₂H₂₂N₂O₅).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	1	Total	C	N	O	0	0
			29	22	2	5		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Poly [ADP-ribose] polymerase 1

Chain A: 




• Molecule 1: Poly [ADP-ribose] polymerase 1

Chain D: 




• Molecule 2: Poly [ADP-ribose] polymerase 1

Chain C: 




• Molecule 2: Poly [ADP-ribose] polymerase 1


Chain F: 



• Molecule 3: DNA (26-MER)



• Molecule 3: DNA (26-MER)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.10Å 113.08Å 296.23Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 3.36 20.00 – 3.36	Depositor EDS
% Data completeness (in resolution range)	98.9 (20.00-3.36) 98.6 (20.00-3.36)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.99 (at 3.33Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.322 , 0.349 0.322 , 0.356	Depositor DCC
R_{free} test set	1627 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	136.5	Xtriage
Anisotropy	0.161	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.25 , 115.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	11775	wwPDB-VP
Average B, all atoms (Å ²)	213.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 2UT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/1795	0.63	0/2416
1	D	0.37	0/1795	0.63	0/2416
2	C	0.39	0/3660	0.64	0/4940
2	F	0.38	0/3660	0.64	0/4940
3	M	0.31	0/594	0.72	2/915 (0.2%)
3	N	0.29	0/594	0.65	2/915 (0.2%)
All	All	0.37	0/12098	0.64	4/16542 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	N	1	DG	C5'-C4'-O4'	6.56	119.23	109.40
3	M	1	DG	C5'-C4'-O4'	5.91	118.26	109.40
3	N	1	DG	C5'-C4'-C3'	5.57	123.25	114.90
3	M	1	DG	C5'-C4'-C3'	5.29	122.84	114.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1755	0	1712	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	1755	0	1712	6	0
2	C	3586	0	3604	21	0
2	F	3586	0	3604	18	0
3	M	530	0	292	1	0
3	N	530	0	292	1	0
4	A	2	0	0	0	0
4	D	2	0	0	0	0
5	C	29	0	22	2	0
All	All	11775	0	11238	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:574:LEU:HD11	2:F:610:ALA:HB1	1.69	0.73
2:C:574:LEU:HD11	2:C:610:ALA:HB1	1.70	0.72
1:A:36:ALA:HB2	1:A:51:TRP:CE3	2.39	0.57
1:D:36:ALA:HB2	1:D:51:TRP:CE3	2.39	0.57
2:C:831:LEU:HD12	2:C:1005:LEU:HD23	1.88	0.54
2:F:831:LEU:HD12	2:F:1005:LEU:HD23	1.88	0.54
2:C:898:ALA:HB2	5:C:1101:2UT:H2	1.91	0.53
2:F:814:ILE:HG21	2:F:836:ILE:HD12	1.92	0.52
2:C:689:TYR:HB3	2:C:764:MET:HE2	1.92	0.51
2:C:814:ILE:HG21	2:C:836:ILE:HD12	1.92	0.51
2:F:679:VAL:HG11	2:F:775:TYR:CZ	2.46	0.50
2:C:679:VAL:HG11	2:C:775:TYR:CZ	2.47	0.50
2:F:919:ILE:HG22	2:F:1005:LEU:HD11	1.94	0.49
2:C:919:ILE:HG22	2:C:1005:LEU:HD11	1.94	0.49
2:F:689:TYR:HB3	2:F:764:MET:HE2	1.92	0.49
2:F:696:MET:N	2:F:697:PRO:HD2	2.28	0.49
2:C:696:MET:N	2:C:697:PRO:HD2	2.28	0.48
1:D:315:VAL:HG13	1:D:316:THR:HG22	1.95	0.48
2:C:929:MET:HE2	2:C:949:LYS:HB2	1.95	0.48
2:F:929:MET:HE2	2:F:949:LYS:HB2	1.95	0.48
1:A:315:VAL:HG13	1:A:316:THR:HG22	1.96	0.47
2:C:861:TRP:CD2	2:C:921:LEU:HD21	2.49	0.47
1:D:319:THR:HG21	2:F:731:ASP:HA	1.96	0.47
3:M:22:DT:H2"	3:M:23:DA:C8	2.50	0.47
2:C:862:HIS:ND1	5:C:1101:2UT:O	2.48	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:861:TRP:CD2	2:F:921:LEU:HD21	2.49	0.46
1:A:275:GLY:HA2	3:N:23:DA:H3'	1.99	0.44
2:F:799:THR:HG22	2:F:841:ARG:HA	2.00	0.43
2:C:539:LEU:HD11	2:C:601:LEU:HD21	2.01	0.43
1:D:319:THR:CG2	2:F:731:ASP:HA	2.48	0.42
1:D:64:ILE:HD13	1:D:71:VAL:CG2	2.50	0.42
2:C:814:ILE:CG2	2:C:1003:LEU:HD21	2.50	0.42
1:A:64:ILE:HD13	1:A:71:VAL:CG2	2.50	0.42
2:C:799:THR:HG22	2:C:841:ARG:HA	2.00	0.42
2:F:539:LEU:HD11	2:F:601:LEU:HD21	2.00	0.41
2:C:682:MET:HG2	2:C:867:THR:HG22	2.02	0.41
2:C:734:ASN:O	2:C:738:THR:OG1	2.38	0.41
2:F:682:MET:HG2	2:F:867:THR:HG22	2.03	0.41
2:F:696:MET:N	2:F:697:PRO:CD	2.83	0.41
2:C:990:ILE:HD12	2:C:990:ILE:N	2.36	0.41
1:A:319:THR:HG22	2:C:734:ASN:HD22	1.86	0.41
2:C:618:TYR:CE2	2:C:622:THR:HG21	2.56	0.41
2:C:696:MET:N	2:C:697:PRO:CD	2.83	0.41
2:F:814:ILE:CG2	2:F:1003:LEU:HD21	2.50	0.41
1:D:36:ALA:HB2	1:D:51:TRP:CZ3	2.56	0.41
2:F:990:ILE:N	2:F:990:ILE:HD12	2.36	0.41
2:C:618:TYR:O	2:C:622:THR:HG22	2.21	0.40
1:A:36:ALA:HB2	1:A:51:TRP:CZ3	2.56	0.40
2:F:618:TYR:O	2:F:622:THR:HG22	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	219/267 (82%)	212 (97%)	7 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	219/267 (82%)	212 (97%)	7 (3%)	0	100	100
2	C	450/505 (89%)	423 (94%)	27 (6%)	0	100	100
2	F	450/505 (89%)	422 (94%)	28 (6%)	0	100	100
All	All	1338/1544 (87%)	1269 (95%)	69 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/237 (80%)	180 (95%)	9 (5%)	23	49
1	D	189/237 (80%)	181 (96%)	8 (4%)	26	52
2	C	391/443 (88%)	375 (96%)	16 (4%)	27	53
2	F	391/443 (88%)	375 (96%)	16 (4%)	27	53
All	All	1160/1360 (85%)	1111 (96%)	49 (4%)	26	52

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	21	CYS
1	A	22	LYS
1	A	37	ILE
1	A	40	GLN
1	A	48	VAL
1	A	236	LYS
1	A	301	GLN
1	A	316	THR
1	A	319	THR
2	C	592	VAL
2	C	594	THR
2	C	599	ASN
2	C	662	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	C	698	LEU
2	C	730	LEU
2	C	738	THR
2	C	758	VAL
2	C	769	LEU
2	C	798	LYS
2	C	835	ASP
2	C	847	ARG
2	C	887	THR
2	C	890	MET
2	C	916	ILE
2	C	1002	LEU
1	D	21	CYS
1	D	22	LYS
1	D	37	ILE
1	D	40	GLN
1	D	48	VAL
1	D	301	GLN
1	D	316	THR
1	D	319	THR
2	F	592	VAL
2	F	594	THR
2	F	599	ASN
2	F	662	LYS
2	F	698	LEU
2	F	730	LEU
2	F	738	THR
2	F	758	VAL
2	F	769	LEU
2	F	798	LYS
2	F	835	ASP
2	F	847	ARG
2	F	887	THR
2	F	890	MET
2	F	916	ILE
2	F	1002	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (35) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	HIS
1	A	62	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	83	GLN
1	A	329	ASN
2	C	541	HIS
2	C	567	ASN
2	C	573	GLN
2	C	599	ASN
2	C	767	ASN
2	C	793	ASN
2	C	856	ASN
2	C	868	ASN
2	C	906	ASN
2	C	961	ASN
2	C	980	ASN
2	C	1008	ASN
1	D	40	GLN
1	D	50	HIS
1	D	62	HIS
1	D	83	GLN
1	D	329	ASN
2	F	541	HIS
2	F	567	ASN
2	F	573	GLN
2	F	599	ASN
2	F	627	HIS
2	F	767	ASN
2	F	793	ASN
2	F	820	ASN
2	F	856	ASN
2	F	868	ASN
2	F	906	ASN
2	F	961	ASN
2	F	980	ASN
2	F	1008	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	2UT	C	1101	-	32,32,32	0.26	0	44,44,44	3.32	8 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	2UT	C	1101	-	-	4/14/34/34	0/4/4/4

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	1101	2UT	O1-C3-C4	-13.57	112.75	124.39
5	C	1101	2UT	C4-C3-C17	11.12	135.37	126.25
5	C	1101	2UT	C18-C17-C3	-7.45	100.75	104.08
5	C	1101	2UT	O4-C17-C3	6.83	132.27	126.21
5	C	1101	2UT	C2-C18-C17	6.52	109.63	106.91
5	C	1101	2UT	O1-C3-C17	3.64	111.79	109.47
5	C	1101	2UT	C2-O1-C3	2.42	107.18	105.52
5	C	1101	2UT	C2-C1-C	-2.40	121.86	125.74

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	C	1101	2UT	C9-C10-N1-C14
5	C	1101	2UT	O-C-C1-C2
5	C	1101	2UT	N-C-C1-C2
5	C	1101	2UT	C9-C10-N1-C11

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	1101	2UT	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	222/267 (83%)	-0.28	1 (0%) 87 78	81, 174, 261, 285	1 (0%)
1	D	222/267 (83%)	-0.51	0 100 100	112, 243, 321, 342	1 (0%)
2	C	456/505 (90%)	-0.19	2 (0%) 88 81	135, 195, 248, 286	0
2	F	456/505 (90%)	-0.33	2 (0%) 88 81	169, 241, 300, 331	0
3	M	26/26 (100%)	-0.50	0 100 100	161, 187, 226, 228	0
3	N	26/26 (100%)	-0.41	0 100 100	157, 203, 226, 230	0
All	All	1408/1596 (88%)	-0.31	5 (0%) 88 81	81, 214, 293, 342	2 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	771	ILE	3.1
1	A	264	LEU	2.6
2	F	771	ILE	2.5
2	C	570	TYR	2.5
2	F	943	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	2UT	C	1101	29/29	0.75	0.10	121,127,140,140	0
4	ZN	D	401	1/1	0.94	0.15	118,118,118,118	0
4	ZN	D	402	1/1	0.96	0.13	132,132,132,132	0
4	ZN	A	402	1/1	0.97	0.17	62,62,62,62	0
4	ZN	A	401	1/1	0.98	0.15	78,78,78,78	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.