



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 06:46 AM UTC

PDB ID : 5OQ5 / pdb_00005oq5
Title : Structure of CHK1 8-pt. mutant complex with aminopyrimido-benzodiazepin
one LRRK2 inhibitor
Authors : Dokurno, P.; Williamson, D.S.; Acheson-Dossang, P.; Chen, I.; Murray, J.B.;
Shaw, T.; Surgenor, A.E.
Deposited on : 2017-08-10
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	297	6% 67% 19% • 11%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2447 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Serine/threonine-protein kinase Chk1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	265	Total	C	N	O	S	0	0	0
			2106	1350	364	381	11			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	59	LEU	ASN	engineered mutation	UNP O14757
A	68	ILE	VAL	engineered mutation	UNP O14757
A	84	MET	LEU	engineered mutation	UNP O14757
A	86	LEU	TYR	engineered mutation	UNP O14757
A	87	ALA	CYS	engineered mutation	UNP O14757
A	91	SER	GLU	engineered mutation	UNP O14757
A	134	HIS	GLU	engineered mutation	UNP O14757
A	147	ALA	SER	engineered mutation	UNP O14757
A	290	HIS	-	expression tag	UNP O14757
A	291	HIS	-	expression tag	UNP O14757
A	292	HIS	-	expression tag	UNP O14757
A	293	HIS	-	expression tag	UNP O14757
A	294	HIS	-	expression tag	UNP O14757
A	295	HIS	-	expression tag	UNP O14757
A	296	HIS	-	expression tag	UNP O14757
A	297	HIS	-	expression tag	UNP O14757

- Molecule 2 is 2-[(2-methoxy-4-{[4-(4-methylpiperazin-1-yl)piperidin-1-yl]carbonyl}phenyl)amino]-5,11-dimethyl-5,11-dihydro-6H-pyrimido[4,5-b][1,4]benzodiazepin-6-one (CCD ID: 4K4) (formula: C₃₁H₃₈N₈O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			42	31	8	3		

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	287	Total 287	O 287	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Serine/threonine-protein kinase Chk1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	45.40Å 66.57Å 55.08Å 90.00° 101.58° 90.00°	Depositor
Resolution (Å)	20.00 – 1.40 20.00 – 1.40	Depositor EDS
% Data completeness (in resolution range)	96.6 (20.00-1.40) 96.6 (20.00-1.40)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.169 , 0.185 (Not available) , 0.202	Depositor DCC
R_{free} test set	3013 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	21.2	Xtriage
Anisotropy	0.055	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 45.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2447	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 4K4, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.87	42/2156 (1.9%)	1.51	13/2926 (0.4%)

All (42) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	181	ARG	CA-C	-9.53	1.41	1.52
1	A	237	ALA	CA-C	8.89	1.62	1.52
1	A	171	LEU	CA-CB	7.91	1.63	1.53
1	A	211	PRO	N-CA	-7.37	1.38	1.47
1	A	177	GLU	CA-C	-7.14	1.43	1.52
1	A	91	SER	N-CA	7.09	1.54	1.45
1	A	268	PRO	CA-CB	7.08	1.62	1.53
1	A	267	LYS	CA-C	-7.06	1.45	1.52
1	A	247	VAL	N-CA	7.01	1.55	1.46
1	A	180	LYS	N-CA	6.50	1.54	1.45
1	A	246	LEU	CA-C	6.50	1.60	1.53
1	A	234	ILE	CA-CB	6.28	1.61	1.54
1	A	106	ASP	CA-C	6.14	1.60	1.52
1	A	156	ARG	C-N	6.11	1.41	1.33
1	A	211	PRO	CA-C	6.11	1.58	1.53
1	A	85	GLU	C-O	6.02	1.31	1.23
1	A	160	ARG	N-CA	-6.02	1.38	1.46
1	A	253	ARG	N-CA	6.01	1.53	1.45
1	A	182	ARG	CA-CB	5.83	1.62	1.53
1	A	132	LYS	CA-CB	5.75	1.62	1.54
1	A	160	ARG	CA-C	5.72	1.59	1.52
1	A	264	TRP	C-N	5.71	1.41	1.33
1	A	215	CYS	CA-CB	5.63	1.60	1.53
1	A	173	TYR	N-CA	5.63	1.53	1.46
1	A	158	ASN	CG-OD1	5.63	1.34	1.23
1	A	156	ARG	N-CA	5.59	1.53	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	134	HIS	C-O	5.41	1.30	1.24
1	A	191	VAL	CA-CB	5.39	1.61	1.54
1	A	90	GLY	CA-C	5.39	1.58	1.52
1	A	141	ARG	CA-C	5.26	1.60	1.52
1	A	147	ALA	N-CA	5.25	1.52	1.45
1	A	121	LEU	C-O	5.24	1.30	1.24
1	A	126	ILE	CB-CG1	5.21	1.63	1.53
1	A	219	SER	N-CA	5.20	1.52	1.46
1	A	126	ILE	CA-C	5.17	1.58	1.52
1	A	152	ALA	CA-CB	-5.17	1.44	1.53
1	A	220	ASP	C-N	5.14	1.40	1.33
1	A	187	GLU	N-CA	5.12	1.51	1.46
1	A	138	LEU	CA-C	5.10	1.58	1.52
1	A	252	ALA	CA-C	5.07	1.59	1.52
1	A	70	PHE	CA-CB	-5.06	1.46	1.53
1	A	139	ASP	N-CA	5.03	1.52	1.45

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	167	MET	CB-CA-C	-6.65	100.28	110.26
1	A	267	LYS	CA-C-O	6.56	125.59	119.76
1	A	141	ARG	NE-CZ-NH2	6.45	125.01	119.20
1	A	219	SER	N-CA-C	5.59	117.37	111.28
1	A	3	VAL	CA-C-N	-5.58	115.01	120.98
1	A	3	VAL	C-N-CA	-5.58	115.01	120.98
1	A	248	GLU	N-CA-C	5.54	117.00	111.07
1	A	175	ALA	O-C-N	5.35	128.44	121.64
1	A	256	ILE	N-CA-CB	5.23	113.80	110.50
1	A	239	LEU	CA-C-O	-5.18	115.06	120.55
1	A	120	TYR	O-C-N	-5.09	116.83	122.07
1	A	156	ARG	NE-CZ-NH1	5.07	126.57	121.50
1	A	97	GLU	O-C-N	-5.05	117.02	121.31

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2106	0	2079	19	0
2	A	42	0	38	2	0
3	A	12	0	13	0	0
4	A	287	0	0	11	5
All	All	2447	0	2130	21	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:LYS:HG3	4:A:402:HOH:O	1.54	1.05
1:A:134:HIS:N	4:A:402:HOH:O	2.01	0.91
1:A:32:GLU:OE1	4:A:403:HOH:O	2.07	0.71
1:A:134:HIS:HB2	4:A:402:HOH:O	1.90	0.71
1:A:213:ASP:HA	4:A:441:HOH:O	1.92	0.68
1:A:10:ASP:OD1	1:A:29:ARG:NH1	2.31	0.63
1:A:180:LYS:HE3	4:A:430:HOH:O	1.99	0.61
1:A:52:ILE:HD12	1:A:52:ILE:N	2.18	0.58
1:A:8:ASP:OD1	1:A:8:ASP:N	2.32	0.57
1:A:10:ASP:OD1	1:A:29:ARG:NE	2.39	0.55
2:A:301:4K4:CAN	2:A:301:4K4:CAQ	2.84	0.55
1:A:162:ARG:NH1	4:A:407:HOH:O	2.38	0.52
1:A:177:GLU:HB2	1:A:181:ARG:HD3	1.95	0.47
2:A:301:4K4:CAN	2:A:301:4K4:H23	2.45	0.45
1:A:260:LYS:HE2	4:A:605:HOH:O	2.15	0.45
1:A:181:ARG:HD2	4:A:626:HOH:O	2.17	0.43
1:A:238:PRO:HD3	1:A:264:TRP:CD1	2.53	0.43
1:A:29:ARG:HG2	4:A:535:HOH:O	2.18	0.42
1:A:140:GLU:OE2	1:A:141:ARG:NH2	2.52	0.42
1:A:141:ARG:HH21	1:A:141:ARG:HG3	1.85	0.42
1:A:68:ILE:HG21	4:A:564:HOH:O	2.20	0.41

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:643:HOH:O	4:A:676:HOH:O[2_555]	1.68	0.52
4:A:636:HOH:O	4:A:637:HOH:O[1_554]	1.86	0.34
4:A:581:HOH:O	4:A:597:HOH:O[2_545]	1.94	0.26
4:A:611:HOH:O	4:A:670:HOH:O[2_555]	1.96	0.24
4:A:632:HOH:O	4:A:670:HOH:O[2_555]	2.09	0.11

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/297 (88%)	253 (97%)	8 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/259 (86%)	211 (94%)	13 (6%)	18	1

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	ASP
1	A	11	LEU
1	A	12	VAL
1	A	25	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	40	VAL
1	A	78	ASN
1	A	82	LEU
1	A	141	ARG
1	A	162	ARG
1	A	174	VAL
1	A	213	ASP
1	A	224	LYS
1	A	232	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	GLN
1	A	24	GLN
1	A	73	HIS
1	A	78	ASN
1	A	80	GLN
1	A	143	ASN
1	A	158	ASN
1	A	165	ASN
1	A	216	GLN
1	A	243	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MES	A	302	-	12,12,12	2.01	4 (33%)	15,16,16	2.30	6 (40%)
2	4K4	A	301	-	45,47,47	1.84	12 (26%)	60,68,68	1.85	13 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MES	A	302	-	-	0/6/14/14	0/1/1/1
2	4K4	A	301	-	-	4/18/38/38	0/6/6/6

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	4K4	CBA-NBM	4.50	1.44	1.34
3	A	302	MES	C8-S	4.32	1.83	1.77
2	A	301	4K4	CAQ-NBM	4.08	1.54	1.47
2	A	301	4K4	CAR-NBM	3.62	1.53	1.47
2	A	301	4K4	OAZ-CAA	3.47	1.52	1.42
2	A	301	4K4	CBH-NBP	3.38	1.46	1.43
2	A	301	4K4	C5-NBO	-3.21	1.34	1.42
2	A	301	4K4	CAP-CAR	2.74	1.59	1.52
3	A	302	MES	O2S-S	2.68	1.52	1.45
2	A	301	4K4	C2-N1	-2.61	1.31	1.34
3	A	302	MES	O1-C2	2.50	1.52	1.42
2	A	301	4K4	CAO-CBK	-2.33	1.46	1.52
2	A	301	4K4	C6-N1	2.28	1.39	1.34
3	A	302	MES	O3S-S	2.22	1.55	1.47
2	A	301	4K4	C2-N3	2.09	1.40	1.34
2	A	301	4K4	CBK-NBN	2.03	1.53	1.48

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	4K4	C5-C4-NBP	6.49	124.32	120.14
3	A	302	MES	O3S-S-C8	-5.13	95.97	106.00
2	A	301	4K4	N1-C2-N3	-4.34	122.23	126.42
2	A	301	4K4	CAJ-CBC-NAY	3.76	128.83	121.32
2	A	301	4K4	CAL-CBH-NBP	-3.75	114.81	120.66
2	A	301	4K4	CAQ-NBM-CAR	3.44	119.70	112.68
3	A	302	MES	O2S-S-C8	3.27	111.67	106.73
2	A	301	4K4	CAH-CAG-CAK	3.04	124.00	120.24
3	A	302	MES	O1-C2-C3	-2.85	105.63	111.77
3	A	302	MES	C7-N4-C5	-2.82	103.72	111.24
2	A	301	4K4	CAJ-CBC-CBE	-2.82	115.37	119.02
2	A	301	4K4	CAG-CAK-CBG	-2.77	114.79	119.80
2	A	301	4K4	C6-C5-C4	-2.69	115.33	117.77
3	A	302	MES	O3S-S-O1S	2.55	117.79	111.40
2	A	301	4K4	CBB-CAN-CBE	2.44	122.87	119.60
3	A	302	MES	O2S-S-O1S	-2.29	106.37	113.82
2	A	301	4K4	CAL-CBH-CBG	2.14	122.44	119.16
2	A	301	4K4	CAK-CBG-CBH	2.13	121.25	118.81
2	A	301	4K4	C6-N1-C2	2.05	118.62	115.81

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	4K4	CAP-CBK-NBN-CAV
2	A	301	4K4	CAO-CBK-NBN-CAV
2	A	301	4K4	CAP-CBK-NBN-CAU
2	A	301	4K4	CAO-CBK-NBN-CAU


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	4K4	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	265/297 (89%)	0.28	18 (6%) 23 24	14, 26, 65, 90	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	30	VAL	4.5
1	A	27	VAL	4.3
1	A	49	PRO	3.2
1	A	20	TYR	3.2
1	A	3	VAL	3.1
1	A	79	ILE	3.0
1	A	6	VAL	2.9
1	A	31	THR	2.9
1	A	5	PHE	2.7
1	A	12	VAL	2.6
1	A	19	ALA	2.5
1	A	48	CYS	2.4
1	A	83	PHE	2.4
1	A	4	PRO	2.3
1	A	52	ILE	2.1
1	A	81	TYR	2.1
1	A	9	TRP	2.0
1	A	78	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MES	A	302	12/12	0.91	0.10	30,35,48,49	0
2	4K4	A	301	42/42	0.93	0.11	23,33,69,71	12

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 4K4 A 301:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.