



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 09:45 AM UTC

PDB ID : 1ORW / pdb_00001orw
Title : Crystal Structure of Porcine Dipeptidyl Peptidase IV (CD26) in Complex with a Peptidomimetic Inhibitor
Authors : Engel, M.; Hoffmann, T.; Wagner, L.; Wermann, M.; Heiser, U.; Kiefersauer, R.; Huber, R.; Bode, W.; Demuth, H.U.; Brandstetter, H.
Deposited on : 2003-03-16
Resolution : 2.84 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1520 (2.86-2.82)
Clashscore	190562	1559 (2.86-2.82)
Ramachandran outliers	187476	1517 (2.86-2.82)
Sidechain outliers	187428	1518 (2.86-2.82)
RSRZ outliers	180081	1521 (2.86-2.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	728	0% 61% 35% •
1	B	728	60% 36% •
1	C	728	2% 53% 42% • •
1	D	728	58% 37% • •
2	E	2	50% 50%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	2	 100%
2	H	2	 100%
2	I	2	 100%
2	K	2	 50% 50%
2	L	2	 100%
2	M	2	 100%
3	G	3	 33% 33% 33%
3	J	3	 33% 67%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	PHI	C	805	-	-	X	-
6	PHI	D	807	-	-	X	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 25262 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called dipeptidyl peptidase IV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	728	Total	C	N	O	S	71	0	0
			5966	3825	986	1132	23			
1	B	728	Total	C	N	O	S	42	0	0
			5966	3825	986	1132	23			
1	C	728	Total	C	N	O	S	62	0	0
			5966	3825	986	1132	23			
1	D	728	Total	C	N	O	S	36	0	0
			5966	3825	986	1132	23			


- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	F	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	H	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	I	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	K	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	L	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	M	2	Total	C	N	O	0	0	0
			28	16	2	10			


- Molecule 3 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-b

eta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	G	3	Total	C	N	O	0	0	0
			39	22	2	15			
3	J	3	Total	C	N	O	0	0	0
			39	22	2	15			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	3	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	O	S	0	0
			5	4	1		
5	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is IODO-PHENYLALANINE (CCD ID: PHI) (formula: $C_9H_{10}INO_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	I	N	O	0	0
			12	9	1	1	1		
6	B	1	Total	C	I	N	O	0	0
			12	9	1	1	1		
6	C	1	Total	C	I	N	O	0	0
			12	9	1	1	1		
6	D	1	Total	C	I	N	O	0	0
			12	9	1	1	1		

- Molecule 7 is (2S)-PYRROLIDIN-2-YLMETHYLAMINE (CCD ID: P2Y) (formula: $C_5H_{12}N_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	N	0	0
			7	5	2		
7	B	1	Total	C	N	0	0
			7	5	2		
7	C	1	Total	C	N	0	0
			7	5	2		
7	D	1	Total	C	N	0	0
			7	5	2		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	206	Total	O	0	0
			206	206		
8	B	226	Total	O	0	0
			226	226		
8	C	213	Total	O	0	0
			213	213		
8	D	173	Total	O	0	0
			173	173		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: dipeptidyl peptidase IV





• Molecule 1: dipeptidyl peptidase IV






• Molecule 1: dipeptidyl peptidase IV





- Molecule 1: dipeptidyl peptidase IV



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  100%MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  100%MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain K:  50% 50%MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain L:  100%MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M:  100%MAG1
MAG2

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  33% 33% 33%MAG1
MAG2
EMAG

- Molecule 3: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  33% 67%

MAG1
MAG2
EMJ3

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	61.97Å 117.71Å 133.56Å 112.65° 94.81° 91.17°	Depositor
Resolution (Å)	29.79 – 2.84 29.79 – 2.84	Depositor EDS
% Data completeness (in resolution range)	96.3 (29.79-2.84) 97.6 (29.79-2.84)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.12 (at 2.85Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.187 , 0.246 0.177 , 0.235	Depositor DCC
R_{free} test set	7805 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	30.5	Xtriage
Anisotropy	0.613	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 57.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	25262	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PHI, BMA, NAG, P2Y, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.49	0/6141	0.98	27/8353 (0.3%)
1	B	0.50	0/6141	0.99	25/8353 (0.3%)
1	C	0.52	0/6141	1.01	37/8353 (0.4%)
1	D	0.46	0/6141	0.97	32/8353 (0.4%)
All	All	0.49	0/24564	0.99	121/33412 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

The worst 5 of 121 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	104	ASP	N-CA-C	-10.75	87.90	110.80
1	D	378	GLU	N-CA-C	-9.77	101.36	113.18
1	C	103	ASN	N-CA-C	9.44	130.91	110.80
1	A	378	GLU	N-CA-C	-9.44	101.18	112.89
1	C	205	GLU	N-CA-C	9.09	120.95	111.14

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	468	TYR	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5966	0	5662	216	0
1	B	5966	0	5662	224	0
1	C	5966	0	5661	280	0
1	D	5966	0	5662	257	0
2	E	28	0	25	0	0
2	F	28	0	25	1	0
2	H	28	0	25	0	0
2	I	28	0	25	0	0
2	K	28	0	25	0	0
2	L	28	0	25	0	0
2	M	28	0	25	1	0
3	G	39	0	34	1	0
3	J	39	0	34	2	0
4	A	56	0	52	1	0
4	B	56	0	52	5	0
4	C	70	0	65	2	0
4	D	28	0	26	1	0
5	A	5	0	0	0	0
5	B	5	0	0	0	0
5	C	5	0	0	0	0
5	D	5	0	0	0	0
6	A	12	0	9	3	0
6	B	12	0	9	4	0
6	C	12	0	9	7	0
6	D	12	0	9	7	0
7	A	7	0	9	2	0
7	B	7	0	9	3	0
7	C	7	0	9	1	0
7	D	7	0	9	2	0
8	A	206	0	0	10	0
8	B	226	0	0	15	0
8	C	213	0	0	18	0
8	D	173	0	0	5	0
All	All	25262	0	23157	956	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 956 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:358:ARG:NH2	6:D:807:PHI:I	2.51	1.13
1:B:358:ARG:NH2	6:B:803:PHI:I	2.57	1.08
1:A:358:ARG:NH2	6:A:801:PHI:I	2.60	1.05
1:C:535:ASP:C	1:C:536:LYS:HD3	1.83	1.04
1:B:358:ARG:NH1	6:B:803:PHI:I	2.65	0.99

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	726/728 (100%)	656 (90%)	58 (8%)	12 (2%)	7	15
1	B	726/728 (100%)	670 (92%)	50 (7%)	6 (1%)	16	31
1	C	726/728 (100%)	652 (90%)	59 (8%)	15 (2%)	5	12
1	D	726/728 (100%)	656 (90%)	64 (9%)	6 (1%)	16	31
All	All	2904/2912 (100%)	2634 (91%)	231 (8%)	39 (1%)	9	20

5 of 39 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	40	ARG
1	B	40	ARG
1	C	103	ASN
1	C	534	PHE
1	C	535	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	652/652 (100%)	624 (96%)	28 (4%)	26	50
1	B	652/652 (100%)	624 (96%)	28 (4%)	26	50
1	C	652/652 (100%)	619 (95%)	33 (5%)	21	44
1	D	652/652 (100%)	619 (95%)	33 (5%)	21	44
All	All	2608/2608 (100%)	2486 (95%)	122 (5%)	23	47

5 of 122 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	90	LEU
1	D	536	LYS
1	C	418	ILE
1	D	507	VAL
1	D	701	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 115 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	761	GLN
1	D	679	ASN
1	C	388	GLN
1	D	612	GLN
1	D	247	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

20 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	E	1	2,1	14,14,15	0.46	0	17,19,21	0.63	0
2	NAG	E	2	2	14,14,15	0.55	0	17,19,21	0.78	1 (5%)
2	NAG	F	1	2,1	14,14,15	0.55	0	17,19,21	0.71	0
2	NAG	F	2	2	14,14,15	0.56	0	17,19,21	0.76	1 (5%)
3	NAG	G	1	3,1	14,14,15	0.56	0	17,19,21	0.86	1 (5%)
3	NAG	G	2	3	14,14,15	0.64	0	17,19,21	0.78	0
3	BMA	G	3	3	11,11,12	0.51	0	15,15,17	0.46	0
2	NAG	H	1	2,1	14,14,15	0.60	0	17,19,21	0.77	0
2	NAG	H	2	2	14,14,15	0.54	0	17,19,21	0.80	0
2	NAG	I	1	2,1	14,14,15	0.55	0	17,19,21	0.65	0
2	NAG	I	2	2	14,14,15	0.65	0	17,19,21	0.67	0
3	NAG	J	1	3,1	14,14,15	0.67	0	17,19,21	0.81	0
3	NAG	J	2	3	14,14,15	0.58	0	17,19,21	0.55	0
3	BMA	J	3	3	11,11,12	0.69	0	15,15,17	0.63	0
2	NAG	K	1	2,1	14,14,15	0.51	0	17,19,21	0.67	1 (5%)
2	NAG	K	2	2	14,14,15	0.54	0	17,19,21	0.64	0
2	NAG	L	1	2,1	14,14,15	0.58	0	17,19,21	1.01	2 (11%)
2	NAG	L	2	2	14,14,15	0.64	0	17,19,21	0.82	1 (5%)
2	NAG	M	1	2,1	14,14,15	0.54	0	17,19,21	0.75	0
2	NAG	M	2	2	14,14,15	0.58	0	17,19,21	1.37	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	E	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	E	2	2	-	0/6/23/26	0/1/1/1
2	NAG	F	1	2,1	-	2/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	F	2	2	-	4/6/23/26	0/1/1/1
3	NAG	G	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	G	2	3	-	2/6/23/26	0/1/1/1
3	BMA	G	3	3	-	2/2/19/22	0/1/1/1
2	NAG	H	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	H	2	2	-	0/6/23/26	0/1/1/1
2	NAG	I	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	I	2	2	-	0/6/23/26	0/1/1/1
3	NAG	J	1	3,1	-	4/6/23/26	0/1/1/1
3	NAG	J	2	3	-	3/6/23/26	0/1/1/1
3	BMA	J	3	3	-	2/2/19/22	0/1/1/1
2	NAG	K	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	K	2	2	-	4/6/23/26	0/1/1/1
2	NAG	L	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	L	2	2	-	1/6/23/26	0/1/1/1
2	NAG	M	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	M	2	2	-	3/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	M	2	NAG	C1-C2-N2	3.18	115.45	110.43
2	L	2	NAG	C2-N2-C7	-2.47	119.59	122.90
2	M	2	NAG	O5-C1-C2	-2.41	107.56	111.29
2	E	2	NAG	C2-N2-C7	-2.33	119.77	122.90
2	L	1	NAG	C2-N2-C7	-2.24	119.90	122.90

There are no chirality outliers.

5 of 35 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	L	1	NAG	C8-C7-N2-C2
2	L	1	NAG	O7-C7-N2-C2
2	M	2	NAG	C1-C2-N2-C7
3	J	2	NAG	C3-C2-N2-C7
3	J	2	NAG	C8-C7-N2-C2


There are no ring outliers.


6 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	J	1	NAG	1	0
3	G	2	NAG	1	0
2	M	1	NAG	1	0
3	J	2	NAG	2	0
2	F	1	NAG	1	0
3	G	1	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





















5.6 Ligand geometry

27 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	SO4	C	902	-	4,4,4	0.42	0	6,6,6	0.63	0
4	NAG	A	771(A)	1	14,14,15	0.62	0	17,19,21	0.68	0
4	NAG	C	768(A)	1	14,14,15	0.60	0	17,19,21	0.74	1 (5%)
4	NAG	D	767(A)	1	14,14,15	0.61	0	17,19,21	0.71	1 (5%)
4	NAG	C	771(A)	1	14,14,15	0.64	0	17,19,21	1.11	2 (11%)
5	SO4	D	903	-	4,4,4	0.47	0	6,6,6	0.65	0
6	PHI	A	801	7	11,12,13	0.81	0	10,15,17	0.27	0
6	PHI	B	803	7	11,12,13	0.51	0	10,15,17	0.15	0
5	SO4	A	900	-	4,4,4	0.50	0	6,6,6	0.62	0
4	NAG	D	773(A)	1	14,14,15	0.66	0	17,19,21	0.76	0
4	NAG	C	770(A)	1	14,14,15	0.64	0	17,19,21	0.64	0
4	NAG	A	768(A)	1	14,14,15	0.62	0	17,19,21	0.82	1 (5%)
4	NAG	B	772(A)	1	14,14,15	0.63	0	17,19,21	0.78	0
6	PHI	C	805	7	11,12,13	0.82	0	10,15,17	0.36	0
5	SO4	B	901	-	4,4,4	0.47	0	6,6,6	0.62	0
4	NAG	B	771(A)	1	14,14,15	0.48	0	17,19,21	0.74	0
7	P2Y	C	806	6,1	6,7,7	1.23	1 (16%)	4,8,8	0.99	0
7	P2Y	B	804	6,1	6,7,7	1.42	1 (16%)	4,8,8	0.88	0
4	NAG	B	767(A)	1	14,14,15	0.55	0	17,19,21	0.76	1 (5%)
6	PHI	D	807	7	11,12,13	0.97	0	10,15,17	0.90	0
4	NAG	A	772(A)	1	14,14,15	0.64	0	17,19,21	0.76	0
4	NAG	C	769(A)	1	14,14,15	0.45	0	17,19,21	0.77	1 (5%)
7	P2Y	A	802	6,1	6,7,7	1.18	1 (16%)	4,8,8	1.24	0
4	NAG	C	767(A)	1	14,14,15	0.56	0	17,19,21	0.88	0
4	NAG	B	773(A)	1	14,14,15	0.41	0	17,19,21	1.03	1 (5%)
7	P2Y	D	808	6,1	6,7,7	1.39	1 (16%)	4,8,8	1.10	0
4	NAG	A	767(A)	1	14,14,15	0.59	0	17,19,21	0.68	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	771(A)	1	-	3/6/23/26	0/1/1/1
4	NAG	C	768(A)	1	-	3/6/23/26	0/1/1/1
4	NAG	D	767(A)	1	-	2/6/23/26	0/1/1/1
4	NAG	C	771(A)	1	-	2/6/23/26	0/1/1/1
6	PHI	A	801	7	-	0/5/6/8	0/1/1/1
6	PHI	B	803	7	-	0/5/6/8	0/1/1/1
4	NAG	D	773(A)	1	-	4/6/23/26	0/1/1/1
4	NAG	C	770(A)	1	-	4/6/23/26	0/1/1/1
4	NAG	A	768(A)	1	-	4/6/23/26	0/1/1/1
4	NAG	B	772(A)	1	-	4/6/23/26	0/1/1/1
6	PHI	C	805	7	-	0/5/6/8	0/1/1/1
4	NAG	B	771(A)	1	-	0/6/23/26	0/1/1/1
7	P2Y	C	806	6,1	-	2/2/9/9	0/1/1/1
7	P2Y	B	804	6,1	-	0/2/9/9	0/1/1/1
4	NAG	B	767(A)	1	-	0/6/23/26	0/1/1/1
6	PHI	D	807	7	-	0/5/6/8	0/1/1/1
4	NAG	A	772(A)	1	-	2/6/23/26	0/1/1/1
4	NAG	C	769(A)	1	-	0/6/23/26	0/1/1/1
7	P2Y	A	802	6,1	-	1/2/9/9	0/1/1/1
4	NAG	C	767(A)	1	-	2/6/23/26	0/1/1/1
4	NAG	B	773(A)	1	-	2/6/23/26	0/1/1/1
7	P2Y	D	808	6,1	-	2/2/9/9	0/1/1/1
4	NAG	A	767(A)	1	-	4/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	804	P2Y	C-NA	-3.19	1.32	1.48
7	D	808	P2Y	C-NA	-2.93	1.34	1.48
7	A	802	P2Y	C-NA	-2.65	1.35	1.48
7	C	806	P2Y	C-NA	-2.47	1.36	1.48

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	773(A)	NAG	C2-N2-C7	-3.12	118.71	122.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	769(A)	NAG	C2-N2-C7	-2.31	119.81	122.90
4	A	768(A)	NAG	C2-N2-C7	-2.21	119.94	122.90
4	B	767(A)	NAG	C2-N2-C7	-2.13	120.04	122.90
4	D	767(A)	NAG	C2-N2-C7	-2.11	120.07	122.90

There are no chirality outliers.

5 of 41 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	768(A)	NAG	C8-C7-N2-C2
4	A	768(A)	NAG	O7-C7-N2-C2
4	A	771(A)	NAG	C8-C7-N2-C2
4	A	771(A)	NAG	O7-C7-N2-C2
4	B	772(A)	NAG	C8-C7-N2-C2

There are no ring outliers.

14 monomers are involved in 36 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	767(A)	NAG	1	0
6	A	801	PHI	3	0
6	B	803	PHI	4	0
4	B	772(A)	NAG	2	0
6	C	805	PHI	7	0
7	C	806	P2Y	1	0
7	B	804	P2Y	3	0
4	B	767(A)	NAG	2	0
6	D	807	PHI	7	0
7	A	802	P2Y	2	0
4	C	767(A)	NAG	2	0
4	B	773(A)	NAG	1	0
7	D	808	P2Y	2	0
4	A	767(A)	NAG	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	724/728 (99%)	-0.30	7 (0%) 79 75	7, 27, 58, 95	11 (1%)
1	B	728/728 (100%)	-0.42	2 (0%) 90 89	6, 25, 51, 67	12 (1%)
1	C	726/728 (99%)	-0.35	13 (1%) 67 60	6, 24, 54, 102	12 (1%)
1	D	728/728 (100%)	-0.24	3 (0%) 88 87	8, 30, 57, 80	10 (1%)
All	All	2906/2912 (99%)	-0.33	25 (0%) 81 76	6, 26, 55, 102	45 (1%)

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	94	THR	4.2
1	D	82	GLU	3.7
1	D	83	TYR	3.6
1	A	101	SER	3.5
1	C	98	LEU	3.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	BMA	J	3	11/12	0.74	0.13	53,56,57,58	0
2	NAG	K	2	14/15	0.75	0.14	47,55,59,61	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	F	2	14/15	0.75	0.14	47,51,54,55	0
2	NAG	E	2	14/15	0.77	0.13	48,52,54,55	0
2	NAG	H	2	14/15	0.77	0.14	46,52,58,60	0
3	BMA	G	3	11/12	0.80	0.10	53,54,56,56	0
2	NAG	L	2	14/15	0.81	0.12	60,61,62,62	0
3	NAG	J	2	14/15	0.83	0.13	58,60,66,67	0
2	NAG	M	2	14/15	0.84	0.11	50,52,54,54	0
2	NAG	I	2	14/15	0.87	0.09	28,33,37,40	0
2	NAG	M	1	14/15	0.87	0.10	33,36,41,46	0
2	NAG	L	1	14/15	0.87	0.12	50,54,55,59	0
3	NAG	G	2	14/15	0.88	0.10	46,47,49,51	0
3	NAG	G	1	14/15	0.89	0.11	45,47,49,49	0
3	NAG	J	1	14/15	0.89	0.11	54,57,58,59	0
2	NAG	K	1	14/15	0.93	0.08	27,34,40,46	0
2	NAG	F	1	14/15	0.93	0.08	30,32,39,41	0
2	NAG	H	1	14/15	0.93	0.08	26,34,38,44	0
2	NAG	E	1	14/15	0.94	0.08	26,28,34,41	0
2	NAG	I	1	14/15	0.95	0.06	21,28,34,36	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain F:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain H:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around Chain I:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain K:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain L:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain M:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain G:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	D	767(A)	14/15	0.41	0.18	79,82,83,83	0
4	NAG	C	768(A)	14/15	0.54	0.15	78,84,87,87	0
4	NAG	A	767(A)	14/15	0.71	0.14	82,82,83,83	0
4	NAG	B	767(A)	14/15	0.75	0.12	64,66,70,71	0
4	NAG	B	772(A)	14/15	0.76	0.13	43,46,53,54	0
4	NAG	D	773(A)	14/15	0.79	0.14	46,49,54,55	0
4	NAG	C	771(A)	14/15	0.81	0.13	35,38,43,43	0
4	NAG	A	768(A)	14/15	0.82	0.11	65,66,68,68	0
4	NAG	C	767(A)	14/15	0.84	0.12	55,56,58,58	0
4	NAG	A	771(A)	14/15	0.85	0.12	38,39,40,42	3
4	NAG	C	770(A)	14/15	0.86	0.11	35,38,42,43	0
4	NAG	A	772(A)	14/15	0.88	0.10	35,38,40,40	0
6	PHI	A	801	12/13	0.88	0.20	46,65,80,97	0
6	PHI	D	807	12/13	0.89	0.19	50,67,81,99	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NAG	B	773(A)	14/15	0.90	0.10	37,40,44,45	0
4	NAG	C	769(A)	14/15	0.91	0.09	28,31,32,35	0
6	PHI	B	803	12/13	0.92	0.16	42,59,72,88	0
6	PHI	C	805	12/13	0.92	0.17	44,59,72,89	0
4	NAG	B	771(A)	14/15	0.92	0.08	19,21,23,26	0
7	P2Y	B	804	7/7	0.93	0.13	36,37,37,38	0
7	P2Y	D	808	7/7	0.94	0.14	42,43,45,47	0
5	SO4	A	900	5/5	0.95	0.09	50,50,51,52	0
5	SO4	D	903	5/5	0.96	0.07	47,48,48,49	0
7	P2Y	C	806	7/7	0.96	0.09	33,35,38,40	0
7	P2Y	A	802	7/7	0.96	0.11	35,39,41,42	0
5	SO4	B	901	5/5	0.98	0.07	42,43,43,44	0
5	SO4	C	902	5/5	0.99	0.05	35,37,37,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.