



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:50 PM UTC

PDB ID : 2OR2 / pdb_00002or2
Title : Structure of the W47A/W242A Mutant of Bacterial Phosphatidylinositol-Specific Phospholipase C
Authors : Shao, C.; Shi, X.; Wehbi, H.; Zambonelli, C.; Head, J.F.; Seaton, B.A.; Roberts, M.F.
Deposited on : 2007-02-01
Resolution : 1.84 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1296 (1.84-1.84)
Clashscore	190562	1329 (1.84-1.84)
Ramachandran outliers	187476	1318 (1.84-1.84)
Sidechain outliers	187428	1318 (1.84-1.84)
RSRZ outliers	180081	1296 (1.84-1.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	296	8% 81% 19%
1	B	296	8% 82% 17% .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5382 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1-phosphatidylinositol phosphodiesterase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	296	Total	C	N	O	S	0	0	0
			2403	1526	400	470	7			
1	B	296	Total	C	N	O	S	0	0	0
			2403	1526	400	470	7			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	47	ALA	TRP	engineered mutation	UNP P08954
A	242	ALA	TRP	engineered mutation	UNP P08954
B	47	ALA	TRP	engineered mutation	UNP P08954
B	242	ALA	TRP	engineered mutation	UNP P08954


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	274	Total	O	0	0
			274	274		
2	B	302	Total	O	0	0
			302	302		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 1-phosphatidylinositol phosphodiesterase



- Molecule 1: 1-phosphatidylinositol phosphodiesterase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	64.42Å 70.03Å 154.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.84 50.00 – 1.84	Depositor EDS
% Data completeness (in resolution range)	95.6 (50.00-1.84) 95.6 (50.00-1.84)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.54 (at 1.84Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.186 , 0.215 0.186 , 0.215	Depositor DCC
R_{free} test set	4952 reflections (8.18%)	wwPDB-VP
Wilson B-factor (Å ²)	17.5	Xtriage
Anisotropy	0.795	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 48.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5382	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.33	0/2461	0.90	6/3336 (0.2%)
1	B	0.35	0/2461	0.87	3/3336 (0.1%)
All	All	0.34	0/4922	0.89	9/6672 (0.1%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	275	TYR	N-CA-C	8.43	121.56	111.02
1	B	183	THR	N-CA-C	-6.69	97.61	108.52
1	B	34	SER	N-CA-C	6.55	119.25	111.33
1	A	37	PHE	N-CA-C	6.25	118.90	111.71
1	A	192	VAL	N-CA-C	5.76	116.36	107.78
1	A	139	ASP	CA-C-N	5.47	125.14	119.56
1	A	139	ASP	C-N-CA	5.47	125.14	119.56
1	A	183	THR	N-CA-C	-5.21	100.03	108.52
1	B	37	PHE	N-CA-C	5.08	118.74	112.54

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2403	0	2323	49	0
1	B	2403	0	2323	47	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	274	0	0	6	0
2	B	302	0	0	3	0
All	All	5382	0	4646	93	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:194:VAL:HG22	1:A:228:LEU:HB3	1.44	0.98
1:B:59:MET:HE1	1:B:98:ALA:HA	1.54	0.89
1:B:73:THR:HB	1:B:121:MET:HE2	1.57	0.87
1:B:74:ASP:H	1:B:121:MET:HE3	1.43	0.83
1:A:190:GLN:HE21	1:A:191:ASN:ND2	1.81	0.78
1:B:190:GLN:CD	1:B:190:GLN:H	1.93	0.77
1:B:215:MET:HE1	1:B:260:ILE:HD11	1.70	0.74
1:A:190:GLN:H	1:A:190:GLN:CD	1.96	0.74
1:A:218:THR:HG22	1:A:267:ARG:HG2	1.70	0.74
1:A:129:SER:O	1:A:133:GLU:HG3	1.89	0.72
1:A:267:ARG:HD2	2:A:339:HOH:O	1.88	0.72
1:A:69:ARG:HB3	1:A:117:GLU:HB2	1.70	0.71
1:A:76:ASN:HD21	1:A:125:GLU:HG2	1.57	0.69
1:B:267:ARG:HD2	2:B:540:HOH:O	1.92	0.67
1:A:1:ALA:HA	1:A:16:PRO:HA	1.78	0.65
1:B:74:ASP:H	1:B:121:MET:CE	2.11	0.64
1:A:4:VAL:HA	1:A:296:ILE:CD1	2.28	0.63
1:B:73:THR:HA	1:B:121:MET:HG3	1.80	0.62
1:A:248:TYR:OH	1:B:42:PRO:HB2	2.00	0.62
1:A:216:ASP:HA	1:A:219:MET:HE3	1.82	0.61
1:B:190:GLN:H	1:B:190:GLN:NE2	1.98	0.61
1:B:133:GLU:HA	1:B:137:PHE:HB3	1.83	0.60
1:B:244:SER:N	1:B:245:PRO:HD2	2.17	0.60
1:B:69:ARG:HB3	1:B:117:GLU:HB2	1.84	0.59
1:B:92:HIS:HD2	2:B:576:HOH:O	1.85	0.59
1:B:218:THR:HG22	1:B:267:ARG:HG2	1.83	0.59
1:A:73:THR:HG22	1:A:121:MET:HE3	1.85	0.58
1:B:253:ASN:HB2	1:B:254:PRO:HD3	1.86	0.58
1:B:73:THR:CB	1:B:121:MET:HE2	2.29	0.58
1:B:41:ASN:OD1	1:B:43:ILE:HG22	2.04	0.57
1:B:129:SER:O	1:B:133:GLU:HG3	2.05	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:243:ASN:C	1:B:245:PRO:HD2	2.31	0.54
1:A:47:ALA:HB2	1:B:241:ALA:O	2.08	0.54
1:B:218:THR:CG2	1:B:267:ARG:HG2	2.38	0.53
1:A:76:ASN:ND2	1:A:125:GLU:HG2	2.22	0.53
1:A:218:THR:CG2	1:A:267:ARG:HG2	2.36	0.52
1:A:4:VAL:HA	1:A:296:ILE:HD13	1.91	0.52
1:A:86:TYR:CE2	1:A:88:TYR:HA	2.44	0.52
1:A:180:ASP:O	1:A:182:GLU:HG3	2.10	0.52
1:A:254:PRO:HG3	1:A:280:TRP:HB3	1.92	0.52
1:B:133:GLU:HA	1:B:137:PHE:CB	2.40	0.51
1:A:1:ALA:CA	1:A:16:PRO:HA	2.41	0.51
1:B:175:ASN:ND2	2:B:318:HOH:O	2.42	0.51
1:A:253:ASN:HB2	1:A:254:PRO:HD3	1.92	0.50
1:B:254:PRO:HG3	1:B:280:TRP:HB3	1.93	0.50
1:A:219:MET:SD	1:A:265:PRO:HB3	2.51	0.50
1:A:188:VAL:CG2	1:A:194:VAL:HG21	2.42	0.49
1:A:121:MET:HE2	2:A:426:HOH:O	2.12	0.49
1:A:1:ALA:HA	1:A:16:PRO:CA	2.44	0.48
1:A:4:VAL:HA	1:A:296:ILE:HD12	1.95	0.48
1:A:243:ASN:C	1:A:245:PRO:HD2	2.39	0.48
1:B:71:ARG:CZ	1:B:121:MET:HG2	2.44	0.48
1:A:10:TRP:CH2	1:A:286:GLN:HG3	2.49	0.47
1:A:244:SER:N	1:A:245:PRO:HD2	2.28	0.47
1:B:236:SER:O	1:B:237:SER:C	2.56	0.47
1:A:82:HIS:HA	2:A:451:HOH:O	2.13	0.47
1:B:1:ALA:HA	1:B:6:GLU:OE1	2.15	0.47
1:B:190:GLN:NE2	1:B:190:GLN:N	2.63	0.47
1:B:74:ASP:N	1:B:121:MET:HE3	2.22	0.47
1:B:59:MET:HE3	1:B:110:ILE:CD1	2.46	0.46
1:A:177:TYR:CE2	1:A:179:PRO:HD3	2.51	0.46
1:A:188:VAL:HG23	1:A:194:VAL:CG2	2.46	0.45
1:A:1:ALA:CB	1:A:16:PRO:HA	2.47	0.45
1:A:188:VAL:HG23	1:A:194:VAL:HG23	1.96	0.45
1:A:7:LEU:O	1:A:10:TRP:CZ3	2.70	0.45
1:A:133:GLU:HA	1:A:137:PHE:HB3	1.97	0.44
1:B:59:MET:HE2	1:B:101:PHE:CD2	2.53	0.44
1:B:48:GLY:C	1:B:49:MET:HE2	2.42	0.44
1:B:86:TYR:CZ	1:B:88:TYR:HA	2.53	0.44
1:B:237:SER:HB2	1:B:248:TYR:HE2	1.82	0.44
1:A:215:MET:HG2	1:A:219:MET:HE2	2.00	0.44
1:B:259:ASP:OD1	1:B:263:LYS:HE3	2.18	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:185:THR:HA	1:A:194:VAL:O	2.18	0.43
1:A:239:GLY:H	1:B:45:GLN:HG2	1.82	0.43
1:A:242:ALA:O	1:A:244:SER:N	2.51	0.43
1:A:189:ASN:ND2	2:A:423:HOH:O	2.51	0.43
1:B:244:SER:N	1:B:245:PRO:CD	2.81	0.43
1:A:286:GLN:OE1	1:A:290:ARG:NH2	2.52	0.43
1:B:237:SER:HB2	1:B:248:TYR:CE2	2.54	0.43
1:A:161:LEU:C	1:A:161:LEU:HD23	2.44	0.42
1:A:7:LEU:O	1:A:10:TRP:HZ3	2.01	0.42
1:A:10:TRP:CZ2	1:A:286:GLN:HG3	2.54	0.42
2:A:509:HOH:O	1:B:241:ALA:HB1	2.20	0.42
1:A:54:ASP:HB2	2:A:345:HOH:O	2.19	0.42
1:B:59:MET:HE2	1:B:110:ILE:HD11	2.02	0.42
1:B:183:THR:HA	1:B:196:VAL:O	2.19	0.42
1:A:194:VAL:CG2	1:A:228:LEU:HD23	2.50	0.41
1:B:36:THR:HB	1:B:49:MET:SD	2.59	0.41
1:B:180:ASP:O	1:B:182:GLU:HG3	2.20	0.41
1:B:28:ILE:HA	1:B:29:PRO:HD3	1.89	0.41
1:B:202:VAL:O	1:B:237:SER:OG	2.38	0.41
1:B:190:GLN:CD	1:B:190:GLN:N	2.72	0.41
1:A:188:VAL:CG2	1:A:194:VAL:CG2	2.99	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	294/296 (99%)	282 (96%)	11 (4%)	1 (0%)	36	25
1	B	294/296 (99%)	279 (95%)	14 (5%)	1 (0%)	36	25
All	All	588/592 (99%)	561 (95%)	25 (4%)	2 (0%)	36	25

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	237	SER
1	A	244	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/267 (100%)	265 (99%)	2 (1%)	76	68
1	B	267/267 (100%)	266 (100%)	1 (0%)	84	78
All	All	534/534 (100%)	531 (99%)	3 (1%)	78	72

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	81	HIS
1	A	223	GLU
1	B	190	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	HIS
1	A	191	ASN
1	A	264	ASN
1	A	273	GLN
1	B	45	GLN
1	B	92	HIS
1	B	174	ASN
1	B	190	GLN
1	B	253	ASN
1	B	258	ASN
1	B	286	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	296/296 (100%)	0.28	24 (8%) 18 19	11, 21, 41, 64	0
1	B	296/296 (100%)	0.14	25 (8%) 17 18	10, 18, 45, 64	0
All	All	592/592 (100%)	0.21	49 (8%) 17 18	10, 19, 44, 64	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	ALA	9.3
1	B	1	ALA	8.3
1	B	241	ALA	7.3
1	A	238	GLY	6.7
1	B	242	ALA	6.7
1	A	242	ALA	6.6
1	B	237	SER	6.0
1	B	189	ASN	5.4
1	A	241	ALA	4.8
1	A	10	TRP	4.7
1	B	243	ASN	4.4
1	B	238	GLY	4.4
1	A	243	ASN	4.3
1	A	237	SER	4.2
1	A	239	GLY	4.1
1	B	240	THR	4.1
1	A	240	THR	3.7
1	B	190	GLN	3.6
1	B	239	GLY	3.3
1	A	5	ASN	3.2
1	A	188	VAL	3.1
1	B	2	SER	3.0
1	A	191	ASN	3.0
1	A	189	ASN	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	43	ILE	2.9
1	B	40	GLN	2.9
1	A	2	SER	2.8
1	A	219	MET	2.7
1	A	168	ASN	2.6
1	B	42	PRO	2.6
1	A	262	GLN	2.6
1	B	124	ALA	2.5
1	B	5	ASN	2.4
1	B	121	MET	2.4
1	A	203	ASN	2.4
1	B	236	SER	2.4
1	B	85	LEU	2.4
1	A	125	GLU	2.4
1	B	123	GLY	2.4
1	B	191	ASN	2.3
1	B	41	ASN	2.3
1	B	125	GLU	2.3
1	B	4	VAL	2.3
1	B	48	GLY	2.2
1	A	119	GLU	2.2
1	A	177	TYR	2.1
1	A	221	ASN	2.1
1	A	206	GLU	2.0
1	A	43	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.