



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 03:20 PM UTC

PDB ID : 3ORX / pdb_00003orx
Title : PDK1 mutant bound to allosteric disulfide fragment inhibitor 1F8
Authors : Sadowsky, J.D.; Wells, J.A.
Deposited on : 2010-09-08
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	5% 66% 20% • 10%
1	B	316	5% 64% 21% • 11%
1	C	316	6% 67% 19% • 10%
1	D	316	5% 68% 18% • 10%
1	E	316	10% 63% 19% • 13%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	316	13% 64% 17% 6% 13%
1	G	316	11% 60% 21% 6% 13%
1	H	316	13% 65% 18% • 13%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 19043 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3-phosphoinositide-dependent protein kinase 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	284	Total	C	N	O	P	S	0	0	0
			2271	1476	374	412	1	8			
1	B	281	Total	C	N	O	P	S	0	0	0
			2262	1470	368	415	1	8			
1	C	284	Total	C	N	O	P	S	0	0	0
			2264	1469	369	417	1	8			
1	D	283	Total	C	N	O	P	S	0	0	0
			2256	1468	370	409	1	8			
1	E	274	Total	C	N	O	S		0	0	0
			2163	1415	354	386	8				
1	F	275	Total	C	N	O	P	S	0	0	0
			2132	1391	348	384	1	8			
1	G	275	Total	C	N	O	P	S	0	0	0
			2155	1406	352	388	1	8			
1	H	275	Total	C	N	O	P	S	0	0	0
			2114	1379	342	384	1	8			

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	44	GLY	-	expression tag	UNP O15530
A	45	ALA	-	expression tag	UNP O15530
A	46	MET	-	expression tag	UNP O15530
A	47	ASP	-	expression tag	UNP O15530
A	48	PRO	-	expression tag	UNP O15530
A	49	GLU	-	expression tag	UNP O15530
A	50	PHE	-	expression tag	UNP O15530
A	148	CYS	THR	engineered mutation	UNP O15530
B	44	GLY	-	expression tag	UNP O15530
B	45	ALA	-	expression tag	UNP O15530
B	46	MET	-	expression tag	UNP O15530
B	47	ASP	-	expression tag	UNP O15530
B	48	PRO	-	expression tag	UNP O15530

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	49	GLU	-	expression tag	UNP O15530
B	50	PHE	-	expression tag	UNP O15530
B	148	CYS	THR	engineered mutation	UNP O15530
C	44	GLY	-	expression tag	UNP O15530
C	45	ALA	-	expression tag	UNP O15530
C	46	MET	-	expression tag	UNP O15530
C	47	ASP	-	expression tag	UNP O15530
C	48	PRO	-	expression tag	UNP O15530
C	49	GLU	-	expression tag	UNP O15530
C	50	PHE	-	expression tag	UNP O15530
C	148	CYS	THR	engineered mutation	UNP O15530
D	44	GLY	-	expression tag	UNP O15530
D	45	ALA	-	expression tag	UNP O15530
D	46	MET	-	expression tag	UNP O15530
D	47	ASP	-	expression tag	UNP O15530
D	48	PRO	-	expression tag	UNP O15530
D	49	GLU	-	expression tag	UNP O15530
D	50	PHE	-	expression tag	UNP O15530
D	148	CYS	THR	engineered mutation	UNP O15530
E	44	GLY	-	expression tag	UNP O15530
E	45	ALA	-	expression tag	UNP O15530
E	46	MET	-	expression tag	UNP O15530
E	47	ASP	-	expression tag	UNP O15530
E	48	PRO	-	expression tag	UNP O15530
E	49	GLU	-	expression tag	UNP O15530
E	50	PHE	-	expression tag	UNP O15530
E	148	CYS	THR	engineered mutation	UNP O15530
F	44	GLY	-	expression tag	UNP O15530
F	45	ALA	-	expression tag	UNP O15530
F	46	MET	-	expression tag	UNP O15530
F	47	ASP	-	expression tag	UNP O15530
F	48	PRO	-	expression tag	UNP O15530
F	49	GLU	-	expression tag	UNP O15530
F	50	PHE	-	expression tag	UNP O15530
F	148	CYS	THR	engineered mutation	UNP O15530
G	44	GLY	-	expression tag	UNP O15530
G	45	ALA	-	expression tag	UNP O15530
G	46	MET	-	expression tag	UNP O15530
G	47	ASP	-	expression tag	UNP O15530
G	48	PRO	-	expression tag	UNP O15530
G	49	GLU	-	expression tag	UNP O15530
G	50	PHE	-	expression tag	UNP O15530

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	148	CYS	THR	engineered mutation	UNP O15530
H	44	GLY	-	expression tag	UNP O15530
H	45	ALA	-	expression tag	UNP O15530
H	46	MET	-	expression tag	UNP O15530
H	47	ASP	-	expression tag	UNP O15530
H	48	PRO	-	expression tag	UNP O15530
H	49	GLU	-	expression tag	UNP O15530
H	50	PHE	-	expression tag	UNP O15530
H	148	CYS	THR	engineered mutation	UNP O15530

- Molecule 2 is 2-methyl-N-(2-sulfanylethyl)-1-benzofuran-3-carboxamide (CCD ID: 1F8) (formula: C₁₂H₁₃NO₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			16	12	1	2	1		
2	B	1	Total	C	N	O	S	0	0
			16	12	1	2	1		
2	C	1	Total	C	N	O	S	0	0
			16	12	1	2	1		
2	D	1	Total	C	N	O	S	0	0
			16	12	1	2	1		
2	E	1	Total	C	N	O	S	0	0
			16	12	1	2	1		
2	F	1	Total	C	N	O	S	0	0
			16	12	1	2	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	G	1	Total	C	N	O	S	0	0
			16	12	1	2	1		
2	H	1	Total	C	N	O	S	0	0
			16	12	1	2	1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		
3	B	1	Total	Cl	0	0
			1	1		
3	C	1	Total	Cl	0	0
			1	1		
3	D	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	204	Total	O	0	0
			204	204		
4	B	213	Total	O	0	0
			213	213		
4	C	194	Total	O	0	0
			194	194		
4	D	187	Total	O	0	0
			187	187		
4	E	120	Total	O	0	0
			120	120		
4	F	121	Total	O	0	0
			121	121		
4	G	121	Total	O	0	0
			121	121		
4	H	134	Total	O	0	0
			134	134		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3-phosphoinositide-dependent protein kinase 1




- Molecule 1: 3-phosphoinositide-dependent protein kinase 1




- Molecule 1: 3-phosphoinositide-dependent protein kinase 1







• Molecule 1: 3-phosphoinositide-dependent protein kinase 1



• Molecule 1: 3-phosphoinositide-dependent protein kinase 1




• Molecule 1: 3-phosphoinositide-dependent protein kinase 1



T359


• Molecule 1: 3-phosphoinositide-dependent protein kinase 1

Chain G: 11% 60% 21% 6% 13%



• Molecule 1: 3-phosphoinositide-dependent protein kinase 1

Chain H: 13% 65% 18% 13%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	40.29Å 115.87Å 145.72Å 91.76° 89.99° 95.41°	Depositor
Resolution (Å)	45.25 – 2.20 45.25 – 2.20	Depositor EDS
% Data completeness (in resolution range)	96.8 (45.25-2.20) 96.8 (45.25-2.20)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.96 (at 2.20Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 2010_01_09_2330)	Depositor
R, R_{free}	0.208 , 0.243 0.204 , 0.239	Depositor DCC
R_{free} test set	6446 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	29.7	Xtriage
Anisotropy	0.652	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 59.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.073 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	19043	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, SEP, 1F8

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	6/2315 (0.3%)	0.88	3/3125 (0.1%)
1	B	0.70	3/2306 (0.1%)	0.88	2/3112 (0.1%)
1	C	0.75	4/2308 (0.2%)	0.89	3/3120 (0.1%)
1	D	0.68	2/2300 (0.1%)	0.87	2/3106 (0.1%)
1	E	0.96	4/2217 (0.2%)	0.97	8/3003 (0.3%)
1	F	0.87	3/2175 (0.1%)	0.92	7/2950 (0.2%)
1	G	1.03	6/2199 (0.3%)	0.97	4/2980 (0.1%)
1	H	0.88	1/2158 (0.0%)	0.93	5/2932 (0.2%)
All	All	0.84	29/17978 (0.2%)	0.91	34/24328 (0.1%)

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	203	HIS	C-O	-7.23	1.15	1.24
1	C	205	ASP	C-O	-7.08	1.15	1.23
1	B	205	ASP	C-O	-7.05	1.16	1.23
1	G	280	PRO	C-O	-6.62	1.19	1.24
1	A	205	ASP	C-O	-6.51	1.16	1.24

The worst 5 of 34 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	153	GLU	N-CA-C	8.53	120.66	111.36
1	H	153	GLU	N-CA-C	7.52	119.56	111.36
1	G	203	HIS	N-CA-C	7.50	119.24	111.14
1	E	154	LYS	N-CA-C	7.29	120.49	109.41
1	G	173	LYS	N-CA-C	7.22	118.94	111.14

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2271	0	2250	56	0
1	B	2262	0	2245	56	0
1	C	2264	0	2218	58	0
1	D	2256	0	2233	46	0
1	E	2163	0	2101	63	0
1	F	2132	0	2040	58	0
1	G	2155	0	2069	71	0
1	H	2114	0	1982	58	0
2	A	16	0	12	3	0
2	B	16	0	12	3	0
2	C	16	0	12	3	0
2	D	16	0	12	3	0
2	E	16	0	12	2	0
2	F	16	0	12	2	0
2	G	16	0	12	3	0
2	H	16	0	12	2	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	204	0	0	4	0
4	B	213	0	0	9	0
4	C	194	0	0	11	0
4	D	187	0	0	5	0
4	E	120	0	0	1	0
4	F	121	0	0	3	0
4	G	121	0	0	2	0
4	H	134	0	0	3	0
All	All	19043	0	17234	445	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 445 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:230:LEU:HD21	1:A:235:LYS:HA	1.28	1.12
1:A:323:LYS:NZ	1:G:310:ARG:HH21	1.47	1.10
1:F:253:LEU:HD23	1:F:253:LEU:N	1.69	1.04
1:E:253:LEU:N	1:E:253:LEU:HD23	1.74	1.01
1:A:323:LYS:HZ1	1:G:310:ARG:HH21	1.00	1.00

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	279/316 (88%)	274 (98%)	5 (2%)	0	100	100
1	B	276/316 (87%)	271 (98%)	5 (2%)	0	100	100
1	C	279/316 (88%)	273 (98%)	6 (2%)	0	100	100
1	D	278/316 (88%)	272 (98%)	5 (2%)	1 (0%)	30	34
1	E	270/316 (85%)	260 (96%)	10 (4%)	0	100	100
1	F	271/316 (86%)	260 (96%)	11 (4%)	0	100	100
1	G	271/316 (86%)	261 (96%)	10 (4%)	0	100	100
1	H	271/316 (86%)	261 (96%)	10 (4%)	0	100	100
All	All	2195/2528 (87%)	2132 (97%)	62 (3%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	244	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/273 (88%)	219 (92%)	20 (8%)	10	11
1	B	241/273 (88%)	222 (92%)	19 (8%)	11	13
1	C	238/273 (87%)	220 (92%)	18 (8%)	12	14
1	D	237/273 (87%)	219 (92%)	18 (8%)	12	14
1	E	220/273 (81%)	201 (91%)	19 (9%)	10	10
1	F	211/273 (77%)	183 (87%)	28 (13%)	4	3
1	G	215/273 (79%)	190 (88%)	25 (12%)	5	5
1	H	205/273 (75%)	186 (91%)	19 (9%)	8	9
All	All	1806/2184 (83%)	1640 (91%)	166 (9%)	8	9

5 of 166 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	257	LYS
1	G	303	GLU
1	F	345	VAL
1	G	173	LYS
1	H	106	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 26 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	349	ASN
1	D	286	ASN
1	H	247	GLN
1	D	247	GLN
1	D	349	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

7 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	SEP	B	241	1	8,9,10	1.62	1 (12%)	7,12,14	1.34	2 (28%)
1	SEP	F	241	1	8,9,10	1.66	1 (12%)	7,12,14	1.16	1 (14%)
1	SEP	D	241	1	8,9,10	1.70	1 (12%)	7,12,14	1.38	2 (28%)
1	SEP	A	241	1	8,9,10	1.65	1 (12%)	7,12,14	1.47	2 (28%)
1	SEP	H	241	1	8,9,10	1.68	1 (12%)	7,12,14	1.07	0
1	SEP	C	241	1	8,9,10	1.06	1 (12%)	7,12,14	1.91	2 (28%)
1	SEP	G	241	1	8,9,10	1.69	1 (12%)	7,12,14	1.39	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	B	241	1	-	0/6/8/10	-
1	SEP	F	241	1	-	5/6/8/10	-
1	SEP	D	241	1	-	0/6/8/10	-
1	SEP	A	241	1	-	0/6/8/10	-
1	SEP	H	241	1	-	4/6/8/10	-
1	SEP	C	241	1	-	2/6/8/10	-
1	SEP	G	241	1	-	2/6/8/10	-

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	241	SEP	P-O1P	3.74	1.62	1.50
1	H	241	SEP	P-O1P	3.71	1.62	1.50
1	F	241	SEP	P-O1P	3.66	1.61	1.50
1	A	241	SEP	P-O1P	3.65	1.61	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	241	SEP	P-O1P	3.64	1.61	1.50

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	241	SEP	OG-CB-CA	3.65	111.70	108.14
1	G	241	SEP	OG-CB-CA	2.96	111.03	108.14
1	C	241	SEP	O2P-P-OG	2.72	113.77	106.67
1	A	241	SEP	OG-CB-CA	2.66	110.73	108.14
1	B	241	SEP	OG-CB-CA	2.57	110.65	108.14

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	C	241	SEP	CB-OG-P-O3P
1	F	241	SEP	CB-OG-P-O2P
1	F	241	SEP	CB-OG-P-O3P
1	G	241	SEP	N-CA-CB-OG
1	G	241	SEP	C-CA-CB-OG

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	241	SEP	1	0
1	D	241	SEP	1	0
1	A	241	SEP	1	0
1	C	241	SEP	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 4 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	1F8	G	1	1	16,17,17	3.09	2 (12%)	20,23,23	2.64	8 (40%)
2	1F8	C	1	1	16,17,17	3.13	2 (12%)	20,23,23	2.63	9 (45%)
2	1F8	F	1	1	16,17,17	3.11	2 (12%)	20,23,23	2.69	8 (40%)
2	1F8	H	1	1	16,17,17	3.10	2 (12%)	20,23,23	2.65	8 (40%)
2	1F8	B	1	1	16,17,17	3.06	2 (12%)	20,23,23	2.67	9 (45%)
2	1F8	E	1	1	16,17,17	3.12	2 (12%)	20,23,23	2.63	8 (40%)
2	1F8	A	1	1	16,17,17	3.16	2 (12%)	20,23,23	2.65	9 (45%)
2	1F8	D	1	1	16,17,17	3.11	2 (12%)	20,23,23	2.63	9 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	1F8	G	1	1	-	1/8/8/8	0/2/2/2
2	1F8	C	1	1	-	1/8/8/8	0/2/2/2
2	1F8	F	1	1	-	1/8/8/8	0/2/2/2
2	1F8	H	1	1	-	1/8/8/8	0/2/2/2
2	1F8	B	1	1	-	1/8/8/8	0/2/2/2
2	1F8	E	1	1	-	1/8/8/8	0/2/2/2
2	1F8	A	1	1	-	1/8/8/8	0/2/2/2
2	1F8	D	1	1	-	1/8/8/8	0/2/2/2

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1	1F8	C20-C24	12.07	1.49	1.36
2	C	1	1F8	C20-C24	11.93	1.49	1.36
2	E	1	1F8	C20-C24	11.92	1.48	1.36
2	F	1	1F8	C20-C24	11.90	1.48	1.36
2	D	1	1F8	C20-C24	11.87	1.48	1.36

The worst 5 of 68 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	1	1F8	O23-C24-C20	-7.41	105.20	111.22
2	E	1	1F8	O23-C24-C20	-7.32	105.28	111.22
2	G	1	1F8	O23-C24-C20	-7.25	105.33	111.22
2	H	1	1F8	O23-C24-C20	-7.20	105.37	111.22
2	B	1	1F8	O23-C24-C20	-6.98	105.55	111.22

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	1	1F8	SD-C15-C16-N17
2	E	1	1F8	C15-C16-N17-C18
2	F	1	1F8	C15-C16-N17-C18
2	G	1	1F8	C15-C16-N17-C18
2	H	1	1F8	C15-C16-N17-C18

There are no ring outliers.

8 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	1	1F8	3	0
2	C	1	1F8	3	0
2	F	1	1F8	2	0
2	H	1	1F8	2	0
2	B	1	1F8	3	0
2	E	1	1F8	2	0
2	A	1	1F8	3	0
2	D	1	1F8	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	283/316 (89%)	0.29	15 (5%) 32 29	16, 32, 69, 100	0
1	B	280/316 (88%)	0.29	17 (6%) 27 24	17, 32, 70, 88	0
1	C	283/316 (89%)	0.34	19 (6%) 24 21	16, 32, 73, 98	0
1	D	282/316 (89%)	0.31	16 (5%) 29 26	17, 33, 72, 96	0
1	E	274/316 (86%)	0.76	33 (12%) 9 6	17, 41, 85, 109	0
1	F	274/316 (86%)	0.78	40 (14%) 6 4	18, 41, 86, 108	0
1	G	274/316 (86%)	0.77	34 (12%) 8 6	17, 42, 85, 109	0
1	H	274/316 (86%)	0.77	42 (15%) 5 4	17, 42, 86, 109	0
All	All	2224/2528 (87%)	0.54	216 (9%) 13 10	16, 37, 82, 109	0

The worst 5 of 216 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	124	VAL	5.1
1	F	82	PHE	5.1
1	A	242	PHE	4.9
1	F	125	PRO	4.8
1	E	243	VAL	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	H	241	10/11	0.59	0.17	64,81,89,93	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	SEP	G	241	10/11	0.72	0.19	66,78,100,109	0
1	SEP	F	241	10/11	0.72	0.21	70,81,94,96	0
1	SEP	D	241	10/11	0.85	0.14	45,60,82,85	0
1	SEP	C	241	10/11	0.87	0.14	48,64,81,86	0
1	SEP	B	241	10/11	0.89	0.12	46,61,81,86	0
1	SEP	A	241	10/11	0.90	0.13	45,61,78,84	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	1F8	F	1	16/16	0.63	0.23	87,97,109,111	0
2	1F8	H	1	16/16	0.64	0.22	87,98,109,112	0
2	1F8	E	1	16/16	0.65	0.23	87,96,108,111	0
2	1F8	G	1	16/16	0.67	0.23	87,98,108,110	0
2	1F8	A	1	16/16	0.67	0.23	72,88,110,120	0
2	1F8	C	1	16/16	0.70	0.22	71,88,109,120	0
2	1F8	B	1	16/16	0.72	0.21	72,89,110,121	0
2	1F8	D	1	16/16	0.73	0.19	73,88,110,120	0
3	CL	B	5	1/1	0.96	0.08	44,44,44,44	0
3	CL	D	360	1/1	0.97	0.10	50,50,50,50	0
3	CL	A	2	1/1	0.98	0.09	45,45,45,45	0
3	CL	C	6	1/1	0.99	0.05	46,46,46,46	0

6.5 Other polymers [i](#)

There are no such residues in this entry.