



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 10:33 AM UTC

PDB ID : 3OS7 / pdb_00003os7
Title : Crystal structure of a galactose mutarotase-like protein (CA_C0697) from CLOSTRIDIUM ACETOBUTYLICUM at 1.80 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2010-09-08
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	341	 94% . .
1	B	341	 96% .
1	C	341	 94% 5% .
1	D	341	 94% 5% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 13142 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called galactose mutarotase-like protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	338	Total	C	N	O	S	Se	0	13	0
			2778	1781	453	535	5	4			
1	B	338	Total	C	N	O	S	Se	0	20	0
			2810	1812	451	536	7	4			
1	C	338	Total	C	N	O	S	Se	0	21	0
			2818	1820	456	533	5	4			
1	D	338	Total	C	N	O	S	Se	0	22	0
			2848	1831	463	544	6	4			

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q97L65
B	0	GLY	-	expression tag	UNP Q97L65
C	0	GLY	-	expression tag	UNP Q97L65
D	0	GLY	-	expression tag	UNP Q97L65

- Molecule 2 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C₈H₁₈O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			13	8	5		
2	A	1	Total	C	O	0	0
			10	6	4		
2	B	1	Total	C	O	0	0
			13	8	5		
2	C	1	Total	C	O	0	0
			13	8	5		
2	C	1	Total	C	O	0	0
			10	6	4		
2	D	1	Total	C	O	0	0
			13	8	5		
2	D	1	Total	C	O	0	0
			7	4	3		

- Molecule 3 is L(+)-TARTARIC ACID (CCD ID: TLA) (formula: C₄H₆O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			10	4	6		
3	B	1	Total	C	O	0	0
			10	4	6		
3	C	1	Total	C	O	0	0
			10	4	6		
3	D	1	Total	C	O	0	0
			10	4	6		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	C	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	406	Total O 406 406	0	3

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	439	Total 440	O 440	0	1
5	C	356	Total 357	O 357	0	4
5	D	488	Total 490	O 490	0	6

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: galactose mutarotase-like protein

Chain A:  94% . .




- Molecule 1: galactose mutarotase-like protein

Chain B:  96% . .




- Molecule 1: galactose mutarotase-like protein

Chain C:  94% 5% .



- Molecule 1: galactose mutarotase-like protein

Chain D:  94% 5% .



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	209.53Å 41.55Å 211.64Å 90.00° 119.58° 90.00°	Depositor
Resolution (Å)	29.55 – 1.80 29.55 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.0 (29.55-1.80) 98.9 (29.55-1.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.62 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.143 , 0.181 0.151 , 0.188	Depositor DCC
R_{free} test set	7348 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	16.4	Xtriage
Anisotropy	0.144	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.013 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	13142	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, TLA, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/2872	0.76	0/3871
1	B	0.77	0/2930	0.79	0/3945
1	C	0.75	0/2941	0.76	0/3958
1	D	0.82	2/2959 (0.1%)	0.81	0/3978
All	All	0.77	2/11702 (0.0%)	0.78	0/15752

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	291	MSE	SE-CE	-8.36	1.70	1.95
1	D	238	ALA	CA-CB	5.54	1.60	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2778	0	2739	13	0
1	B	2810	0	2817	11	0
1	C	2818	0	2840	18	0
1	D	2848	0	2855	10	0
2	A	23	0	31	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	13	0	18	0	0
2	C	23	0	31	4	0
2	D	20	0	27	1	0
3	A	10	0	4	0	0
3	B	10	0	4	0	0
3	C	10	0	4	0	0
3	D	10	0	4	0	0
4	A	12	0	18	2	0
4	B	28	0	42	5	0
4	C	24	0	36	1	0
4	D	12	0	18	0	0
5	A	406	0	0	5	0
5	B	440	0	0	4	0
5	C	357	0	0	8	0
5	D	490	0	0	3	0
All	All	13142	0	11488	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:13:ILE:CG2	1:B:19[B]:CYS:SG	2.54	0.96
1:B:13:ILE:HG21	1:B:19[B]:CYS:SG	2.09	0.92
1:B:13:ILE:HG22	1:B:19[B]:CYS:SG	2.17	0.84
4:B:346:EDO:H22	5:B:1697:HOH:O	1.82	0.78
1:C:21[B]:ARG:HG2	1:C:30:LEU:HD13	1.70	0.73
1:D:261:ARG:HH22	2:D:344:PG4:H41	1.55	0.72
1:B:258:LYS:HD2	5:B:1251:HOH:O	1.93	0.67
1:A:102:LYS:HB3	5:A:1600:HOH:O	1.94	0.67
1:D:62:GLU:CG	5:D:1647:HOH:O	2.42	0.66
1:C:49:VAL:HG22	1:C:192[B]:ILE:HG12	1.80	0.64
1:C:317:VAL:O	2:C:347:PG4:H62	1.99	0.62
1:D:62:GLU:HG3	5:D:1647:HOH:O	1.98	0.62
1:A:192[B]:ILE:HG22	1:A:193:GLU:O	2.02	0.59
1:C:240:ALA:O	5:C:1648:HOH:O	2.18	0.54
1:D:176[B]:GLU:HG3	1:D:322:THR:HB	1.90	0.53
1:C:129[B]:LEU:HD22	1:C:157[B]:GLU:HG2	1.92	0.52
1:D:314[B]:ASP:OD1	1:D:317:VAL:HG23	2.09	0.52
1:A:125[B]:GLN:O	1:A:126[B]:ASP:HB2	2.10	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:LYS:NZ	5:A:1021:HOH:O	2.42	0.50
1:B:135:ASP:H	4:B:346:EDO:C2	2.25	0.50
2:C:347:PG4:H41	5:C:916:HOH:O	2.13	0.49
1:C:49:VAL:HG22	1:C:192[B]:ILE:CG1	2.42	0.48
1:A:92[A]:LYS:CG	5:A:1667:HOH:O	2.62	0.48
1:C:5:ASP:OD2	5:C:1693:HOH:O	2.19	0.48
1:A:312:LYS:H	4:A:341:EDO:H22	1.79	0.47
1:C:283:LYS:NZ	2:C:347:PG4:O1	2.48	0.46
1:C:317:VAL:O	2:C:347:PG4:C6	2.64	0.45
1:C:231:GLU:HG2	5:C:1132:HOH:O	2.17	0.44
1:D:6:LEU:HD21	1:D:123[A]:ILE:CG2	2.47	0.44
1:C:321[A]:LYS:HB3	1:C:321[A]:LYS:HE3	1.54	0.44
1:B:135:ASP:H	4:B:346:EDO:H21	1.83	0.43
1:A:116:TRP:CZ3	1:A:154:LEU:HG	2.54	0.43
1:A:49:VAL:HG22	1:A:192[B]:ILE:HG12	2.01	0.43
1:B:7:TRP:HA	1:B:123[A]:ILE:HD11	2.01	0.43
1:B:88[B]:LYS:HE3	1:B:88[B]:LYS:HB3	1.86	0.43
1:B:283:LYS:O	4:B:347:EDO:H21	2.18	0.42
1:C:321[B]:LYS:HE3	5:C:1708[B]:HOH:O	2.19	0.42
1:A:312:LYS:H	4:A:341:EDO:C2	2.32	0.42
1:C:258:LYS:HD2	5:C:1632:HOH:O	2.20	0.42
1:B:176[A]:GLU:CG	1:B:322:THR:HB	2.49	0.42
1:D:164[A]:LYS:NZ	5:D:1193:HOH:O	2.49	0.42
1:C:151:GLN:O	4:C:342:EDO:H21	2.20	0.42
1:D:314[B]:ASP:OD2	1:D:316:SER:OG	2.23	0.41
1:A:314[A]:ASP:OD1	1:A:316:SER:OG	2.36	0.41
1:A:100:GLU:HG2	5:A:1600:HOH:O	2.20	0.41
1:A:62:GLU:HG3	5:A:1653:HOH:O	2.19	0.41
1:C:211[B]:LYS:HD3	5:C:1218:HOH:O	2.21	0.41
1:B:193:GLU:HB2	5:B:1012:HOH:O	2.20	0.40
1:D:185:SER:O	1:D:298:CYS:HA	2.20	0.40
4:B:347:EDO:O1	5:B:1304:HOH:O	2.21	0.40
1:C:74:PRO:HA	5:C:364:HOH:O	2.21	0.40
1:A:192[A]:ILE:HG21	1:A:270:LYS:HG3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	349/341 (102%)	338 (97%)	11 (3%)	0	100	100
1	B	356/341 (104%)	343 (96%)	13 (4%)	0	100	100
1	C	357/341 (105%)	344 (96%)	13 (4%)	0	100	100
1	D	358/341 (105%)	344 (96%)	14 (4%)	0	100	100
All	All	1420/1364 (104%)	1369 (96%)	51 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	313/302 (104%)	309 (99%)	4 (1%)	61	54
1	B	321/302 (106%)	319 (99%)	2 (1%)	78	77
1	C	321/302 (106%)	316 (98%)	5 (2%)	55	47
1	D	325/302 (108%)	324 (100%)	1 (0%)	86	86
All	All	1280/1208 (106%)	1268 (99%)	12 (1%)	76	67

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	125[A]	GLN
1	A	125[B]	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	152	PHE
1	A	193	GLU
1	B	152	PHE
1	B	193	GLU
1	C	152	PHE
1	C	193[A]	GLU
1	C	193[B]	GLU
1	C	321[A]	LYS
1	C	321[B]	LYS
1	D	152	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	151	GLN
1	C	271	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

30 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	EDO	C	346	-	3,3,3	0.51	0	2,2,2	0.43	0
4	EDO	A	341	-	3,3,3	0.27	0	2,2,2	1.02	0
4	EDO	A	342	-	3,3,3	0.48	0	2,2,2	0.27	0
2	PG4	C	400	-	12,12,12	0.50	0	11,11,11	0.55	0
4	EDO	B	342	-	3,3,3	0.53	0	2,2,2	0.42	0
2	PG4	A	400	-	12,12,12	0.61	0	11,11,11	0.50	0
2	PG4	D	400	-	12,12,12	0.44	0	11,11,11	0.46	0
4	EDO	D	342	-	3,3,3	0.60	0	2,2,2	0.27	0
3	TLA	B	401	-	9,9,9	1.21	0	12,12,12	1.08	2 (16%)
4	EDO	D	341	-	3,3,3	0.32	0	2,2,2	0.67	0
4	EDO	B	345	-	3,3,3	0.37	0	2,2,2	0.35	0
4	EDO	B	343	-	3,3,3	0.38	0	2,2,2	0.57	0
2	PG4	B	400	-	12,12,12	0.50	0	11,11,11	0.48	0
4	EDO	C	341	-	3,3,3	0.41	0	2,2,2	0.45	0
4	EDO	C	345	-	3,3,3	0.51	0	2,2,2	0.24	0
4	EDO	C	343	-	3,3,3	0.44	0	2,2,2	0.43	0
3	TLA	D	401	-	9,9,9	1.25	0	12,12,12	1.14	1 (8%)
3	TLA	A	401	-	9,9,9	1.35	1 (11%)	12,12,12	1.23	1 (8%)
2	PG4	C	347	-	9,9,12	0.59	0	8,8,11	0.64	0
4	EDO	B	347	-	3,3,3	0.61	0	2,2,2	0.32	0
4	EDO	C	342	-	3,3,3	0.51	0	2,2,2	0.23	0
4	EDO	B	341	-	3,3,3	0.51	0	2,2,2	0.27	0
4	EDO	A	343	-	3,3,3	0.35	0	2,2,2	0.55	0
4	EDO	B	344	-	3,3,3	0.41	0	2,2,2	0.48	0
4	EDO	C	344	-	3,3,3	0.41	0	2,2,2	0.20	0
2	PG4	D	344	-	6,6,12	0.57	0	5,5,11	0.67	0
2	PG4	A	344	-	9,9,12	0.53	0	8,8,11	0.46	0
4	EDO	D	343	-	3,3,3	0.59	0	2,2,2	0.17	0
3	TLA	C	401	-	9,9,9	1.33	0	12,12,12	1.36	1 (8%)
4	EDO	B	346	-	3,3,3	0.54	0	2,2,2	0.24	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	C	346	-	-	1/1/1/1	-
4	EDO	A	341	-	-	0/1/1/1	-
4	EDO	A	342	-	-	0/1/1/1	-
2	PG4	C	400	-	-	4/10/10/10	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	B	342	-	-	1/1/1/1	-
2	PG4	A	400	-	-	4/10/10/10	-
2	PG4	D	400	-	-	3/10/10/10	-
4	EDO	D	342	-	-	1/1/1/1	-
3	TLA	B	401	-	-	0/12/12/12	-
4	EDO	D	341	-	-	0/1/1/1	-
4	EDO	B	345	-	-	1/1/1/1	-
4	EDO	B	343	-	-	0/1/1/1	-
2	PG4	B	400	-	-	5/10/10/10	-
4	EDO	C	341	-	-	0/1/1/1	-
4	EDO	C	345	-	-	1/1/1/1	-
4	EDO	C	343	-	-	0/1/1/1	-
3	TLA	D	401	-	-	0/12/12/12	-
3	TLA	A	401	-	-	0/12/12/12	-
2	PG4	C	347	-	-	3/7/7/10	-
4	EDO	B	347	-	-	1/1/1/1	-
4	EDO	C	342	-	-	0/1/1/1	-
4	EDO	B	341	-	-	0/1/1/1	-
4	EDO	A	343	-	-	0/1/1/1	-
4	EDO	B	344	-	-	0/1/1/1	-
4	EDO	C	344	-	-	0/1/1/1	-
2	PG4	D	344	-	-	3/4/4/10	-
2	PG4	A	344	-	-	3/7/7/10	-
4	EDO	D	343	-	-	0/1/1/1	-
3	TLA	C	401	-	-	0/12/12/12	-
4	EDO	B	346	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	401	TLA	C2-C1	-2.47	1.49	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	TLA	O41-C4-C3	2.31	119.74	113.31
3	C	401	TLA	O41-C4-C3	2.31	119.73	113.31
3	B	401	TLA	O41-C4-C3	2.30	119.70	113.31
3	D	401	TLA	O11-C1-C2	2.08	119.08	113.31
3	B	401	TLA	O4-C4-C3	-2.02	116.22	121.62

There are no chirality outliers.

All (31) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	344	PG4	O2-C3-C4-O3
2	C	347	PG4	O2-C3-C4-O3
2	D	344	PG4	O1-C1-C2-O2
4	B	345	EDO	O1-C1-C2-O2
4	B	347	EDO	O1-C1-C2-O2
4	C	345	EDO	O1-C1-C2-O2
2	B	400	PG4	O4-C7-C8-O5
2	B	400	PG4	C8-C7-O4-C6
2	D	400	PG4	C5-C6-O4-C7
2	A	344	PG4	C6-C5-O3-C4
2	A	400	PG4	C5-C6-O4-C7
2	C	400	PG4	C5-C6-O4-C7
2	A	344	PG4	C4-C3-O2-C2
2	D	344	PG4	C4-C3-O2-C2
2	C	400	PG4	O1-C1-C2-O2
2	A	400	PG4	O1-C1-C2-O2
2	C	400	PG4	O3-C5-C6-O4
2	D	400	PG4	O2-C3-C4-O3
4	C	346	EDO	O1-C1-C2-O2
2	C	347	PG4	C1-C2-O2-C3
2	D	344	PG4	O2-C3-C4-O3
2	C	347	PG4	C4-C3-O2-C2
4	D	342	EDO	O1-C1-C2-O2
2	B	400	PG4	O2-C3-C4-O3
2	A	400	PG4	O2-C3-C4-O3
2	D	400	PG4	O4-C7-C8-O5
4	B	342	EDO	O1-C1-C2-O2
2	A	400	PG4	O3-C5-C6-O4
2	B	400	PG4	O3-C5-C6-O4
2	B	400	PG4	O1-C1-C2-O2
2	C	400	PG4	O2-C3-C4-O3

There are no ring outliers.

6 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	341	EDO	2	0
2	C	347	PG4	4	0
4	B	347	EDO	2	0
4	C	342	EDO	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	344	PG4	1	0
4	B	346	EDO	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	334/341 (97%)	-0.60	0	100 100	11, 18, 33, 47	13 (3%)
1	B	334/341 (97%)	-0.69	0	100 100	7, 15, 30, 44	20 (5%)
1	C	334/341 (97%)	-0.42	2 (0%)	85 86	9, 19, 38, 49	21 (6%)
1	D	334/341 (97%)	-0.71	3 (0%)	81 81	6, 14, 29, 46	22 (6%)
All	All	1336/1364 (97%)	-0.61	5 (0%)	88 89	6, 16, 34, 49	76 (5%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	3	ASN	2.8
1	C	3	ASN	2.7
1	D	124	ASP	2.6
1	D	125[A]	GLN	2.5
1	C	241	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	C	345	4/4	0.72	0.20	53,57,58,60	0
4	EDO	B	344	4/4	0.74	0.17	53,56,58,58	0
2	PG4	A	344	10/13	0.75	0.20	49,54,59,60	0
2	PG4	C	347	10/13	0.77	0.21	56,61,65,66	0
4	EDO	B	346	4/4	0.77	0.21	34,44,45,52	0
2	PG4	D	344	7/13	0.77	0.17	38,43,53,55	0
4	EDO	B	347	4/4	0.80	0.28	34,40,43,52	0
4	EDO	C	344	4/4	0.81	0.15	56,59,60,60	0
4	EDO	C	346	4/4	0.81	0.17	34,40,42,46	0
4	EDO	A	341	4/4	0.82	0.14	30,36,43,53	0
4	EDO	C	343	4/4	0.85	0.14	53,54,55,55	0
4	EDO	C	342	4/4	0.85	0.18	40,41,42,44	0
4	EDO	A	343	4/4	0.86	0.13	46,50,52,52	0
4	EDO	D	341	4/4	0.86	0.12	27,30,33,38	0
3	TLA	C	401	10/10	0.90	0.09	34,44,46,49	0
4	EDO	D	342	4/4	0.90	0.15	24,31,32,33	0
4	EDO	B	345	4/4	0.91	0.12	43,45,48,48	0
4	EDO	B	343	4/4	0.91	0.11	29,30,35,37	0
4	EDO	C	341	4/4	0.92	0.10	37,43,44,52	0
4	EDO	D	343	4/4	0.92	0.11	40,41,44,44	0
3	TLA	A	401	10/10	0.93	0.07	30,37,45,46	0
3	TLA	D	401	10/10	0.93	0.07	27,36,38,43	0
3	TLA	B	401	10/10	0.93	0.08	27,33,40,41	0
2	PG4	C	400	13/13	0.94	0.10	25,34,39,46	0
4	EDO	A	342	4/4	0.94	0.08	31,33,35,36	0
2	PG4	A	400	13/13	0.94	0.08	26,31,40,43	0
4	EDO	B	342	4/4	0.94	0.12	27,29,31,35	0
2	PG4	D	400	13/13	0.94	0.09	25,29,37,40	0
2	PG4	B	400	13/13	0.94	0.09	21,29,37,42	0
4	EDO	B	341	4/4	0.96	0.06	18,20,23,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.