



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 04:53 PM UTC

PDB ID : 3OSP / pdb_00003osp
Title : Structure of rev1
Authors : Nair, D.T.; Aggarwal, A.K.
Deposited on : 2010-09-09
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	434	3% 71% 24% ..
2	T	16	6% 56% 31% 12%
3	P	12	42% 25% 33%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 4158 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA repair protein REV1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	428	Total	C	N	O	S	0	0	0
			3419	2179	585	632	23			


- Molecule 2 is a DNA chain called 5'-D(*TP*AP*AP*(3DR)P*GP*TP*AP*GP*GP*GP*GP*AP*GP*GP*AP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	T	16	Total	C	N	O	P	0	0	0
			327	155	66	91	15			

- Molecule 3 is a DNA chain called 5'-D(*AP*TP*CP*CP*TP*CP*CP*CP*CP*TP*AP*(D OC))-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	12	Total	C	N	O	P	0	0	0
			231	113	37	70	11			

- Molecule 4 is 2'-DEOXYCYTIDINE-5'-TRIPHOSPHATE (CCD ID: DCP) (formula: C₉H₁₆N₃O₁₃P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			28	9	3	13	3		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Mg	0	0
			2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	139	Total	O	0	0
			139	139		
6	T	9	Total	O	0	0
			9	9		
6	P	3	Total	O	0	0
			3	3		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: DNA repair protein REV1



• Molecule 2: 5'-D(*TP*AP*AP*(3DR)P*GP*TP*AP*GP*GP*GP*GP*AP*GP*GP*AP*T)-3'



• Molecule 3: 5'-D(*AP*TP*CP*CP*TP*CP*CP*CP*CP*TP*AP*(DOC))-3'



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	180.02Å 200.24Å 55.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.50 50.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) (50.00-2.50) 94.0 (50.00-2.50)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.03 (at 2.51Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.204 , 0.255 0.210 , 0.251	Depositor DCC
R_{free} test set	2688 reflections (7.58%)	wwPDB-VP
Wilson B-factor (Å ²)	36.6	Xtriage
Anisotropy	0.717	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4158	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, DOC, DCP, 3DR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.27	7/3487 (0.2%)	1.33	20/4700 (0.4%)
2	T	0.85	1/356 (0.3%)	1.63	4/548 (0.7%)
3	P	0.88	0/236	1.79	5/360 (1.4%)
All	All	1.22	8/4079 (0.2%)	1.40	29/5608 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	708	ALA	CA-CB	-8.32	1.39	1.53
1	A	685	MET	SD-CE	-7.76	1.60	1.79
1	A	523	MET	SD-CE	-7.52	1.60	1.79
1	A	529	TYR	C-O	-5.59	1.16	1.23
1	A	520	ALA	CA-C	-5.37	1.45	1.52
1	A	694	SER	CA-C	5.30	1.59	1.52
1	A	389	ILE	CA-CB	-5.30	1.47	1.54
2	T	2	DA	O3'-P	5.01	1.68	1.61

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	525	LYS	CA-C-N	-18.18	97.11	119.84
1	A	525	LYS	C-N-CA	-18.18	97.11	119.84

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	T	6	DT	C5'-C4'-C3'	-10.98	98.44	114.90
1	A	455	LEU	CA-C-N	-6.49	109.14	121.54
1	A	455	LEU	C-N-CA	-6.49	109.14	121.54
1	A	532	THR	N-CA-C	6.29	124.21	110.80
2	T	9	DG	P-O5'-C5'	-6.27	110.60	120.00
1	A	711	MET	CG-SD-CE	-6.17	87.33	100.90
1	A	515	VAL	CB-CA-C	-6.16	104.09	111.97
1	A	661	THR	CB-CA-C	-6.16	97.20	109.33
3	P	11	DA	C4'-C3'-O3'	-6.10	100.85	110.00
1	A	525	LYS	N-CA-C	5.86	122.76	109.81
1	A	460	LEU	N-CA-C	5.65	117.69	108.76
3	P	7	DC	C4'-C3'-O3'	5.59	118.39	110.00
3	P	8	DC	OP1-P-OP2	5.44	136.32	120.00
1	A	733	ASN	N-CA-C	5.43	119.62	112.89
1	A	688	CYS	N-CA-C	5.42	117.42	109.24
2	T	6	DT	C6-N1-C1'	5.34	127.36	119.35
1	A	433	SER	N-CA-C	5.31	116.95	108.41
1	A	525	LYS	C-N-CD	5.27	146.61	125.00
2	T	8	DG	P-O3'-C3'	5.22	128.03	120.20
3	P	1	DA	P-O3'-C3'	5.19	127.98	120.20
1	A	666	LEU	CA-CB-CG	5.12	134.21	116.30
1	A	381	ALA	CA-C-N	5.11	128.36	121.05
1	A	381	ALA	C-N-CA	5.11	128.36	121.05
1	A	699	ILE	N-CA-CB	-5.07	106.22	111.64
1	A	540	GLU	N-CA-C	-5.07	105.84	111.36
3	P	11	DA	C5'-C4'-O4'	5.05	116.97	109.40
1	A	382	CYS	N-CA-C	5.03	117.64	110.24

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	347	TYR	Peptide
1	A	524	ALA	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3419	0	3455	74	0
2	T	327	0	178	7	0
3	P	231	0	137	8	0
4	A	28	0	12	2	0
5	A	2	0	0	0	0
6	A	139	0	0	2	0
6	P	3	0	0	0	0
6	T	9	0	0	0	0
All	All	4158	0	3782	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (86) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:501:THR:HG21	1:A:504:CYS:HB2	1.44	0.97
3:P:11:DA:H2''	3:P:12:DOC:H5''	1.45	0.97
2:T:3:DA:H5''	2:T:4:3DR:H5'	1.51	0.93
1:A:497:ILE:O	1:A:501:THR:HB	1.70	0.92
3:P:7:DC:OP1	3:P:7:DC:H4'	1.74	0.86
1:A:501:THR:CG2	1:A:504:CYS:HB2	2.14	0.78
1:A:326:HIS:HD2	2:T:6:DT:OP1	1.67	0.77
1:A:501:THR:CG2	1:A:504:CYS:H	2.01	0.74
1:A:364:ASP:OD2	1:A:505:THR:HG22	1.88	0.72
1:A:501:THR:HG22	1:A:504:CYS:H	1.56	0.70
1:A:318:SER:O	1:A:322:HIS:HD2	1.74	0.70
1:A:525:LYS:O	1:A:527:ASN:N	2.26	0.68
1:A:669:MET:HE3	1:A:685:MET:HE1	1.74	0.67
1:A:668:LEU:HD12	1:A:668:LEU:C	2.20	0.66
3:P:1:DA:H1'	3:P:2:DT:C5'	2.26	0.66
1:A:365:CYS:HA	4:A:739:DCP:O1B	1.96	0.65
1:A:635:ASN:C	1:A:635:ASN:HD22	2.04	0.64
1:A:525:LYS:O	1:A:526:PRO:C	2.35	0.64
1:A:460:LEU:HD12	1:A:460:LEU:C	2.23	0.64
1:A:635:ASN:ND2	1:A:638:GLN:H	1.97	0.62
1:A:362:ASP:OD1	1:A:525:LYS:NZ	2.31	0.62
1:A:669:MET:HG3	1:A:685:MET:HE1	1.82	0.61
1:A:573:ASP:O	1:A:577:ARG:HG3	2.01	0.60
1:A:510:CYS:O	1:A:531:ILE:O	2.21	0.58
1:A:520:ALA:HB1	1:A:530:ASN:HD22	1.67	0.58
1:A:674:ASP:CB	1:A:687:ARG:HH22	2.17	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:632:ARG:NH1	1:A:678:GLU:OE1	2.37	0.57
1:A:661:THR:CG2	1:A:732:PHE:HB3	2.33	0.57
2:T:3:DA:H5''	2:T:4:3DR:C5'	2.29	0.57
1:A:348:THR:HG23	1:A:614:PRO:HG2	1.88	0.56
1:A:593:MET:HE3	1:A:597:LEU:HG	1.87	0.56
1:A:323:SER:O	1:A:327:HIS:HD2	1.89	0.55
1:A:349:LYS:HB3	1:A:612:TYR:O	2.07	0.55
1:A:640:ASP:OD2	1:A:716:ARG:NH2	2.40	0.54
1:A:501:THR:HG21	1:A:504:CYS:CB	2.28	0.54
1:A:365:CYS:HA	4:A:739:DCP:PB	2.48	0.53
1:A:661:THR:HG23	1:A:732:PHE:HB3	1.89	0.53
1:A:331:TRP:HE1	1:A:442:GLN:HE21	1.56	0.53
3:P:7:DC:H2''	3:P:8:DC:H5'	1.90	0.53
1:A:364:ASP:OD2	1:A:505:THR:CG2	2.55	0.53
1:A:674:ASP:HB2	1:A:687:ARG:HH22	1.75	0.52
1:A:604:ASP:OD1	1:A:607:SER:OG	2.26	0.52
1:A:339:PHE:CD2	1:A:618:LEU:HD12	2.46	0.51
2:T:3:DA:C5'	2:T:4:3DR:H5'	2.33	0.51
1:A:501:THR:CG2	1:A:504:CYS:CB	2.86	0.51
3:P:2:DT:H2''	3:P:3:DC:OP2	2.10	0.51
1:A:495:GLN:OE1	1:A:499:GLN:NE2	2.45	0.50
1:A:523:MET:HE3	1:A:530:ASN:OD1	2.13	0.49
1:A:619:GLN:HG3	6:A:841:HOH:O	2.14	0.48
3:P:1:DA:H1'	3:P:2:DT:H5''	1.95	0.48
1:A:363:PHE:CD1	1:A:467:ASP:HB2	2.49	0.47
1:A:635:ASN:C	1:A:635:ASN:ND2	2.71	0.47
1:A:402:SER:HA	1:A:414:ASN:OD1	2.15	0.47
1:A:674:ASP:HB2	1:A:687:ARG:NH2	2.29	0.47
1:A:405:TYR:OH	1:A:525:LYS:HG2	2.16	0.46
1:A:667:LYS:HB3	1:A:729:ALA:HB3	1.98	0.46
1:A:460:LEU:CD1	1:A:462:LEU:HG	2.45	0.46
1:A:322:HIS:CE1	2:T:3:DA:N6	2.84	0.46
1:A:364:ASP:O	1:A:365:CYS:C	2.59	0.45
1:A:654:LEU:HD22	1:A:659:LYS:O	2.16	0.45
1:A:487:ALA:O	1:A:491:GLU:HG3	2.16	0.45
1:A:501:THR:CG2	1:A:504:CYS:N	2.75	0.45
1:A:599:LEU:HA	1:A:599:LEU:HD23	1.73	0.45
1:A:407:ALA:O	1:A:412:ILE:HB	2.16	0.45
2:T:5:DG:H2''	2:T:6:DT:H5''	1.99	0.45
3:P:7:DC:OP1	3:P:7:DC:C4'	2.54	0.45
1:A:674:ASP:HB3	1:A:687:ARG:HH22	1.80	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:531:ILE:O	1:A:532:THR:C	2.60	0.44
1:A:669:MET:CE	1:A:685:MET:HE1	2.46	0.43
1:A:318:SER:O	1:A:322:HIS:CD2	2.62	0.43
1:A:539:GLU:HG2	1:A:572:ASN:HD21	1.82	0.43
1:A:669:MET:CG	1:A:685:MET:HE1	2.46	0.43
1:A:668:LEU:HD12	1:A:668:LEU:O	2.18	0.43
1:A:547:LEU:HB2	1:A:558:LEU:HD23	2.01	0.43
1:A:619:GLN:CG	6:A:841:HOH:O	2.67	0.42
1:A:661:THR:HG21	1:A:732:PHE:HB3	2.02	0.42
1:A:520:ALA:CB	1:A:530:ASN:HD22	2.31	0.42
1:A:635:ASN:HD21	1:A:638:GLN:H	1.68	0.42
1:A:632:ARG:NH2	1:A:684:GLY:O	2.53	0.41
1:A:458:PHE:CG	1:A:471:CYS:HB3	2.56	0.41
1:A:322:HIS:CG	2:T:3:DA:C6	3.09	0.41
3:P:1:DA:H1'	3:P:2:DT:H5'	2.02	0.41
1:A:360:HIS:CD2	1:A:521:LEU:HD22	2.56	0.40
1:A:665:THR:OG1	1:A:733:ASN:ND2	2.51	0.40
1:A:470:VAL:O	1:A:470:VAL:HG12	2.18	0.40
1:A:493:ILE:HG22	1:A:497:ILE:HD12	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	424/434 (98%)	408 (96%)	14 (3%)	2 (0%)	24 43

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	306	ARG
1	A	532	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	385/391 (98%)	347 (90%)	38 (10%)	7 16

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	305	LYS
1	A	328	LEU
1	A	337	ASP
1	A	344	ILE
1	A	346	LYS
1	A	348	THR
1	A	353	LYS
1	A	362	ASP
1	A	375	ARG
1	A	386	ARG
1	A	431	LEU
1	A	466	ILE
1	A	477	ASP
1	A	484	THR
1	A	495	GLN
1	A	501	THR
1	A	505	THR
1	A	506	VAL
1	A	515	VAL
1	A	556	SER
1	A	558	LEU
1	A	559	SER
1	A	574	LEU
1	A	580	LEU
1	A	583	LEU
1	A	589	SER
1	A	591	LEU
1	A	607	SER
1	A	619	GLN
1	A	635	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	653	LYS
1	A	654	LEU
1	A	661	THR
1	A	666	LEU
1	A	695	SER
1	A	723	MET
1	A	724	GLU
1	A	728	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	322	HIS
1	A	326	HIS
1	A	327	HIS
1	A	397	ASN
1	A	420	GLN
1	A	440	GLN
1	A	442	GLN
1	A	459	ASN
1	A	495	GLN
1	A	499	GLN
1	A	530	ASN
1	A	572	ASN
1	A	602	GLN
1	A	635	ASN
1	A	648	GLN
1	A	655	ASN
1	A	663	GLN
1	A	731	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	3DR	T	4	2	8,11,12	1.69	1 (12%)	7,14,17	2.31	2 (28%)
3	DOC	P	12	2,3	16,19,20	0.41	0	20,26,29	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3DR	T	4	2	-	1/3/15/16	0/1/1/1
3	DOC	P	12	2,3	-	4/7/18/19	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	T	4	3DR	O5'-C5'	-3.90	1.32	1.44

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	T	4	3DR	O4'-C4'-C3'	-5.02	96.34	103.73
2	T	4	3DR	O3'-C3'-C2'	-2.51	105.54	111.43

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	P	12	DOC	O4'-C1'-N1-C2
3	P	12	DOC	O4'-C1'-N1-C6
2	T	4	3DR	O4'-C4'-C5'-O5'
3	P	12	DOC	C2'-C1'-N1-C6
3	P	12	DOC	C2'-C1'-N1-C2

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	T	4	3DR	3	0
3	P	12	DOC	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	DCP	A	739	5	28,29,29	2.81	9 (32%)	41,45,45	2.10	9 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DCP	A	739	5	-	4/22/34/34	0/2/2/2

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	739	DCP	PA-O3A	9.28	1.69	1.59
4	A	739	DCP	C1'-N1	5.42	1.61	1.48
4	A	739	DCP	PA-O5'	-5.11	1.39	1.59
4	A	739	DCP	PB-O3A	4.99	1.64	1.59
4	A	739	DCP	PA-O1A	-3.38	1.39	1.50
4	A	739	DCP	PG-O1G	2.92	1.59	1.50
4	A	739	DCP	O5'-C5'	2.71	1.54	1.44
4	A	739	DCP	C2-N1	2.48	1.45	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	739	DCP	C4-N3	2.19	1.38	1.34

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	739	DCP	C2'-C1'-N1	-7.52	95.00	113.81
4	A	739	DCP	O4'-C1'-N1	5.38	117.42	107.86
4	A	739	DCP	O5'-C5'-C4'	3.78	121.86	108.99
4	A	739	DCP	O2B-PB-O3B	3.46	116.62	107.27
4	A	739	DCP	O4'-C4'-C5'	3.23	119.69	109.33
4	A	739	DCP	O4'-C4'-C3'	2.34	110.99	105.65
4	A	739	DCP	O3A-PB-O1B	-2.32	103.73	110.70
4	A	739	DCP	C1'-N1-C2	-2.21	114.03	117.83
4	A	739	DCP	O3B-PG-O1G	-2.21	99.40	111.04

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	739	DCP	PG-O3B-PB-O1B
4	A	739	DCP	PB-O3A-PA-O1A
4	A	739	DCP	PG-O3B-PB-O2B
4	A	739	DCP	PB-O3A-PA-O2A


There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	739	DCP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	428/434 (98%)	-0.16	12 (2%) 55 50	16, 31, 55, 77	0
2	T	15/16 (93%)	-0.41	1 (6%) 24 21	21, 30, 59, 79	0
3	P	11/12 (91%)	-0.27	0 100 100	26, 32, 65, 65	0
All	All	454/462 (98%)	-0.17	13 (2%) 53 49	16, 31, 57, 79	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	477	ASP	4.2
1	A	484	THR	4.0
1	A	347	TYR	3.8
1	A	476	PRO	3.7
2	T	16	DT	3.2
1	A	305	LYS	3.0
1	A	353	LYS	2.7
1	A	348	THR	2.7
1	A	386	ARG	2.5
1	A	526	PRO	2.3
1	A	525	LYS	2.2
1	A	736	VAL	2.1
1	A	349	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	DOC	P	12	18/19	0.86	0.18	52,60,67,67	0
2	3DR	T	4	11/12	0.97	0.07	21,24,29,29	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	MG	A	297	1/1	0.90	0.11	53,53,53,53	0
4	DCP	A	739	28/28	0.96	0.09	26,29,31,32	0
5	MG	A	298	1/1	0.98	0.07	21,21,21,21	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around DCP A 739:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.