



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 05:10 AM UTC

PDB ID : 5OSX / pdb_00005osx
Title : Translation initiation factor 4E in complex with m7G(5'S)ppp(5'S)G mRNA 5' cap analog
Authors : Warminski, M.; Nowak, E.; Wojtczak, B.A.; Fac-Dabrowska, K.; Kubacka, D.; Nowicka, A.; Sikorski, P.J.; Kowalska, J.; Jemielity, J.; Nowotny, M.
Deposited on : 2017-08-18
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1188 (1.92-1.92)
Clashscore	190562	1209 (1.92-1.92)
Ramachandran outliers	187476	1195 (1.92-1.92)
Sidechain outliers	187428	1195 (1.92-1.92)
RSRZ outliers	180081	1188 (1.92-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	190	5% 87% 7% 6%
1	B	190	6% 81% 13% 6%
1	C	190	27% 70% 26%
1	D	190	24% 59% 8% 33%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 5091 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Eukaryotic translation initiation factor 4E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	178	Total	C	N	O	S	0	3	0
			1428	915	246	261	6			
1	B	178	Total	C	N	O	S	0	1	0
			1412	904	245	258	5			
1	C	141	Total	C	N	O	S	0	0	0
			1016	655	174	184	3			
1	D	128	Total	C	N	O	S	0	2	0
			951	606	165	176	4			

- Molecule 2 is [(2 {S},3 {S},4 {R},5 {R})-5-(2-azanyl-7-methyl-6-oxidanylidene-3 {H}-purin-7-ium-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methysulfanyl-[[[(3 {R},4 {S})-5-(2-azanyl-6-oxidanylidene-1 {H}-purin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methysulfanyl-oxidanyl-phosphoryl]oxy-oxidanyl-phosphoryl]oxy-phosphinic acid (CCD ID: AKW) (formula: C₂₁H₃₀N₁₀O₁₆P₃S₂).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	S	0	0
			33	11	5	12	3	2		
2	B	1	Total	C	N	O	P	S	0	0
			33	11	5	12	3	2		

- Molecule 3 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	K	0	0
			1	1		



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	83	Total	O	0	0
			83	83		
4	B	92	Total	O	0	0
			92	92		
4	C	21	Total	O	0	0
			21	21		
4	D	21	Total	O	0	0
			21	21		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Eukaryotic translation initiation factor 4E



- Molecule 1: Eukaryotic translation initiation factor 4E




- Molecule 1: Eukaryotic translation initiation factor 4E



- Molecule 1: Eukaryotic translation initiation factor 4E





HIS
ALA
ASP
THR
ALA
THR
LYS
SER
GLY
SER
THR
THR
LYS
ASN
ARG
PHE
VAL
VAL

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	37.84Å 37.85Å 145.99Å 86.56° 84.56° 77.05°	Depositor
Resolution (Å)	48.41 – 1.92 48.41 – 1.92	Depositor EDS
% Data completeness (in resolution range)	95.1 (48.41-1.92) 95.5 (48.41-1.92)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.13 (at 1.91Å)	Xtriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, R_{free}	0.224 , 0.269 0.226 , 0.269	Depositor DCC
R_{free} test set	2100 reflections (3.65%)	wwPDB-VP
Wilson B-factor (Å ²)	28.3	Xtriage
Anisotropy	0.585	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 67.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	0.068 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5091	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 17.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AKW, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/1474	0.54	0/2002
1	B	0.39	0/1452	0.56	0/1976
1	C	0.24	0/1037	0.42	0/1415
1	D	0.22	0/981	0.43	1/1341 (0.1%)
All	All	0.32	0/4944	0.50	1/6734 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	143	ASP	CB-CA-C	-5.99	109.68	116.63

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1428	0	1330	7	0
1	B	1412	0	1305	13	0
1	C	1016	0	859	5	0
1	D	951	0	794	10	0
2	A	33	0	0	0	0
2	B	33	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
4	A	83	0	0	0	0
4	B	92	0	0	3	0
4	C	21	0	0	0	0
4	D	21	0	0	0	0
All	All	5091	0	4288	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:47:PHE:HB2	1:A:63:ILE:HD11	1.60	0.83
1:C:65:LYS:NZ	1:C:96:ASP:OD1	2.26	0.69
1:D:46:TRP:HB2	1:D:92:SER:HB2	1.76	0.68
1:B:51:ASP:OD2	1:B:54:LYS:NZ	2.27	0.67
1:B:137:LEU:HD22	1:B:149:VAL:HG11	1.80	0.64
1:B:47:PHE:HB2	1:B:63:ILE:HD11	1.86	0.58
1:D:88:GLY:N	1:D:156:VAL:O	2.31	0.53
1:A:46:TRP:HB2	1:A:92:SER:HB2	1.92	0.51
1:B:122:ARG:NH2	1:B:157:ARG:O	2.45	0.50
1:C:83:SER:OG	1:C:127:ASP:OD2	2.28	0.48
1:D:47:PHE:HA	1:D:90:ASP:O	2.13	0.48
1:B:50:ASN:ND2	4:B:408:HOH:O	2.47	0.47
1:D:46:TRP:N	1:D:92:SER:O	2.36	0.47
1:B:55:THR:HG23	4:B:456:HOH:O	2.16	0.46
1:B:123:ARG:HD3	4:B:459:HOH:O	2.16	0.45
1:D:46:TRP:CG	1:D:100:PRO:HG3	2.51	0.45
1:A:83:SER:OG	1:A:127:ASP:OD2	2.35	0.45
1:B:46:TRP:HB2	1:B:92:SER:HB2	1.99	0.45
1:B:169:GLU:OE2	1:B:171:GLU:HB2	2.17	0.45
1:B:35:ILE:HG22	1:B:36:LYS:O	2.18	0.44
1:C:85:LEU:HD11	1:C:91:TYR:CE2	2.53	0.43
1:B:111:GLY:HA3	1:B:199:SER:HA	2.01	0.43
1:A:177:THR:O	1:A:181:ARG:HG3	2.18	0.43
1:C:122:ARG:NH2	1:C:161:ASP:OD1	2.51	0.43
1:D:87:PRO:HA	1:D:156:VAL:HG12	2.01	0.43
1:C:122:ARG:HE	1:C:122:ARG:HB2	1.69	0.42
1:B:83:SER:OG	1:B:127:ASP:OD2	2.35	0.42
1:D:117:LEU:HD22	1:D:121:GLN:NE2	2.34	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:MET:HB3	1:A:101:MET:HE2	1.72	0.42
1:B:66:PHE:CE1	1:B:72:PHE:HB2	2.55	0.42
1:D:116:THR:O	1:D:117:LEU:HD23	2.19	0.42
1:A:42:ARG:HD2	1:A:96:ASP:OD2	2.20	0.41
1:D:162:LYS:HE3	1:D:162:LYS:HB2	1.74	0.41
1:A:46:TRP:CZ2	1:A:62:LEU:HD12	2.55	0.41
1:D:121:GLN:HG2	1:D:125:ASP:OD2	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	177/190 (93%)	173 (98%)	4 (2%)	0	100	100
1	B	175/190 (92%)	171 (98%)	4 (2%)	0	100	100
1	C	133/190 (70%)	124 (93%)	9 (7%)	0	100	100
1	D	126/190 (66%)	122 (97%)	4 (3%)	0	100	100
All	All	611/760 (80%)	590 (97%)	21 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	145/169 (86%)	144 (99%)	1 (1%)	76	73
1	B	142/169 (84%)	141 (99%)	1 (1%)	76	73
1	C	84/169 (50%)	84 (100%)	0	100	100
1	D	82/169 (48%)	81 (99%)	1 (1%)	63	57
All	All	453/676 (67%)	450 (99%)	3 (1%)	76	73

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	213	ASN
1	B	148	ASP
1	D	64	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	107	ASN
1	A	172	ASN
1	B	84	ASN
1	B	155	ASN
1	C	84	ASN
1	D	155	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	AKW	B	301	-	30,35,57	2.19	9 (30%)	35,56,90	1.57	7 (20%)
2	AKW	A	301	-	30,35,57	2.29	9 (30%)	35,56,90	1.56	10 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AKW	B	301	-	-	0/15/38/64	0/3/3/6
2	AKW	A	301	-	-	0/15/38/64	0/3/3/6

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	AKW	O12-C11	7.66	1.37	1.23
2	A	301	AKW	O12-C11	7.62	1.37	1.23
2	A	301	AKW	C8-N2	-4.65	1.33	1.39
2	A	301	AKW	C9-N1	-4.26	1.32	1.37
2	B	301	AKW	C8-N2	-4.15	1.34	1.39
2	B	301	AKW	C9-N1	-3.47	1.33	1.37
2	B	301	AKW	P1-O1	-3.30	1.48	1.56
2	A	301	AKW	P3-O8	-3.21	1.48	1.56
2	B	301	AKW	P3-O8	-3.02	1.49	1.56
2	A	301	AKW	P1-O1	-3.01	1.49	1.56
2	A	301	AKW	P2-O6	2.68	1.62	1.59
2	A	301	AKW	C11-N5	-2.66	1.33	1.38
2	A	301	AKW	O9-C5	2.59	1.48	1.42
2	A	301	AKW	P2-O3	2.33	1.62	1.59
2	B	301	AKW	P2-O3	2.26	1.61	1.59
2	B	301	AKW	O9-C5	2.25	1.47	1.42
2	B	301	AKW	P2-O6	2.21	1.61	1.59
2	B	301	AKW	C11-N5	-2.19	1.34	1.38

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	AKW	O4-P2-O3	3.96	117.98	107.27
2	A	301	AKW	C7-N2-C8	3.37	130.99	126.80
2	A	301	AKW	C7-N2-C6	-2.97	120.30	124.79
2	A	301	AKW	O4-P2-O3	2.88	115.04	107.27
2	B	301	AKW	O8-P3-O7	-2.86	103.62	109.86
2	B	301	AKW	O3-P2-O5	-2.76	102.40	110.70
2	B	301	AKW	C7-N2-C8	2.53	129.95	126.80
2	A	301	AKW	O4-P2-O6	2.41	113.78	107.27
2	A	301	AKW	O1-P1-O3	2.39	113.74	107.27
2	B	301	AKW	C7-N2-C6	-2.32	121.28	124.79
2	B	301	AKW	O4-P2-O6	2.24	113.34	107.27
2	A	301	AKW	O3-P2-O5	-2.24	103.97	110.70
2	A	301	AKW	O8-P3-O7	-2.18	105.11	109.86
2	A	301	AKW	N4-C10-N3	2.13	121.26	116.76
2	A	301	AKW	C11-C8-N2	2.03	134.78	131.46
2	A	301	AKW	O6-P2-O5	-2.02	104.62	110.70
2	B	301	AKW	N1-C6-N2	-2.01	107.61	112.48


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	178/190 (93%)	0.60	10 (5%) 30 35	18, 35, 61, 85	4 (2%)
1	B	178/190 (93%)	0.68	11 (6%) 26 31	21, 34, 60, 89	1 (0%)
1	C	141/190 (74%)	1.74	52 (36%) 1 0	29, 61, 91, 101	1 (0%)
1	D	128/190 (67%)	1.75	45 (35%) 1 0	24, 62, 86, 98	3 (2%)
All	All	625/760 (82%)	1.11	118 (18%) 3 4	18, 42, 82, 101	9 (1%)

All (118) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	113	TRP	7.6
1	C	104	ASP	4.9
1	D	102	TRP	4.8
1	C	179	ILE	4.4
1	C	177	THR	4.4
1	C	102	TRP	4.2
1	B	193	ILE	4.0
1	B	34	TYR	4.0
1	C	175	ALA	3.9
1	C	182	VAL	3.9
1	D	113	TRP	3.9
1	D	58	ALA	3.9
1	C	142	PHE	3.9
1	D	88	GLY	3.7
1	B	32	GLU	3.7
1	C	114	LEU	3.7
1	D	32	GLU	3.7
1	D	85	LEU	3.6
1	D	48	PHE	3.6
1	B	203	THR	3.5
1	C	180	GLY	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	167	THR	3.5
1	C	48	PHE	3.5
1	D	114	LEU	3.5
1	C	145	TYR	3.4
1	D	149	VAL	3.4
1	A	203	THR	3.4
1	D	165	ILE	3.4
1	D	110	GLY	3.3
1	A	193	ILE	3.3
1	C	165	ILE	3.3
1	D	103	GLU	3.3
1	B	33	HIS	3.3
1	D	111	GLY	3.3
1	C	183	TYR	3.2
1	D	112	ARG	3.2
1	A	191	PRO	3.1
1	D	156	VAL	3.1
1	C	117	LEU	3.1
1	C	40	GLN	3.1
1	C	120	GLN	3.1
1	C	174	ASP	3.0
1	D	47	PHE	3.0
1	D	148	ASP	3.0
1	D	101	MET	3.0
1	C	173	ARG	2.9
1	C	187	LEU	2.9
1	C	164	ALA	2.9
1	C	46	TRP	2.9
1	C	188	GLY	2.9
1	B	201	ALA	2.9
1	D	86	MET	2.9
1	D	117	LEU	2.9
1	D	116	THR	2.8
1	C	97	GLY	2.8
1	A	33	HIS	2.8
1	D	150	CYS	2.7
1	C	167	THR	2.7
1	C	60	LEU	2.7
1	C	32	GLU	2.7
1	B	175	ALA	2.6
1	C	86	MET	2.6
1	D	129	PHE	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	37[A]	HIS	2.6
1	D	163	ILE	2.6
1	B	191	PRO	2.6
1	A	32	GLU	2.6
1	A	125	ASP	2.6
1	C	178	HIS	2.5
1	D	87	PRO	2.5
1	C	148	ASP	2.5
1	D	166	TRP	2.5
1	D	100	PRO	2.5
1	C	150	CYS	2.5
1	D	160	GLY	2.5
1	C	35	ILE	2.5
1	D	33	HIS	2.5
1	C	149	VAL	2.5
1	D	91	TYR	2.5
1	C	33	HIS	2.5
1	D	98	ILE	2.5
1	B	217	VAL	2.5
1	A	194	VAL	2.4
1	C	116	THR	2.4
1	C	100	PRO	2.4
1	D	105	GLU	2.4
1	D	136	CYS	2.4
1	C	181	ARG	2.3
1	C	154	VAL	2.3
1	C	158	ALA	2.3
1	D	97	GLY	2.3
1	D	137	LEU	2.3
1	C	115	ILE	2.3
1	D	57	GLN	2.3
1	C	129	PHE	2.2
1	C	147	ASP	2.2
1	C	54	LYS	2.2
1	D	60	LEU	2.2
1	D	62	LEU	2.2
1	C	144	ASP	2.2
1	D	74	ALA	2.2
1	C	51	ASP	2.2
1	A	160	GLY	2.2
1	A	217	VAL	2.1
1	B	136	CYS	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	94	PHE	2.1
1	D	142	PHE	2.1
1	D	115	ILE	2.1
1	B	214	ARG	2.1
1	C	98	ILE	2.1
1	D	144	ASP	2.1
1	C	53	SER	2.1
1	A	212	LYS	2.1
1	C	160	GLY	2.1
1	C	101	MET	2.1
1	D	158	ALA	2.1
1	C	62	LEU	2.0
1	C	143	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	AKW	A	301	33/52	0.94	0.09	27,41,61,66	0
2	AKW	B	301	33/52	0.94	0.08	30,38,56,59	0
3	K	B	302	1/1	0.97	0.08	64,64,64,64	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around AKW A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around AKW B 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.