



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 05:11 PM UTC

PDB ID : 8OS6 / pdb_00008os6
Title : Structure of a GFRA1/GDNF LICAM complex
Authors : Houghton, F.M.; Adams, S.E.; Briggs, D.C.; McDonald, N.Q.
Deposited on : 2023-04-18
Resolution : 2.66 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	260	71%8%20%
1	C	260	2% 72%7%22%
1	E	260	2% 72%7%21%
1	G	260	3% 70%9%22%
1	I	260	2% 68%8%23%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	K	260	
1	M	260	
1	O	260	
1	Q	260	
1	S	260	
2	B	99	
2	D	99	
2	F	99	
2	H	99	
2	J	99	
2	L	99	
2	N	99	
2	P	99	
2	R	99	
2	T	99	
3	U	2	
3	Y	2	
3	Z	2	
3	b	2	
4	V	3	
4	W	3	
4	X	3	
5	a	2	

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 46489 atoms, of which 22664 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GDNF family receptor alpha.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	207	Total	C	H	N	O	S	0	0	0
			3128	981	1530	285	310	22			
1	C	204	Total	C	H	N	O	S	0	0	0
			3071	967	1496	281	305	22			
1	E	205	Total	C	H	N	O	S	0	0	0
			3087	971	1507	281	306	22			
1	G	204	Total	C	H	N	O	S	0	0	0
			3073	969	1495	281	306	22			
1	I	201	Total	C	H	N	O	S	0	0	0
			3055	958	1498	278	299	22			
1	K	202	Total	C	H	N	O	S	0	0	0
			3043	959	1483	278	301	22			
1	M	204	Total	C	H	N	O	S	0	0	0
			3086	967	1511	281	305	22			
1	O	205	Total	C	H	N	O	S	0	0	0
			3084	968	1509	282	303	22			
1	Q	203	Total	C	H	N	O	S	0	0	0
			3061	962	1494	279	304	22			
1	S	203	Total	C	H	N	O	S	0	0	0
			3060	964	1491	280	303	22			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	353	GLY	-	expression tag	UNP Q98TT9
A	354	SER	-	expression tag	UNP Q98TT9
A	355	GLU	-	expression tag	UNP Q98TT9
C	353	GLY	-	expression tag	UNP Q98TT9
C	354	SER	-	expression tag	UNP Q98TT9
C	355	GLU	-	expression tag	UNP Q98TT9
E	353	GLY	-	expression tag	UNP Q98TT9
E	354	SER	-	expression tag	UNP Q98TT9
E	355	GLU	-	expression tag	UNP Q98TT9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	353	GLY	-	expression tag	UNP Q98TT9
G	354	SER	-	expression tag	UNP Q98TT9
G	355	GLU	-	expression tag	UNP Q98TT9
I	353	GLY	-	expression tag	UNP Q98TT9
I	354	SER	-	expression tag	UNP Q98TT9
I	355	GLU	-	expression tag	UNP Q98TT9
K	353	GLY	-	expression tag	UNP Q98TT9
K	354	SER	-	expression tag	UNP Q98TT9
K	355	GLU	-	expression tag	UNP Q98TT9
M	353	GLY	-	expression tag	UNP Q98TT9
M	354	SER	-	expression tag	UNP Q98TT9
M	355	GLU	-	expression tag	UNP Q98TT9
O	353	GLY	-	expression tag	UNP Q98TT9
O	354	SER	-	expression tag	UNP Q98TT9
O	355	GLU	-	expression tag	UNP Q98TT9
Q	353	GLY	-	expression tag	UNP Q98TT9
Q	354	SER	-	expression tag	UNP Q98TT9
Q	355	GLU	-	expression tag	UNP Q98TT9
S	353	GLY	-	expression tag	UNP Q98TT9
S	354	SER	-	expression tag	UNP Q98TT9
S	355	GLU	-	expression tag	UNP Q98TT9

- Molecule 2 is a protein called Glial cell line-derived neurotrophic factor.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	97	Total	C	H	N	O	S	0	0	0
			1481	473	720	132	149	7			
2	D	98	Total	C	H	N	O	S	0	0	0
			1535	485	752	138	153	7			
2	F	98	Total	C	H	N	O	S	0	0	0
			1522	485	741	136	153	7			
2	H	96	Total	C	H	N	O	S	0	0	0
			1500	478	732	133	150	7			
2	J	98	Total	C	H	N	O	S	0	0	0
			1518	483	741	135	152	7			
2	L	96	Total	C	H	N	O	S	0	0	0
			1493	475	729	132	150	7			
2	N	95	Total	C	H	N	O	S	0	0	0
			1502	475	739	132	149	7			
2	P	95	Total	C	H	N	O	S	0	0	0
			1487	472	728	131	149	7			
2	R	95	Total	C	H	N	O	S	0	0	0
			1502	475	739	132	149	7			

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	T	95	Total	C	H	N	O	S	0	1	0
			1503	475	740	132	149	7			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	U	2	Total	C	H	N	O		0	0	0
			40	16	12	2	10				
3	Y	2	Total	C	H	N	O		0	0	0
			53	16	25	2	10				
3	Z	2	Total	C	H	N	O		0	0	0
			53	16	25	2	10				
3	b	2	Total	C	H	N	O		0	0	0
			53	16	25	2	10				

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
4	V	3	Total	C	H	N	O		0	0	0
			72	22	34	2	14				
4	W	3	Total	C	H	N	O		0	0	0
			72	22	34	2	14				
4	X	3	Total	C	H	N	O		0	0	0
			72	22	34	2	14				

- Molecule 5 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	a	2	Total	C	H	N	O	0	0	0
			46	14	22	1	9			

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	B	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
6	E	1	Total	C	N	O		0	0
			14	8	1	5			
6	G	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
6	I	1	Total	C	N	O		0	0
			14	8	1	5			
6	K	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
6	L	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
6	M	1	Total	C	N	O		0	0
			14	8	1	5			
6	R	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
6	S	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
6	T	1	Total	C	N	O		0	0
			14	8	1	5			


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	3	Total 3	O 3	0	0
7	C	2	Total 2	O 2	0	0
7	E	3	Total 3	O 3	0	0
7	G	2	Total 2	O 2	0	0
7	I	1	Total 1	O 1	0	0
7	K	2	Total 2	O 2	0	0
7	M	1	Total 1	O 1	0	0
7	O	1	Total 1	O 1	0	0
7	Q	2	Total 2	O 2	0	0
7	S	2	Total 2	O 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GDNF family receptor alpha




- Molecule 1: GDNF family receptor alpha



- Molecule 1: GDNF family receptor alpha





- Molecule 1: GDNF family receptor alpha







- Molecule 1: GDNF family receptor alpha




- Molecule 1: GDNF family receptor alpha



- Molecule 1: GDNF family receptor alpha




- Molecule 1: GDNF family receptor alpha





- Molecule 1: GDNF family receptor alpha







- Molecule 1: GDNF family receptor alpha




- Molecule 2: Glial cell line-derived neurotrophic factor




- Molecule 2: Glial cell line-derived neurotrophic factor




- Molecule 2: Glial cell line-derived neurotrophic factor




- Molecule 2: Glial cell line-derived neurotrophic factor




- Molecule 2: Glial cell line-derived neurotrophic factor




- Molecule 2: Glial cell line-derived neurotrophic factor





- Molecule 2: Glial cell line-derived neurotrophic factor




- Molecule 2: Glial cell line-derived neurotrophic factor




- Molecule 2: Glial cell line-derived neurotrophic factor



- Molecule 2: Glial cell line-derived neurotrophic factor



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain Y:  50% 50%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain Z:  50% 50%



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain b:  50% 50%




- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain V:  100%



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain W:  100%



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain X:  33% 67%



- Molecule 5: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain a:  100%

MCI
FUC2

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	114.14Å 170.04Å 130.79Å 90.00° 96.17° 90.00°	Depositor
Resolution (Å)	90.45 – 2.66 90.45 – 2.66	Depositor EDS
% Data completeness (in resolution range)	88.1 (90.45-2.66) 88.3 (90.45-2.66)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.07 (at 2.65Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.244 , 0.283 0.245 , 0.253	Depositor DCC
R_{free} test set	3869 reflections (2.74%)	wwPDB-VP
Wilson B-factor (Å ²)	75.8	Xtriage
Anisotropy	0.414	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 59.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	46489	wwPDB-VP
Average B, all atoms (Å ²)	90.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FUC, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.10	0/1628	0.25	0/2196
1	C	0.09	0/1605	0.24	0/2166
1	E	0.10	0/1610	0.24	0/2173
1	G	0.10	0/1608	0.24	0/2170
1	I	0.10	0/1586	0.25	0/2139
1	K	0.10	0/1589	0.23	0/2145
1	M	0.10	0/1605	0.25	0/2166
1	O	0.10	0/1605	0.24	0/2166
1	Q	0.11	0/1596	0.25	0/2154
1	S	0.11	0/1599	0.25	0/2158
2	B	0.21	0/775	0.35	1/1046 (0.1%)
2	D	0.09	0/797	0.24	0/1073
2	F	0.09	0/795	0.24	0/1070
2	H	0.10	0/782	0.25	0/1053
2	J	0.09	0/791	0.25	0/1065
2	L	0.08	0/778	0.25	0/1049
2	N	0.09	0/777	0.24	0/1046
2	P	0.08	0/773	0.24	0/1042
2	R	0.08	0/777	0.24	0/1046
2	T	0.08	0/784	0.26	0/1056
All	All	0.10	0/23860	0.25	1/32179 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	145	LYS	CA-CB-CG	5.55	125.20	114.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1598	1530	1533	13	0
1	C	1575	1496	1511	7	0
1	E	1580	1507	1510	12	0
1	G	1578	1495	1512	13	0
1	I	1557	1498	1498	13	0
1	K	1560	1483	1488	14	0
1	M	1575	1511	1511	13	0
1	O	1575	1509	1507	15	0
1	Q	1567	1494	1497	13	0
1	S	1569	1491	1506	21	0
2	B	761	720	720	9	0
2	D	783	752	752	7	0
2	F	781	741	752	8	0
2	H	768	732	741	5	0
2	J	777	741	746	11	0
2	L	764	729	730	17	0
2	N	763	739	739	10	0
2	P	759	728	728	9	0
2	R	763	739	739	5	0
2	T	763	740	732	11	0
3	U	28	12	25	1	0
3	Y	28	25	25	4	0
3	Z	28	25	25	2	0
3	b	28	25	25	1	0
4	V	38	34	34	0	0
4	W	38	34	34	0	0
4	X	38	34	34	7	0
5	a	24	22	22	0	0
6	B	14	13	13	0	0
6	E	14	0	13	1	0
6	G	14	13	13	1	0
6	I	14	0	13	4	0
6	K	14	13	13	1	0
6	L	14	13	13	1	0
6	M	14	0	13	6	0
6	R	14	13	13	1	0
6	S	14	13	13	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	T	14	0	13	2	0
7	A	3	0	0	0	0
7	C	2	0	0	0	0
7	E	3	0	0	0	0
7	G	2	0	0	0	0
7	I	1	0	0	0	0
7	K	2	0	0	0	0
7	M	1	0	0	0	0
7	O	1	0	0	0	0
7	Q	2	0	0	0	0
7	S	2	0	0	0	0
All	All	23825	22664	22806	225	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 225 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:I:401:NAG:H83	1:S:323:ASN:HB2	1.18	1.10
6:I:401:NAG:O3	6:I:401:NAG:H82	1.70	0.90
6:I:401:NAG:H83	1:S:323:ASN:CB	2.03	0.89
6:M:401:NAG:H3	6:M:401:NAG:H83	1.55	0.89
6:I:401:NAG:C8	1:S:323:ASN:HB2	2.02	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	205/260 (79%)	194 (95%)	11 (5%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	202/260 (78%)	191 (95%)	11 (5%)	0	100	100
1	E	203/260 (78%)	198 (98%)	5 (2%)	0	100	100
1	G	202/260 (78%)	193 (96%)	9 (4%)	0	100	100
1	I	197/260 (76%)	193 (98%)	4 (2%)	0	100	100
1	K	198/260 (76%)	194 (98%)	4 (2%)	0	100	100
1	M	202/260 (78%)	198 (98%)	3 (2%)	1 (0%)	24	39
1	O	203/260 (78%)	194 (96%)	9 (4%)	0	100	100
1	Q	199/260 (76%)	194 (98%)	5 (2%)	0	100	100
1	S	201/260 (77%)	195 (97%)	6 (3%)	0	100	100
2	B	95/99 (96%)	93 (98%)	2 (2%)	0	100	100
2	D	96/99 (97%)	92 (96%)	4 (4%)	0	100	100
2	F	96/99 (97%)	94 (98%)	2 (2%)	0	100	100
2	H	94/99 (95%)	92 (98%)	2 (2%)	0	100	100
2	J	96/99 (97%)	92 (96%)	4 (4%)	0	100	100
2	L	94/99 (95%)	90 (96%)	4 (4%)	0	100	100
2	N	93/99 (94%)	88 (95%)	5 (5%)	0	100	100
2	P	93/99 (94%)	89 (96%)	4 (4%)	0	100	100
2	R	93/99 (94%)	87 (94%)	6 (6%)	0	100	100
2	T	94/99 (95%)	91 (97%)	3 (3%)	0	100	100
All	All	2956/3590 (82%)	2852 (96%)	103 (4%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	M	322	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/232 (79%)	176 (96%)	8 (4%)	26	44
1	C	182/232 (78%)	175 (96%)	7 (4%)	29	49
1	E	181/232 (78%)	177 (98%)	4 (2%)	45	67
1	G	182/232 (78%)	178 (98%)	4 (2%)	45	67
1	I	180/232 (78%)	172 (96%)	8 (4%)	25	43
1	K	179/232 (77%)	172 (96%)	7 (4%)	28	48
1	M	182/232 (78%)	175 (96%)	7 (4%)	29	49
1	O	180/232 (78%)	174 (97%)	6 (3%)	33	55
1	Q	181/232 (78%)	170 (94%)	11 (6%)	17	30
1	S	181/232 (78%)	174 (96%)	7 (4%)	28	48
2	B	85/90 (94%)	82 (96%)	3 (4%)	32	52
2	D	89/90 (99%)	88 (99%)	1 (1%)	65	80
2	F	89/90 (99%)	87 (98%)	2 (2%)	45	67
2	H	88/90 (98%)	86 (98%)	2 (2%)	44	66
2	J	88/90 (98%)	85 (97%)	3 (3%)	32	53
2	L	87/90 (97%)	85 (98%)	2 (2%)	44	66
2	N	88/90 (98%)	84 (96%)	4 (4%)	24	42
2	P	87/90 (97%)	84 (97%)	3 (3%)	32	53
2	R	88/90 (98%)	86 (98%)	2 (2%)	44	66
2	T	89/90 (99%)	85 (96%)	4 (4%)	24	42
All	All	2690/3220 (84%)	2595 (96%)	95 (4%)	32	52

5 of 95 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	M	320	SER
1	Q	171	SER
2	N	188	THR
1	O	221	GLN
1	Q	229	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	M	284	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	R	222	HIS
1	Q	323	ASN
1	S	207	HIS
2	D	148	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

19 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	U	1	3,1	14,14,15	0.95	2 (14%)	17,19,21	1.26	2 (11%)
3	NAG	U	2	3	14,14,15	0.27	0	17,19,21	0.52	0
4	NAG	V	1	2,4	14,14,15	0.26	0	17,19,21	0.39	0
4	NAG	V	2	4	14,14,15	0.26	0	17,19,21	0.55	0
4	FUC	V	3	4	10,10,11	0.71	0	14,14,16	0.77	0
4	NAG	W	1	2,4	14,14,15	0.21	0	17,19,21	0.36	0
4	NAG	W	2	4	14,14,15	0.21	0	17,19,21	0.44	0
4	FUC	W	3	4	10,10,11	0.82	0	14,14,16	0.82	0
4	NAG	X	1	2,4	14,14,15	0.38	0	17,19,21	0.96	1 (5%)
4	NAG	X	2	4	14,14,15	0.35	0	17,19,21	1.08	2 (11%)
4	FUC	X	3	4	10,10,11	0.96	0	14,14,16	0.75	0
3	NAG	Y	1	3,2	14,14,15	0.41	0	17,19,21	0.47	0
3	NAG	Y	2	3	14,14,15	0.39	0	17,19,21	0.87	1 (5%)
3	NAG	Z	1	3,2	14,14,15	0.20	0	17,19,21	0.45	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	Z	2	3	14,14,15	0.22	0	17,19,21	0.54	0
5	NAG	a	1	5,1	14,14,15	0.32	0	17,19,21	0.58	0
5	FUC	a	2	5	10,10,11	0.82	0	14,14,16	0.80	0
3	NAG	b	1	3,2	14,14,15	0.26	0	17,19,21	0.50	0
3	NAG	b	2	3	14,14,15	0.30	0	17,19,21	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	U	1	3,1	-	4/6/23/26	0/1/1/1
3	NAG	U	2	3	-	3/6/23/26	0/1/1/1
4	NAG	V	1	2,4	-	0/6/23/26	0/1/1/1
4	NAG	V	2	4	-	0/6/23/26	0/1/1/1
4	FUC	V	3	4	-	-	0/1/1/1
4	NAG	W	1	2,4	-	0/6/23/26	0/1/1/1
4	NAG	W	2	4	-	0/6/23/26	0/1/1/1
4	FUC	W	3	4	-	-	0/1/1/1
4	NAG	X	1	2,4	-	3/6/23/26	0/1/1/1
4	NAG	X	2	4	-	4/6/23/26	0/1/1/1
4	FUC	X	3	4	-	-	0/1/1/1
3	NAG	Y	1	3,2	-	2/6/23/26	0/1/1/1
3	NAG	Y	2	3	-	3/6/23/26	0/1/1/1
3	NAG	Z	1	3,2	-	0/6/23/26	0/1/1/1
3	NAG	Z	2	3	-	0/6/23/26	0/1/1/1
5	NAG	a	1	5,1	-	2/6/23/26	0/1/1/1
5	FUC	a	2	5	-	-	0/1/1/1
3	NAG	b	1	3,2	-	2/6/23/26	0/1/1/1
3	NAG	b	2	3	-	3/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	U	1	NAG	O5-C1	2.46	1.47	1.43
3	U	1	NAG	C1-C2	2.43	1.55	1.52

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	U	1	NAG	C1-O5-C5	3.68	117.11	112.19
4	X	2	NAG	C2-N2-C7	2.95	126.86	122.90
3	Y	2	NAG	C2-N2-C7	2.63	126.43	122.90
4	X	1	NAG	C2-N2-C7	2.54	126.31	122.90
4	X	2	NAG	C1-O5-C5	2.35	115.33	112.19

There are no chirality outliers.

5 of 26 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	Y	2	NAG	C1-C2-N2-C7
4	X	1	NAG	C1-C2-N2-C7
5	a	1	NAG	C4-C5-C6-O6
5	a	1	NAG	O5-C5-C6-O6
3	U	1	NAG	C4-C5-C6-O6


There are no ring outliers.


8 monomers are involved in 15 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	Z	1	NAG	2	0
3	Y	1	NAG	1	0
4	X	3	FUC	1	0
4	X	2	NAG	4	0
3	b	1	NAG	1	0
3	Y	2	NAG	3	0
4	X	1	NAG	4	0
3	U	2	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

















5.6 Ligand geometry

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	I	401	1	14,14,15	0.27	0	17,19,21	0.64	0
6	NAG	B	301	2	14,14,15	0.29	0	17,19,21	0.42	0
6	NAG	R	301	2	14,14,15	0.35	0	17,19,21	0.57	0
6	NAG	S	401	1	14,14,15	0.65	1 (7%)	17,19,21	0.82	2 (11%)
6	NAG	E	401	1	14,14,15	0.37	0	17,19,21	0.53	0
6	NAG	T	301	2	14,14,15	0.37	0	17,19,21	0.53	0
6	NAG	G	401	1	14,14,15	0.42	0	17,19,21	0.62	0
6	NAG	M	401	1	14,14,15	0.33	0	17,19,21	0.55	0
6	NAG	L	301	2	14,14,15	0.47	0	17,19,21	0.81	1 (5%)
6	NAG	K	401	1	14,14,15	0.39	0	17,19,21	0.66	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	I	401	1	-	4/6/23/26	0/1/1/1
6	NAG	B	301	2	-	1/6/23/26	0/1/1/1
6	NAG	R	301	2	-	1/6/23/26	0/1/1/1
6	NAG	S	401	1	-	2/6/23/26	0/1/1/1
6	NAG	E	401	1	-	3/6/23/26	0/1/1/1
6	NAG	T	301	2	-	2/6/23/26	0/1/1/1
6	NAG	G	401	1	-	2/6/23/26	0/1/1/1
6	NAG	M	401	1	-	6/6/23/26	0/1/1/1
6	NAG	L	301	2	-	2/6/23/26	0/1/1/1
6	NAG	K	401	1	-	2/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	S	401	NAG	C1-C2	2.16	1.55	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	L	301	NAG	C1-O5-C5	2.96	116.15	112.19
6	S	401	NAG	C2-N2-C7	2.15	125.78	122.90
6	S	401	NAG	C1-O5-C5	2.04	114.92	112.19

There are no chirality outliers.

5 of 25 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	I	401	NAG	O5-C5-C6-O6
6	E	401	NAG	O5-C5-C6-O6
6	G	401	NAG	C8-C7-N2-C2
6	G	401	NAG	O7-C7-N2-C2
6	I	401	NAG	C8-C7-N2-C2

There are no ring outliers.

9 monomers are involved in 21 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	I	401	NAG	4	0
6	R	301	NAG	1	0
6	S	401	NAG	4	0
6	E	401	NAG	1	0
6	T	301	NAG	2	0
6	G	401	NAG	1	0
6	M	401	NAG	6	0
6	L	301	NAG	1	0
6	K	401	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	207/260 (79%)	0.16	2 (0%) 79 76	64, 75, 109, 129	0
1	C	204/260 (78%)	0.19	5 (2%) 58 52	65, 78, 111, 128	0
1	E	205/260 (78%)	0.24	5 (2%) 59 53	71, 83, 119, 150	0
1	G	204/260 (78%)	0.24	7 (3%) 48 40	66, 81, 109, 120	0
1	I	201/260 (77%)	0.31	6 (2%) 52 45	67, 84, 113, 148	0
1	K	202/260 (77%)	0.32	9 (4%) 38 31	75, 91, 125, 147	0
1	M	204/260 (78%)	0.29	6 (2%) 53 46	75, 86, 111, 129	0
1	O	205/260 (78%)	0.32	11 (5%) 31 25	71, 86, 118, 139	0
1	Q	203/260 (78%)	0.41	13 (6%) 25 19	76, 92, 133, 145	0
1	S	203/260 (78%)	0.33	7 (3%) 48 40	67, 85, 121, 137	0
2	B	97/99 (97%)	0.32	3 (3%) 51 43	71, 95, 123, 127	0
2	D	98/99 (98%)	0.09	2 (2%) 65 59	70, 82, 115, 127	0
2	F	98/99 (98%)	0.27	1 (1%) 79 76	75, 85, 101, 117	0
2	H	96/99 (96%)	0.29	3 (3%) 51 43	72, 91, 111, 135	0
2	J	98/99 (98%)	0.15	2 (2%) 65 59	78, 88, 112, 130	0
2	L	96/99 (96%)	0.45	4 (4%) 40 32	87, 111, 138, 148	0
2	N	95/99 (95%)	0.20	3 (3%) 50 42	79, 93, 113, 126	0
2	P	95/99 (95%)	0.19	2 (2%) 63 57	74, 88, 109, 115	0
2	R	95/99 (95%)	0.35	0 100 100	83, 114, 132, 153	0
2	T	95/99 (95%)	0.36	4 (4%) 40 32	82, 103, 124, 131	0
All	All	3001/3590 (83%)	0.28	95 (3%) 50 42	64, 87, 121, 153	0

The worst 5 of 95 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	323	ASN	6.4
1	A	355	GLU	5.5
1	O	354	SER	4.5
1	Q	290	LEU	4.2
1	K	218	LEU	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	U	2	14/15	0.36	0.19	93,95,97,97	0
4	NAG	W	2	14/15	0.37	0.15	82,87,103,105	0
3	NAG	Y	2	14/15	0.49	0.12	89,91,110,110	0
4	FUC	X	3	10/11	0.49	0.15	80,83,100,100	0
4	FUC	W	3	10/11	0.51	0.15	83,87,104,105	0
4	FUC	V	3	10/11	0.55	0.16	80,86,104,104	0
3	NAG	b	1	14/15	-	-	80,84,101,101	0
3	NAG	b	2	14/15	-	-	83,86,103,105	0
3	NAG	Z	2	14/15	0.55	0.14	83,86,102,104	0
3	NAG	U	1	14/15	0.59	0.16	86,100,121,121	0
4	NAG	X	2	14/15	0.63	0.10	80,82,97,99	0
3	NAG	Z	1	14/15	0.66	0.13	81,85,102,102	0
4	NAG	V	2	14/15	0.75	0.10	76,85,102,107	0
4	NAG	X	1	14/15	0.77	0.11	76,80,97,97	0
4	NAG	W	1	14/15	0.78	0.09	82,87,103,104	0
3	NAG	Y	1	14/15	0.85	0.10	84,89,106,106	0
4	NAG	V	1	14/15	0.90	0.08	69,78,93,93	0
5	NAG	a	1	14/15	-	-	92,105,121,126	0
5	FUC	a	2	10/11	-	-	100,105,126,127	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain U:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain Y:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain Z:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain b:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain V:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around Chain W:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain X:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	NAG	K	401	14/15	0.37	0.21	96,103,123,125	0
6	NAG	I	401	14/15	0.38	0.20	105,112,117,117	0
6	NAG	S	401	14/15	0.50	0.15	95,104,123,125	0
6	NAG	B	301	14/15	0.55	0.13	82,86,102,103	0
6	NAG	R	301	14/15	0.57	0.12	98,106,128,128	0
6	NAG	E	401	14/15	0.57	0.15	95,101,103,105	0
6	NAG	T	301	14/15	0.57	0.18	92,96,97,98	0
6	NAG	G	401	14/15	0.68	0.15	84,93,117,117	0
6	NAG	M	401	14/15	0.69	0.16	95,99,105,105	0
6	NAG	L	301	14/15	0.72	0.11	96,99,119,120	0

6.5 Other polymers [i](#)

There are no such residues in this entry.