



Full wwPDB EM Validation Report ⓘ

Mar 9, 2026 – 10:55 PM UTC

PDB ID : 8OSK / pdb_00008osk
EMDB ID : EMD-17157
Title : Cryo-EM structure of CLOCK-BMAL1 bound to a nucleosomal E-box at position SHL+5.8 (composite map)
Authors : Stoos, L.; Michael, A.K.; Kempf, G.; Cavadini, S.; Thoma, N.H.
Deposited on : 2023-04-19
Resolution : 3.60 Å (reported)
Based on initial models : 4H10, 4F3L

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	12797 (3.10 - 4.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	139	66% 32%
1	E	139	65% 32%
2	B	106	75% 25%
2	F	106	75% 25%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	133	
3	G	133	
4	D	128	
4	H	128	
5	I	153	
6	J	153	
7	M	375	
8	N	384	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 13995 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.1.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	94	Total	C	N	O	S	0	0
			773	488	147	134	4		
1	E	94	Total	C	N	O	S	0	0
			773	488	147	134	4		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP P68431
A	-2	SER	-	expression tag	UNP P68431
A	-1	HIS	-	expression tag	UNP P68431
E	-3	GLY	-	expression tag	UNP P68431
E	-2	SER	-	expression tag	UNP P68431
E	-1	HIS	-	expression tag	UNP P68431

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	80	Total	C	N	O	S	0	0
			641	405	125	110	1		
2	F	80	Total	C	N	O	S	0	0
			641	405	125	110	1		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P62805
B	-2	SER	-	expression tag	UNP P62805
B	-1	HIS	-	expression tag	UNP P62805
F	-3	GLY	-	expression tag	UNP P62805
F	-2	SER	-	expression tag	UNP P62805
F	-1	HIS	-	expression tag	UNP P62805

- Molecule 3 is a protein called Histone H2A type 1-B/E.

Mol	Chain	Residues	Atoms				AltConf	Trace
3	C	104	Total	C	N	O	0	0
			805	508	157	140		
3	G	104	Total	C	N	O	0	0
			805	508	157	140		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	GLY	-	expression tag	UNP P04908
C	-2	SER	-	expression tag	UNP P04908
C	-1	HIS	-	expression tag	UNP P04908
G	-3	GLY	-	expression tag	UNP P04908
G	-2	SER	-	expression tag	UNP P04908
G	-1	HIS	-	expression tag	UNP P04908

- Molecule 4 is a protein called Histone H2B type 1-J.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	92	Total	C	N	O	S	0	0
			719	453	129	135	2		
4	H	94	Total	C	N	O	S	0	0
			736	462	134	138	2		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-3	GLY	-	expression tag	UNP P06899
D	-2	SER	-	expression tag	UNP P06899
D	-1	HIS	-	expression tag	UNP P06899
H	-3	GLY	-	expression tag	UNP P06899
H	-2	SER	-	expression tag	UNP P06899
H	-1	HIS	-	expression tag	UNP P06899

- Molecule 5 is a DNA chain called DNA (124-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
5	I	124	Total	C	N	O	P	0	0
			2520	1197	453	746	124		

- Molecule 6 is a DNA chain called DNA (124-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace
6	J	124	Total	C	N	O	P	0	0
			2564	1211	487	742	124		

- Molecule 7 is a protein called Circadian locomoter output cycles protein kaput.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	M	316	Total	C	N	O		0	0
			1591	959	316	316			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	21	GLY	-	expression tag	UNP O08785
M	22	ALA	-	expression tag	UNP O08785
M	23	MET	-	expression tag	UNP O08785
M	24	ASN	-	expression tag	UNP O08785
M	25	PRO	-	expression tag	UNP O08785

- Molecule 8 is a protein called Basic helix-loop-helix ARNT-like protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	N	285	Total	C	N	O		0	0
			1427	857	285	285			

There are 5 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
N	64	GLY	-	expression tag	UNP Q9WTL8
N	65	ALA	-	expression tag	UNP Q9WTL8
N	66	MET	-	expression tag	UNP Q9WTL8
N	67	ASN	-	expression tag	UNP Q9WTL8
N	68	PRO	-	expression tag	UNP Q9WTL8

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Histone H3.1

Chain A: 




- Molecule 1: Histone H3.1

Chain E: 




- Molecule 2: Histone H4

Chain B: 



- Molecule 2: Histone H4

Chain F: 




- Molecule 3: Histone H2A type 1-B/E

Chain C: 



- Molecule 3: Histone H2A type 1-B/E



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	25483	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	28.349	Depositor
Minimum map value	-9.143	Depositor
Average map value	-0.004	Depositor
Map value standard deviation	0.951	Depositor
Recommended contour level	2.9	Depositor
Map size (Å)	304.2, 304.2, 304.2	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.845, 0.845, 0.845	Depositor

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/783	0.78	4/1050 (0.4%)
1	E	0.62	0/783	0.79	6/1050 (0.6%)
2	B	0.61	0/648	0.79	2/868 (0.2%)
2	F	0.63	0/648	0.76	2/868 (0.2%)
3	C	0.60	0/815	0.84	6/1100 (0.5%)
3	G	0.59	0/815	0.86	8/1100 (0.7%)
4	D	0.57	0/730	0.79	4/982 (0.4%)
4	H	0.59	0/747	0.81	4/1004 (0.4%)
5	I	0.31	0/2821	0.76	0/4346
6	J	0.31	0/2881	0.72	0/4450
7	M	0.41	0/1600	0.98	22/2237 (1.0%)
8	N	0.40	0/1429	0.90	12/1991 (0.6%)
All	All	0.47	0/14700	0.81	70/21046 (0.3%)

There are no bond length outliers.

All (70) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	102	LEU	CA-C-N	7.54	127.54	119.78
4	D	102	LEU	C-N-CA	7.54	127.54	119.78
4	H	49	HIS	CA-C-N	7.24	126.95	119.56
4	H	49	HIS	C-N-CA	7.24	126.95	119.56
7	M	203	ASP	CA-C-N	7.13	127.29	119.87
7	M	203	ASP	C-N-CA	7.13	127.29	119.87
8	N	331	VAL	CA-C-N	7.11	127.07	120.03
8	N	331	VAL	C-N-CA	7.11	127.07	120.03
4	H	102	LEU	CA-C-N	7.02	127.01	119.78
4	H	102	LEU	C-N-CA	7.02	127.01	119.78
7	M	181	THR	CA-C-N	7.01	126.98	119.76
7	M	181	THR	C-N-CA	7.01	126.98	119.76
3	C	116	LEU	CA-C-N	6.91	126.83	119.85
3	C	116	LEU	C-N-CA	6.91	126.83	119.85
3	G	116	LEU	CA-C-N	6.79	126.55	119.76

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	116	LEU	C-N-CA	6.79	126.55	119.76
8	N	431	ASN	CA-C-N	6.71	126.33	119.56
8	N	431	ASN	C-N-CA	6.71	126.33	119.56
7	M	61	LEU	CA-C-N	6.50	127.03	120.14
7	M	61	LEU	C-N-CA	6.50	127.03	120.14
4	D	49	HIS	CA-C-N	6.37	126.29	119.28
4	D	49	HIS	C-N-CA	6.37	126.29	119.28
7	M	259	THR	CA-C-N	6.32	126.23	119.85
7	M	259	THR	C-N-CA	6.32	126.23	119.85
8	N	333	GLN	CA-C-N	6.32	126.28	119.78
8	N	333	GLN	C-N-CA	6.32	126.28	119.78
3	C	25	PHE	CA-C-N	6.30	127.71	119.84
3	C	25	PHE	C-N-CA	6.30	127.71	119.84
7	M	158	ILE	CA-C-N	6.09	126.03	119.76
7	M	158	ILE	C-N-CA	6.09	126.03	119.76
8	N	204	HIS	CA-C-N	6.07	126.51	119.47
8	N	204	HIS	C-N-CA	6.07	126.51	119.47
7	M	206	GLU	CA-C-N	6.01	125.97	119.78
7	M	206	GLU	C-N-CA	6.01	125.97	119.78
3	G	25	PHE	CA-C-N	5.93	127.26	119.84
3	G	25	PHE	C-N-CA	5.93	127.26	119.84
7	M	294	PRO	CA-C-N	5.88	125.56	119.56
7	M	294	PRO	C-N-CA	5.88	125.56	119.56
3	G	47	ALA	CA-C-N	5.88	125.58	119.05
3	G	47	ALA	C-N-CA	5.88	125.58	119.05
8	N	106	VAL	CA-C-N	5.87	126.01	119.32
8	N	106	VAL	C-N-CA	5.87	126.01	119.32
1	E	42	ARG	CA-C-N	5.86	125.79	119.76
1	E	42	ARG	C-N-CA	5.86	125.79	119.76
7	M	365	ARG	CA-C-N	5.79	125.73	119.76
7	M	365	ARG	C-N-CA	5.79	125.73	119.76
3	C	79	ILE	CA-C-N	5.71	126.09	119.47
3	C	79	ILE	C-N-CA	5.71	126.09	119.47
2	B	31	LYS	CA-C-N	5.66	125.78	119.32
2	B	31	LYS	C-N-CA	5.66	125.78	119.32
2	F	31	LYS	CA-C-N	5.65	125.76	119.32
2	F	31	LYS	C-N-CA	5.65	125.76	119.32
3	G	79	ILE	CA-C-N	5.62	125.99	119.47
3	G	79	ILE	C-N-CA	5.62	125.99	119.47
1	A	120	MET	CA-C-N	5.52	125.88	119.47
1	A	120	MET	C-N-CA	5.52	125.88	119.47
7	M	300	LEU	CA-C-N	5.49	125.58	119.32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	M	300	LEU	C-N-CA	5.49	125.58	119.32
1	A	65	LEU	CA-C-N	5.43	125.51	119.32
1	A	65	LEU	C-N-CA	5.43	125.51	119.32
1	E	120	MET	CA-C-N	5.39	125.72	119.47
1	E	120	MET	C-N-CA	5.39	125.72	119.47
7	M	271	GLU	CA-C-N	5.39	125.46	119.32
7	M	271	GLU	C-N-CA	5.39	125.46	119.32
1	E	65	LEU	CA-C-N	5.36	125.00	119.05
1	E	65	LEU	C-N-CA	5.36	125.00	119.05
7	M	145	LEU	CA-C-N	5.24	125.55	119.47
7	M	145	LEU	C-N-CA	5.24	125.55	119.47
8	N	370	LEU	CA-C-N	5.24	125.30	119.32
8	N	370	LEU	C-N-CA	5.24	125.30	119.32

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	773	0	811	0	0
1	E	773	0	811	1	0
2	B	641	0	684	0	0
2	F	641	0	684	0	0
3	C	805	0	861	1	0
3	G	805	0	861	1	0
4	D	719	0	740	0	0
4	H	736	0	758	0	0
5	I	2520	0	1392	1	0
6	J	2564	0	1390	0	0
7	M	1591	0	735	4	0
8	N	1427	0	660	2	0
All	All	13995	0	10387	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (10) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:47:ALA:N	3:C:48:PRO:HD2	2.25	0.52
3:G:47:ALA:N	3:G:48:PRO:CD	2.72	0.51
7:M:293:ALA:N	7:M:294:PRO:CD	2.78	0.46
8:N:367:LEU:C	8:N:369:TYR:H	2.23	0.46
1:E:65:LEU:N	1:E:66:PRO:CD	2.79	0.46
7:M:294:PRO:N	7:M:295:PRO:CD	2.80	0.44
7:M:294:PRO:HB2	7:M:295:PRO:HD3	1.98	0.44
7:M:293:ALA:HB3	7:M:294:PRO:HD3	2.02	0.42
8:N:204:HIS:HA	8:N:205:PRO:HD3	1.93	0.42
5:I:132:DC:H2'	5:I:133:DG:C8	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	92/139 (66%)	92 (100%)	0	0	100	100
1	E	92/139 (66%)	92 (100%)	0	0	100	100
2	B	78/106 (74%)	78 (100%)	0	0	100	100
2	F	78/106 (74%)	78 (100%)	0	0	100	100
3	C	102/133 (77%)	102 (100%)	0	0	100	100
3	G	102/133 (77%)	101 (99%)	1 (1%)	0	100	100
4	D	90/128 (70%)	90 (100%)	0	0	100	100
4	H	92/128 (72%)	92 (100%)	0	0	100	100
7	M	310/375 (83%)	309 (100%)	1 (0%)	0	100	100
8	N	275/384 (72%)	273 (99%)	2 (1%)	0	100	100
All	All	1311/1771 (74%)	1307 (100%)	4 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	82/113 (73%)	82 (100%)	0	100	100
1	E	82/113 (73%)	82 (100%)	0	100	100
2	B	66/81 (82%)	66 (100%)	0	100	100
2	F	66/81 (82%)	66 (100%)	0	100	100
3	C	83/102 (81%)	83 (100%)	0	100	100
3	G	83/102 (81%)	83 (100%)	0	100	100
4	D	78/106 (74%)	78 (100%)	0	100	100
4	H	80/106 (76%)	80 (100%)	0	100	100
7	M	12/345 (4%)	12 (100%)	0	100	100
8	N	7/342 (2%)	7 (100%)	0	100	100
All	All	639/1491 (43%)	639 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
4	H	95	GLN
4	H	109	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-17157. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 180



Y Index: 180




Z Index: 180


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 162



Y Index: 193




Z Index: 188


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 2.9. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 333 nm^3 ; this corresponds to an approximate mass of 301 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.278 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-17157 and PDB model 8OSK. Per-residue inclusion information can be found in section 3 on page 7.


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 2.9 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (2.9).

9.4 Atom inclusion ⓘ



At the recommended contour level, 94% of all backbone atoms, 93% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (2.9) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.9290	 0.2670
A	 0.9250	 0.3220
B	 0.9400	 0.3480
C	 0.9400	 0.3280
D	 0.9690	 0.3280
E	 0.9440	 0.3140
F	 0.9420	 0.3510
G	 0.8830	 0.3310
H	 0.9120	 0.3290
I	 0.9830	 0.3010
J	 0.9800	 0.2930
M	 0.8270	 0.1130
N	 0.8470	 0.0710

