



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:05 PM UTC

PDB ID : 2OTH / pdb_00002oth
Title : Crystal structure of a ternary complex of phospholipase A2 with indomethacin and nimesulide at 2.9 Å resolution
Authors : Kumar, S.; Singh, N.; Sharma, S.; Kaur, P.; Singh, T.P.
Deposited on : 2007-02-08
Resolution : 2.90 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	121	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NIM	A	300	-	X	X	-
3	IMN	A	301	-	X	-	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 1041 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phospholipase A2 VRV-PL-VIIIa.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	121	944	591	159	178	16	0	0	0

- Molecule 2 is 4-NITRO-2-PHENOXYMETHANESULFONANILIDE (CCD ID: NIM) (formula: C₁₃H₁₂N₂O₅S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	21	13	2	5	1	0	0

- Molecule 3 is INDOMETHACIN (CCD ID: IMN) (formula: C₁₉H₁₆ClNO₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Cl	N			O
3	A	1	25	19	1	1	4	0	0

- Molecule 4 is ACETONITRILE (CCD ID: CCN) (formula: C_2H_3N).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	C N		
4	A	1	3	2 1	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	48	Total 48	O 48	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Phospholipase A2 VRV-PL-VIIIa



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 43	Depositor
Cell constants a, b, c, α , β , γ	52.88Å 52.88Å 48.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	17.44 – 2.90	Depositor
% Data completeness (in resolution range)	96.7 (17.44-2.90)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 0.9	Depositor
R, R_{free}	0.189 , 0.232	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1041	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NIM, CCN, IMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	0/964	1.18	7/1292 (0.5%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	111	ASN	CA-CB-CG	-9.30	103.30	112.60
1	A	35	GLY	N-CA-C	8.24	122.24	111.70
1	A	34	LYS	N-CA-C	6.84	119.78	110.35
1	A	19	ILE	N-CA-C	-6.71	100.13	108.05
1	A	39	ASP	N-CA-C	6.55	117.22	108.38
1	A	37	PRO	CB-CA-C	5.30	118.37	111.64
1	A	122	ASP	N-CA-C	5.00	116.73	111.28

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	944	0	909	41	0
2	A	21	0	12	8	0
3	A	25	0	16	2	0
4	A	3	0	3	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	48	0	0	2	0
All	All	1041	0	940	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:69:LYS:CE	2:A:300:NIM:H11A	1.89	1.03
1:A:69:LYS:HE3	2:A:300:NIM:H11A	1.03	1.00
1:A:69:LYS:HE3	2:A:300:NIM:C1	1.98	0.90
3:A:301:IMN:H5	3:A:301:IMN:H15	1.65	0.78
1:A:28:TYR:CE1	1:A:37:PRO:HD3	2.23	0.73
1:A:77:ARG:HA	1:A:81:ALA:O	1.92	0.69
1:A:3:LEU:HD23	1:A:72:ARG:HH11	1.58	0.68
1:A:19:ILE:HD11	2:A:300:NIM:H6	1.74	0.68
1:A:19:ILE:HD11	2:A:300:NIM:C6	2.29	0.61
1:A:124:PHE:CE1	1:A:125:LEU:HG	2.36	0.60
1:A:38:LYS:HA	1:A:38:LYS:HE2	1.86	0.58
1:A:3:LEU:HD23	1:A:72:ARG:NH1	2.20	0.56
1:A:78:VAL:O	1:A:79:ASN:HB2	2.06	0.55
1:A:28:TYR:HE1	1:A:37:PRO:HD3	1.67	0.55
1:A:7:LYS:O	1:A:11:GLU:HG3	2.06	0.55
1:A:1:SER:HA	1:A:68:PRO:O	2.07	0.54
1:A:78:VAL:O	1:A:78:VAL:HG12	2.07	0.54
1:A:12:GLU:HG3	1:A:107:ARG:HB2	1.90	0.54
1:A:9:ILE:HD12	1:A:106:PHE:HD2	1.73	0.53
1:A:1:SER:HB3	1:A:71:ASP:O	2.09	0.53
1:A:9:ILE:HD12	1:A:106:PHE:CD2	2.45	0.51
1:A:37:PRO:HA	1:A:42:ASP:OD2	2.11	0.50
1:A:8:MET:HE2	1:A:82:ILE:HG21	1.94	0.50
1:A:124:PHE:CZ	1:A:125:LEU:HG	2.49	0.48
1:A:56:PRO:HD2	5:A:2080:HOH:O	2.15	0.47
1:A:5:PHE:CE1	1:A:9:ILE:HD11	2.51	0.45
1:A:113:TYR:CD1	1:A:113:TYR:C	2.95	0.44
1:A:52:TYR:CZ	1:A:68:PRO:HB3	2.53	0.44
1:A:22:TYR:O	1:A:29:CYS:HB3	2.18	0.44
3:A:301:IMN:O1	3:A:301:IMN:H161	2.17	0.44
1:A:18:ALA:HB1	2:A:300:NIM:O3	2.18	0.44
1:A:130:LEU:HD23	1:A:130:LEU:HA	1.62	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:105:CYS:SG	4:A:2067:CCN:H22	2.57	0.43
1:A:69:LYS:CE	2:A:300:NIM:C1	2.77	0.42
1:A:2:LEU:HD21	2:A:300:NIM:H13A	2.01	0.42
1:A:2:LEU:HD13	1:A:69:LYS:HA	2.02	0.42
1:A:95:ILE:O	1:A:98:CYS:HB2	2.19	0.42
1:A:2:LEU:O	1:A:3:LEU:C	2.61	0.42
1:A:67:ASN:N	5:A:2113:HOH:O	2.45	0.41
1:A:2:LEU:N	1:A:2:LEU:CD1	2.83	0.41
1:A:48:HIS:CD2	1:A:102:ALA:HB2	2.56	0.40
1:A:1:SER:HB3	1:A:71:ASP:C	2.47	0.40
1:A:1:SER:HB2	1:A:69:LYS:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	119/121 (98%)	113 (95%)	6 (5%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	104/104 (100%)	93 (89%)	11 (11%)	6 22

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LEU
1	A	16	LYS
1	A	20	PRO
1	A	31	TRP
1	A	34	LYS
1	A	74	LYS
1	A	98	CYS
1	A	118	MET
1	A	127	LYS
1	A	129	GLU
1	A	130	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	108	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	CCN	A	2067	-	2,2,2	0.36	0	1,1,1	0.60	0
3	IMN	A	301	-	27,27,27	5.54	23 (85%)	39,39,39	4.16	24 (61%)
2	NIM	A	300	-	22,22,22	6.89	19 (86%)	29,31,31	3.89	13 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IMN	A	301	-	-	10/14/14/14	0/3/3/3
2	NIM	A	300	-	-	0/11/13/13	0/2/2/2

All (42) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	300	NIM	O2-S1	17.76	1.72	1.43
2	A	300	NIM	C3-C4	13.37	1.60	1.38
2	A	300	NIM	S1-N1	10.48	1.77	1.63
3	A	301	IMN	C1-C	10.11	1.54	1.41
2	A	300	NIM	O1-S1	9.40	1.58	1.43
3	A	301	IMN	C8-C7	8.85	1.54	1.36
3	A	301	IMN	C17-C18	8.44	1.64	1.51
3	A	301	IMN	C10-C9	8.43	1.64	1.50
3	A	301	IMN	C15-C14	7.41	1.50	1.38
3	A	301	IMN	C15-C10	7.24	1.50	1.39
3	A	301	IMN	C5-C4	7.18	1.50	1.38
3	A	301	IMN	C13-CL	7.13	1.91	1.74
2	A	300	NIM	C10-C11	6.87	1.53	1.38
3	A	301	IMN	C4-C3	6.76	1.51	1.38
3	A	301	IMN	C9-N	6.56	1.56	1.41
2	A	300	NIM	C13-C8	6.32	1.50	1.38
2	A	300	NIM	C2-C7	6.11	1.53	1.40
3	A	301	IMN	C14-C13	5.72	1.48	1.38
3	A	301	IMN	C5-C	5.45	1.48	1.39
2	A	300	NIM	C10-C9	5.11	1.47	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	300	NIM	C12-C13	4.95	1.47	1.38
3	A	301	IMN	O3-C18	4.81	1.46	1.30
2	A	300	NIM	C6-C7	4.80	1.47	1.38
2	A	300	NIM	O3-N2	4.72	1.30	1.22
2	A	300	NIM	C4-C5	4.69	1.47	1.38
3	A	301	IMN	C2-C3	4.60	1.46	1.39
2	A	300	NIM	C9-C8	4.47	1.47	1.38
2	A	300	NIM	O5-C7	4.39	1.48	1.39
3	A	301	IMN	O1-C9	4.36	1.31	1.22
2	A	300	NIM	C11-C12	4.32	1.47	1.38
2	A	300	NIM	C6-C5	4.07	1.45	1.39
3	A	301	IMN	C12-C13	3.76	1.45	1.38
2	A	300	NIM	C1-S1	3.66	1.83	1.75
2	A	300	NIM	C5-N2	3.33	1.53	1.45
3	A	301	IMN	C1-C7	3.14	1.49	1.44
3	A	301	IMN	O-C3	3.08	1.43	1.37
3	A	301	IMN	C8-N	2.95	1.46	1.40
3	A	301	IMN	C17-C7	-2.92	1.43	1.50
3	A	301	IMN	O2-C18	2.90	1.31	1.22
3	A	301	IMN	C2-C1	2.81	1.44	1.39
2	A	300	NIM	O5-C8	2.62	1.45	1.39
3	A	301	IMN	O-C6	2.10	1.48	1.42

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	300	NIM	C1-S1-N1	-13.13	91.79	106.56
3	A	301	IMN	C10-C9-N	8.56	134.35	117.92
3	A	301	IMN	C17-C7-C8	8.07	139.01	126.86
2	A	300	NIM	O2-S1-C1	7.70	120.45	108.26
3	A	301	IMN	C11-C12-C13	7.60	126.89	119.24
3	A	301	IMN	C14-C13-C12	-7.31	112.19	121.24
3	A	301	IMN	C14-C13-CL	6.81	129.40	119.36
3	A	301	IMN	C17-C7-C1	-6.56	113.54	124.80
3	A	301	IMN	C5-C-C1	-6.49	114.71	121.95
2	A	300	NIM	O3-N2-C5	6.44	127.68	118.82
3	A	301	IMN	C6-O-C3	6.36	131.13	117.50
2	A	300	NIM	O2-S1-N1	6.00	119.08	107.10
3	A	301	IMN	C18-C17-C7	5.70	128.92	113.73
2	A	300	NIM	O1-S1-C1	-5.41	99.70	108.26
3	A	301	IMN	C-N-C8	5.30	112.12	107.80
3	A	301	IMN	O1-C9-C10	-5.00	110.39	120.29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	301	IMN	C16-C8-N	4.86	129.35	122.07
3	A	301	IMN	C2-C1-C	4.72	125.45	119.82
3	A	301	IMN	C15-C10-C9	4.67	132.26	120.28
2	A	300	NIM	C3-C4-C5	3.91	125.26	120.08
3	A	301	IMN	C5-C-N	3.86	138.29	130.65
2	A	300	NIM	C6-C5-N2	3.70	121.93	118.74
3	A	301	IMN	C2-C1-C7	-3.24	127.37	132.81
2	A	300	NIM	C7-C2-N1	-3.19	112.19	118.48
3	A	301	IMN	C15-C10-C11	-3.06	114.68	118.57
2	A	300	NIM	C3-C2-N1	3.00	126.72	120.92
3	A	301	IMN	O1-C9-N	-2.99	115.26	120.18
2	A	300	NIM	C4-C5-C6	-2.86	116.44	120.11
3	A	301	IMN	C11-C10-C9	-2.82	113.06	120.28
2	A	300	NIM	C4-C3-C2	-2.79	114.19	119.57
3	A	301	IMN	C7-C8-N	-2.73	106.25	108.85
2	A	300	NIM	C4-C5-N2	2.63	121.64	119.34
3	A	301	IMN	C9-N-C8	-2.52	120.20	126.55
3	A	301	IMN	C15-C14-C13	2.52	121.78	119.24
3	A	301	IMN	C12-C11-C10	2.44	123.41	120.80
3	A	301	IMN	C16-C8-C7	-2.35	124.40	128.07
2	A	300	NIM	O5-C7-C2	-2.31	111.53	116.62

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	301	IMN	C10-C9-N-C8
3	A	301	IMN	C15-C10-C9-O1
3	A	301	IMN	C11-C10-C9-O1
3	A	301	IMN	C15-C10-C9-N
3	A	301	IMN	C10-C9-N-C
3	A	301	IMN	O1-C9-N-C
3	A	301	IMN	C11-C10-C9-N
3	A	301	IMN	C2-C3-O-C6
3	A	301	IMN	C4-C3-O-C6
3	A	301	IMN	O1-C9-N-C8


There are no ring outliers.

3 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	2067	CCN	1	0
3	A	301	IMN	2	0
2	A	300	NIM	8	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.