



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 12:18 PM UTC

PDB ID : 4OT8 / pdb_00004ot8
Title : X-ray Crystal Structure of Serine Hydroxymethyl Transferase from Burkholderia cenocepacia bound to PLP and Serine
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2014-02-13
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	423	0% 92% 5% .
1	B	423	0% 92% 5% .
1	C	423	2% 95% . .
1	D	423	92% . .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13390 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine hydroxymethyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	414	Total	C	N	O	S	0	5	0
			3111	1963	543	593	12			
1	B	411	Total	C	N	O	S	0	6	0
			3102	1960	542	588	12			
1	C	415	Total	C	N	O	S	0	4	0
			3124	1971	549	591	13			
1	D	408	Total	C	N	O	S	0	4	0
			3072	1938	540	582	12			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	expression tag	UNP B4ECY9
A	-6	ALA	-	expression tag	UNP B4ECY9
A	-5	HIS	-	expression tag	UNP B4ECY9
A	-4	HIS	-	expression tag	UNP B4ECY9
A	-3	HIS	-	expression tag	UNP B4ECY9
A	-2	HIS	-	expression tag	UNP B4ECY9
A	-1	HIS	-	expression tag	UNP B4ECY9
A	0	HIS	-	expression tag	UNP B4ECY9
B	-7	MET	-	expression tag	UNP B4ECY9
B	-6	ALA	-	expression tag	UNP B4ECY9
B	-5	HIS	-	expression tag	UNP B4ECY9
B	-4	HIS	-	expression tag	UNP B4ECY9
B	-3	HIS	-	expression tag	UNP B4ECY9
B	-2	HIS	-	expression tag	UNP B4ECY9
B	-1	HIS	-	expression tag	UNP B4ECY9
B	0	HIS	-	expression tag	UNP B4ECY9
C	-7	MET	-	expression tag	UNP B4ECY9
C	-6	ALA	-	expression tag	UNP B4ECY9
C	-5	HIS	-	expression tag	UNP B4ECY9
C	-4	HIS	-	expression tag	UNP B4ECY9
C	-3	HIS	-	expression tag	UNP B4ECY9


Continued on next page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	HIS	-	expression tag	UNP B4ECY9
C	-1	HIS	-	expression tag	UNP B4ECY9
C	0	HIS	-	expression tag	UNP B4ECY9
D	-7	MET	-	expression tag	UNP B4ECY9
D	-6	ALA	-	expression tag	UNP B4ECY9
D	-5	HIS	-	expression tag	UNP B4ECY9
D	-4	HIS	-	expression tag	UNP B4ECY9
D	-3	HIS	-	expression tag	UNP B4ECY9
D	-2	HIS	-	expression tag	UNP B4ECY9
D	-1	HIS	-	expression tag	UNP B4ECY9
D	0	HIS	-	expression tag	UNP B4ECY9

- PLP
-
- The diagram shows the chemical structure of PLP (Pyridoxal phosphate). It consists of a pyridine ring with a nitrogen atom (N1) at the top. The ring is substituted with a hydroxyl group (OH) at C3, a carboxyl group (COOH) at C4, and a phosphate group (PO3H2) at C5. The phosphate group is shown as a phosphorus atom (P) bonded to four oxygen atoms: one double-bonded (O1P), one single-bonded and hydroxylated (O2P), and two single-bonded and hydroxylated (O3P, O4P). The carboxyl group is shown as a carbon atom (C4A) double-bonded to an oxygen atom (O4A) and single-bonded to a hydroxyl group (OH, O3). The pyridine ring carbons are labeled C2, C3, C4, C5, and C6. The phosphate group carbons are labeled C5A and C4A. The nitrogen atom is labeled N1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	B	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	C	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	D	1	Total 15	C 8	N 1	O 5	P 1	0	0

- 



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			7	3	1	3		
3	A	1	Total	C	N	O	0	0
			7	3	1	3		
3	B	1	Total	C	N	O	0	0
			7	3	1	3		
3	B	1	Total	C	N	O	0	0
			7	3	1	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	261	Total	O	0	0
			261	261		
4	B	268	Total	O	0	0
			268	268		
4	C	172	Total	O	0	0
			172	172		
4	D	192	Total	O	0	0
			192	192		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine hydroxymethyltransferase




- Molecule 1: Serine hydroxymethyltransferase



- Molecule 1: Serine hydroxymethyltransferase



- Molecule 1: Serine hydroxymethyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.22Å 179.84Å 75.64Å 90.00° 115.25° 90.00°	Depositor
Resolution (Å)	42.37 – 2.00 42.37 – 2.00	Depositor EDS
% Data completeness (in resolution range)	91.4 (42.37-2.00) 91.4 (42.37-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 2.00Å)	Xtriage
Refinement program	PHENIX dev_1659	Depositor
R, R_{free}	0.165 , 0.206 0.176 , 0.207	Depositor DCC
R_{free} test set	4895 reflections (4.55%)	wwPDB-VP
Wilson B-factor (Å ²)	27.9	Xtriage
Anisotropy	0.205	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 40.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.037 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13390	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.34	0/3189	0.74	0/4334
1	B	0.33	0/3183	0.75	0/4324
1	C	0.33	0/3200	0.74	0/4351
1	D	0.33	0/3146	0.74	0/4276
All	All	0.33	0/12718	0.74	0/17285

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	2	PHE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3111	0	3043	12	0
1	B	3102	0	3054	11	0
1	C	3124	0	3054	5	0
1	D	3072	0	3020	9	0
2	A	15	0	7	1	0
2	B	15	0	7	0	0
2	C	15	0	7	1	0
2	D	15	0	7	0	0
3	A	14	0	8	2	0
3	B	14	0	8	0	0
4	A	261	0	0	1	0
4	B	268	0	0	2	0
4	C	172	0	0	0	0
4	D	192	0	0	1	0
All	All	13390	0	12215	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:32:GLU:OE1	4:D:1086:HOH:O	2.03	0.76
2:A:501:PLP:HO3	3:A:502:SER:N	1.82	0.76
1:B:32:GLU:OE1	4:B:769:HOH:O	2.04	0.75
1:A:32:GLU:OE1	4:A:840:HOH:O	2.06	0.72
1:A:72:VAL:HB	1:A:264:MET:HE2	1.80	0.64
1:C:72:VAL:HB	1:C:264:MET:HE2	1.80	0.63
1:D:72:VAL:HB	1:D:264:MET:HE2	1.81	0.63
1:B:72:VAL:HB	1:B:264:MET:HE2	1.81	0.62
1:A:186:ARG:HH21	1:A:186:ARG:HG3	1.66	0.59
3:A:503:SER:N	2:C:700:PLP:HO3	2.02	0.56
1:D:122:LEU:HD21	1:D:132:SER:HB2	1.89	0.55
1:D:227:THR:HB	1:D:229:HIS:CE1	2.51	0.46
1:B:36:SER:HB2	1:B:230:LYS:HE3	1.99	0.45
1:C:36:SER:HB2	1:C:230:LYS:HE3	1.99	0.45
1:B:227:THR:HB	1:B:229:HIS:CE1	2.51	0.45
1:A:70:GLU:H	1:A:70:GLU:CD	2.25	0.44
1:B:277:ALA:HA	1:B:282:PHE:CG	2.53	0.44
1:B:118:MET:HA	1:B:144:VAL:O	2.18	0.44
1:C:227:THR:HB	1:C:229:HIS:CE1	2.53	0.44
1:C:277:ALA:HA	1:C:282:PHE:CG	2.53	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9:ILE:HG22	1:A:17:PHE:HB2	2.00	0.43
1:D:321:LEU:HD11	1:D:360:ILE:HG23	1.99	0.43
1:A:277:ALA:HA	1:A:282:PHE:CG	2.54	0.43
1:A:150:GLU:HA	1:B:398:GLU:HG2	2.01	0.43
1:D:48:GLN:HA	1:D:265:HIS:HB2	2.01	0.43
1:D:294[B]:ARG:HE	1:D:298:GLU:CD	2.26	0.43
1:B:288:LYS:HD3	4:B:764:HOH:O	2.19	0.43
1:A:36:SER:HB2	1:A:230:LYS:HE3	2.01	0.42
1:C:321:LEU:HD11	1:C:360:ILE:HG23	2.02	0.42
1:B:214:TYR:CG	1:B:215:PRO:HD2	2.55	0.41
1:B:9:ILE:HG22	1:B:17:PHE:HB2	2.01	0.41
1:D:258:ILE:HG13	1:D:259:GLN:HG3	2.02	0.41
1:A:227:THR:HB	1:A:229:HIS:CE1	2.56	0.41
1:A:321:LEU:HD11	1:A:360:ILE:HG23	2.02	0.41
1:B:122:LEU:HD21	1:B:132:SER:HB2	2.03	0.41
1:A:48:GLN:HA	1:A:265:HIS:HB2	2.03	0.40
1:A:364:SER:N	1:A:365:PRO:CD	2.85	0.40
1:D:36:SER:HB2	1:D:230:LYS:HE3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	417/423 (99%)	408 (98%)	9 (2%)	0	100	100
1	B	415/423 (98%)	408 (98%)	7 (2%)	0	100	100
1	C	417/423 (99%)	409 (98%)	8 (2%)	0	100	100
1	D	410/423 (97%)	400 (98%)	10 (2%)	0	100	100
All	All	1659/1692 (98%)	1625 (98%)	34 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	314/332 (95%)	310 (99%)	4 (1%)	61	68
1	B	314/332 (95%)	312 (99%)	2 (1%)	78	85
1	C	314/332 (95%)	312 (99%)	2 (1%)	78	85
1	D	311/332 (94%)	309 (99%)	2 (1%)	78	85
All	All	1253/1328 (94%)	1243 (99%)	10 (1%)	73	80

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	TYR
1	A	70	GLU
1	A	128	LEU
1	A	345	ASN
1	B	56	TYR
1	B	128	LEU
1	C	56	TYR
1	C	128	LEU
1	D	56	TYR
1	D	128	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	51	GLN
1	B	97	ASN
1	B	345	ASN
1	C	77	GLN
1	D	24	ASN
1	D	345	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	PLP	D	800	3	15,15,16	1.17	2 (13%)	21,22,23	1.32	3 (14%)
3	SER	B	503	2	4,6,6	1.16	1 (25%)	2,7,7	1.83	1 (50%)
3	SER	B	502	2	4,6,6	1.18	1 (25%)	2,7,7	1.83	1 (50%)
2	PLP	C	700	3	15,15,16	1.13	2 (13%)	21,22,23	1.14	1 (4%)
2	PLP	B	501	3	15,15,16	1.14	2 (13%)	21,22,23	1.30	4 (19%)
2	PLP	A	501	3	15,15,16	1.13	2 (13%)	21,22,23	1.16	2 (9%)
3	SER	A	502	2	4,6,6	1.14	1 (25%)	2,7,7	1.96	1 (50%)
3	SER	A	503	2	4,6,6	1.19	1 (25%)	2,7,7	1.88	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	D	800	3	-	0/6/6/8	0/1/1/1
3	SER	B	503	2	-	0/6/6/6	-
3	SER	B	502	2	-	0/6/6/6	-
2	PLP	C	700	3	-	2/6/6/8	0/1/1/1
2	PLP	B	501	3	-	0/6/6/8	0/1/1/1
2	PLP	A	501	3	-	2/6/6/8	0/1/1/1
3	SER	A	502	2	-	0/6/6/6	-
3	SER	A	503	2	-	2/6/6/6	-

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	800	PLP	C2-N1	2.77	1.38	1.33
2	B	501	PLP	C2-N1	2.76	1.38	1.33
2	C	700	PLP	C2-N1	2.64	1.38	1.33
2	A	501	PLP	C2-N1	2.60	1.38	1.33
3	B	502	SER	OXT-C	-2.25	1.23	1.30
3	A	503	SER	OXT-C	-2.25	1.23	1.30
3	B	503	SER	OXT-C	-2.21	1.23	1.30
2	D	800	PLP	C6-N1	2.21	1.38	1.34
3	A	502	SER	OXT-C	-2.19	1.23	1.30
2	A	501	PLP	C6-N1	2.17	1.38	1.34
2	C	700	PLP	C6-N1	2.15	1.38	1.34
2	B	501	PLP	C6-N1	2.07	1.38	1.34

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	800	PLP	O4P-C5A-C5	2.76	114.52	109.36
3	A	502	SER	OXT-C-O	-2.75	117.84	124.08
3	A	503	SER	OXT-C-O	-2.64	118.09	124.08
3	B	502	SER	OXT-C-O	-2.58	118.24	124.08
3	B	503	SER	OXT-C-O	-2.57	118.26	124.08
2	B	501	PLP	C5-C6-N1	-2.42	119.89	123.83
2	C	700	PLP	C5-C6-N1	-2.41	119.91	123.83
2	B	501	PLP	C6-C5-C4	2.38	120.04	118.10
2	D	800	PLP	C5-C6-N1	-2.37	119.98	123.83
2	A	501	PLP	C5-C6-N1	-2.34	120.03	123.83
2	B	501	PLP	O4P-C5A-C5	2.23	113.54	109.36
2	B	501	PLP	C5A-C5-C4	-2.21	118.27	122.64
2	D	800	PLP	C5A-C5-C4	-2.10	118.49	122.64
2	A	501	PLP	O4P-C5A-C5	2.02	113.14	109.36

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	PLP	C4-C5-C5A-O4P
2	C	700	PLP	C4-C5-C5A-O4P
2	C	700	PLP	C6-C5-C5A-O4P
3	A	503	SER	O-C-CA-N
2	A	501	PLP	C6-C5-C5A-O4P
3	A	503	SER	OXT-C-CA-N

There are no ring outliers.

4 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	700	PLP	1	0
2	A	501	PLP	1	0
3	A	502	SER	1	0
3	A	503	SER	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	414/423 (97%)	-0.41	6 (1%) 73 73	8, 21, 39, 84	5 (1%)
1	B	411/423 (97%)	-0.38	3 (0%) 84 84	10, 22, 39, 67	6 (1%)
1	C	415/423 (98%)	-0.09	8 (1%) 66 66	10, 32, 53, 75	4 (0%)
1	D	408/423 (96%)	-0.30	2 (0%) 87 87	10, 27, 45, 52	4 (0%)
All	All	1648/1692 (97%)	-0.30	19 (1%) 76 76	8, 25, 46, 84	19 (1%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	3	ASP	3.3
1	A	5	ALA	3.1
1	A	4	ARG	3.1
1	C	2	PHE	3.1
1	A	2	PHE	2.9
1	A	7	SER	2.8
1	B	5	ALA	2.8
1	C	395	ALA	2.7
1	C	0	HIS	2.7
1	C	363	GLY	2.6
1	C	5	ALA	2.6
1	C	355	PHE	2.4
1	C	1	MET	2.3
1	D	7	SER	2.2
1	C	192	LYS	2.2
1	D	401	ARG	2.1
1	B	7	SER	2.1
1	B	65	TYR	2.1
1	A	9	ILE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SER	B	502	7/7	0.83	0.12	36,37,40,42	0
3	SER	A	503	7/7	0.88	0.13	33,33,40,46	0
3	SER	B	503	7/7	0.90	0.09	34,35,39,41	0
3	SER	A	502	7/7	0.94	0.10	22,22,26,29	0
2	PLP	B	501	15/16	0.94	0.08	25,31,33,34	0
2	PLP	D	800	15/16	0.96	0.07	18,29,31,32	0
2	PLP	C	700	15/16	0.97	0.07	17,27,29,31	0
2	PLP	A	501	15/16	0.98	0.06	17,20,22,22	0

6.5 Other polymers [i](#)

There are no such residues in this entry.