



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:10 PM UTC

PDB ID : 4OTL / pdb_00004otl
Title : X-ray Crystal Structure of Serine Hydroxymethyl Transferase from Burkholderia cenocepacia bound to PLP and Glycine
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2014-02-13
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	423	2% 91% 5% .
1	B	423	2% 90% 7% .
1	C	423	2% 94% . .
1	D	423	2% 95% . .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 13382 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine hydroxymethyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	409	Total	C	N	O	S	0	9	0
			3123	1978	543	590	12			
1	B	409	Total	C	N	O	S	0	7	0
			3113	1967	542	592	12			
1	C	414	Total	C	N	O	S	0	7	0
			3142	1984	555	591	12			
1	D	416	Total	C	N	O	S	0	2	0
			3130	1972	552	593	13			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	expression tag	UNP B4ECY9
A	-6	ALA	-	expression tag	UNP B4ECY9
A	-5	HIS	-	expression tag	UNP B4ECY9
A	-4	HIS	-	expression tag	UNP B4ECY9
A	-3	HIS	-	expression tag	UNP B4ECY9
A	-2	HIS	-	expression tag	UNP B4ECY9
A	-1	HIS	-	expression tag	UNP B4ECY9
A	0	HIS	-	expression tag	UNP B4ECY9
B	-7	MET	-	expression tag	UNP B4ECY9
B	-6	ALA	-	expression tag	UNP B4ECY9
B	-5	HIS	-	expression tag	UNP B4ECY9
B	-4	HIS	-	expression tag	UNP B4ECY9
B	-3	HIS	-	expression tag	UNP B4ECY9
B	-2	HIS	-	expression tag	UNP B4ECY9
B	-1	HIS	-	expression tag	UNP B4ECY9
B	0	HIS	-	expression tag	UNP B4ECY9
C	-7	MET	-	expression tag	UNP B4ECY9
C	-6	ALA	-	expression tag	UNP B4ECY9
C	-5	HIS	-	expression tag	UNP B4ECY9
C	-4	HIS	-	expression tag	UNP B4ECY9
C	-3	HIS	-	expression tag	UNP B4ECY9

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	HIS	-	expression tag	UNP B4ECY9
C	-1	HIS	-	expression tag	UNP B4ECY9
C	0	HIS	-	expression tag	UNP B4ECY9
D	-7	MET	-	expression tag	UNP B4ECY9
D	-6	ALA	-	expression tag	UNP B4ECY9
D	-5	HIS	-	expression tag	UNP B4ECY9
D	-4	HIS	-	expression tag	UNP B4ECY9
D	-3	HIS	-	expression tag	UNP B4ECY9
D	-2	HIS	-	expression tag	UNP B4ECY9
D	-1	HIS	-	expression tag	UNP B4ECY9
D	0	HIS	-	expression tag	UNP B4ECY9

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	C	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	D	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is GLYCINE (CCD ID: GLY) (formula: $C_2H_5NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			5	2	1	2		
3	B	1	Total	C	N	O	0	0
			5	2	1	2		
3	C	1	Total	C	N	O	0	0
			5	2	1	2		
3	D	1	Total	C	N	O	0	0
			5	2	1	2		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	222	Total	O	0	0
			222	222		
5	B	245	Total	O	0	0
			245	245		
5	C	171	Total	O	0	0
			171	171		
5	D	152	Total	O	0	0
			152	152		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine hydroxymethyltransferase




- Molecule 1: Serine hydroxymethyltransferase



- Molecule 1: Serine hydroxymethyltransferase



- Molecule 1: Serine hydroxymethyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.97Å 179.69Å 75.51Å 90.00° 115.43° 90.00°	Depositor
Resolution (Å)	42.56 – 2.00 42.56 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.3 (42.56-2.00) 99.3 (42.56-2.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.76 (at 2.00Å)	Xtriage
Refinement program	PHENIX dev_1659	Depositor
R, R_{free}	0.180 , 0.224 0.186 , 0.226	Depositor DCC
R_{free} test set	5376 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	26.8	Xtriage
Anisotropy	0.182	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 40.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.036 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13382	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PLP, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.36	0/3214	0.75	4/4362 (0.1%)
1	B	0.37	0/3198	0.73	0/4344
1	C	0.34	0/3226	0.74	0/4379
1	D	0.35	0/3199	0.73	0/4345
All	All	0.35	0/12837	0.74	4/17430 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	390	ASN	CA-C-N	5.29	124.90	119.56
1	A	390	ASN	C-N-CA	5.29	124.90	119.56
1	A	95	GLN	CA-C-N	-5.22	114.34	119.93
1	A	95	GLN	C-N-CA	-5.22	114.34	119.93

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3123	0	3093	12	0
1	B	3113	0	3060	16	0
1	C	3142	0	3099	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	3130	0	3068	7	0
2	A	15	0	6	0	0
2	B	15	0	7	2	0
2	C	15	0	7	0	0
2	D	15	0	7	1	0
3	A	5	0	2	0	0
3	B	5	0	2	1	0
3	C	5	0	2	0	0
3	D	5	0	2	0	0
4	B	4	0	6	0	0
5	A	222	0	0	1	0
5	B	245	0	0	2	0
5	C	171	0	0	1	0
5	D	152	0	0	0	0
All	All	13382	0	12361	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:72:VAL:HB	1:B:264:MET:HE2	1.78	0.63
1:C:72:VAL:HB	1:C:264:MET:HE2	1.80	0.63
1:B:230:LYS:HZ1	2:B:501:PLP:C4A	2.15	0.59
1:C:398[A]:GLU:OE1	1:C:401:ARG:NH1	2.32	0.59
1:D:72:VAL:HB	1:D:264:MET:HE2	1.84	0.58
1:D:321:LEU:HD11	1:D:360:ILE:HG23	1.89	0.55
1:B:410:ARG:HD2	1:B:411:PHE:CE2	2.42	0.55
1:B:201:ASP:OD2	2:B:501:PLP:N1	2.43	0.52
1:A:306:ARG:HB3	1:A:320:ASP:HB3	1.92	0.52
1:D:306:ARG:HB3	1:D:320:ASP:HB3	1.92	0.51
1:B:277:ALA:HA	1:B:282:PHE:CG	2.47	0.50
1:A:389:GLU:OE2	5:A:796:HOH:O	2.20	0.49
1:C:229:HIS:ND1	1:C:236[A]:ARG:HA	2.28	0.49
1:B:306:ARG:HB3	1:B:320:ASP:HB3	1.94	0.49
1:C:229:HIS:ND1	1:C:236[B]:ARG:HA	2.29	0.48
1:C:36:SER:HB2	1:C:230:LYS:HZ3	1.78	0.48
1:A:277:ALA:HA	1:A:282:PHE:CG	2.49	0.47
1:D:277:ALA:HA	1:D:282:PHE:CG	2.49	0.47
1:B:9:ILE:HG22	1:B:17:PHE:HB2	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:31:ILE:HG13	1:D:404:VAL:HG13	1.97	0.46
1:D:230:LYS:HZ1	2:D:800:PLP:C4A	2.29	0.46
1:B:58:GLU:HB3	1:B:255:PHE:CZ	2.52	0.44
1:C:321:LEU:HD11	1:C:360:ILE:HG23	1.99	0.44
1:A:9:ILE:HG22	1:A:17:PHE:HB2	2.01	0.43
1:A:236[B]:ARG:HG3	1:C:53:THR:OG1	2.18	0.43
1:B:379[B]:GLN:NE2	5:B:825:HOH:O	2.46	0.43
1:C:58:GLU:HB3	1:C:255:PHE:CZ	2.54	0.42
1:B:229:HIS:ND1	1:B:236[A]:ARG:HA	2.35	0.42
1:B:230:LYS:HZ1	3:B:502:GLY:N	2.18	0.42
1:B:393:ASP:OD1	5:B:828:HOH:O	2.22	0.42
1:A:53:THR:OG1	1:C:236[B]:ARG:HG3	2.20	0.42
1:A:72:VAL:HB	1:A:264:MET:HE2	2.01	0.42
1:C:139:LYS:NZ	5:C:917:HOH:O	2.34	0.42
1:A:28:GLU:O	1:A:408:THR:HB	2.20	0.42
1:A:364:SER:N	1:A:365:PRO:CD	2.83	0.42
1:B:44:VAL:HG13	1:B:269:ALA:HB1	2.02	0.41
1:B:384:ILE:O	1:B:388:LEU:HG	2.20	0.41
1:B:364:SER:N	1:B:365:PRO:CD	2.84	0.41
1:A:321:LEU:HD11	1:A:360:ILE:HG23	2.01	0.41
1:A:56:TYR:HB3	1:A:66[B]:TYR:CE2	2.56	0.40
1:B:31:ILE:HG13	1:B:404:VAL:HG13	2.02	0.40
1:C:214:TYR:CG	1:C:215:PRO:HD2	2.56	0.40
1:A:93:ASN:OD1	1:A:96:PRO:HD3	2.21	0.40
1:D:214:TYR:CG	1:D:215:PRO:HD2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	416/423 (98%)	407 (98%)	9 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	414/423 (98%)	405 (98%)	9 (2%)	0	100	100
1	C	419/423 (99%)	411 (98%)	8 (2%)	0	100	100
1	D	416/423 (98%)	407 (98%)	9 (2%)	0	100	100
All	All	1665/1692 (98%)	1630 (98%)	35 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/332 (96%)	318 (100%)	1 (0%)	86	91
1	B	317/332 (96%)	315 (99%)	2 (1%)	78	85
1	C	317/332 (96%)	315 (99%)	2 (1%)	78	85
1	D	315/332 (95%)	314 (100%)	1 (0%)	86	91
All	All	1268/1328 (96%)	1262 (100%)	6 (0%)	81	87

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	TYR
1	B	56	TYR
1	B	345	ASN
1	C	56	TYR
1	C	345	ASN
1	D	56	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	GLN
1	A	345	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	24	ASN
1	B	77	GLN
1	B	142	ASN
1	D	142	ASN
1	D	345	ASN
1	D	403	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PLP	C	700	3	15,15,16	1.24	2 (13%)	21,22,23	1.69	6 (28%)
2	PLP	A	500	3	15,15,16	1.22	2 (13%)	21,22,23	1.74	6 (28%)
2	PLP	D	800	3	15,15,16	1.20	2 (13%)	21,22,23	1.54	6 (28%)
3	GLY	C	701	2	4,4,4	1.11	1 (25%)	3,4,4	1.39	0
3	GLY	D	801	2	4,4,4	1.14	1 (25%)	3,4,4	1.52	0
4	EDO	B	503	-	3,3,3	0.45	0	2,2,2	0.27	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PLP	B	501	3	15,15,16	1.17	1 (6%)	21,22,23	1.40	3 (14%)
3	GLY	B	502	2	4,4,4	1.16	1 (25%)	3,4,4	1.46	0
3	GLY	A	501	2	4,4,4	1.17	1 (25%)	3,4,4	1.53	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	C	700	3	-	5/6/6/8	0/1/1/1
2	PLP	A	500	3	-	3/6/6/8	0/1/1/1
2	PLP	D	800	3	-	1/6/6/8	0/1/1/1
3	GLY	C	701	2	-	1/2/2/2	-
3	GLY	D	801	2	-	2/2/2/2	-
4	EDO	B	503	-	-	0/1/1/1	-
2	PLP	B	501	3	-	0/6/6/8	0/1/1/1
3	GLY	B	502	2	-	0/2/2/2	-
3	GLY	A	501	2	-	0/2/2/2	-

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	700	PLP	C2-N1	2.83	1.38	1.33
2	A	500	PLP	C2-N1	2.75	1.38	1.33
2	D	800	PLP	C2-N1	2.70	1.38	1.33
2	B	501	PLP	C2-N1	2.60	1.38	1.33
2	D	800	PLP	C6-N1	2.34	1.39	1.34
3	A	501	GLY	OXT-C	-2.23	1.23	1.30
2	A	500	PLP	C6-N1	2.16	1.38	1.34
3	D	801	GLY	OXT-C	-2.15	1.23	1.30
3	B	502	GLY	OXT-C	-2.15	1.23	1.30
3	C	701	GLY	OXT-C	-2.08	1.23	1.30
2	C	700	PLP	C6-N1	2.04	1.38	1.34

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	PLP	C5A-C5-C4	-3.49	115.75	122.64
2	C	700	PLP	O4P-C5A-C5	3.47	115.86	109.36
2	B	501	PLP	O4P-C5A-C5	3.41	115.75	109.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	500	PLP	O4P-C5A-C5	3.32	115.58	109.36
2	D	800	PLP	O4P-C5A-C5	3.16	115.27	109.36
2	C	700	PLP	C5A-C5-C4	-3.13	116.46	122.64
2	C	700	PLP	C5-C6-N1	-2.84	119.21	123.83
2	D	800	PLP	C5A-C5-C4	-2.80	117.11	122.64
2	A	500	PLP	C6-C5-C4	2.78	120.38	118.10
2	A	500	PLP	C4A-C4-C5	-2.77	118.09	120.94
2	A	500	PLP	C5A-C5-C6	2.74	123.82	119.36
2	A	500	PLP	C5-C6-N1	-2.69	119.46	123.83
2	C	700	PLP	C6-C5-C4	2.62	120.24	118.10
2	D	800	PLP	C5-C6-N1	-2.61	119.58	123.83
2	B	501	PLP	C4A-C4-C5	-2.41	118.46	120.94
2	C	700	PLP	C5A-C5-C6	2.39	123.26	119.36
2	B	501	PLP	C5-C6-N1	-2.38	119.97	123.83
2	C	700	PLP	C4A-C4-C5	-2.28	118.59	120.94
2	D	800	PLP	C6-C5-C4	2.21	119.91	118.10
2	D	800	PLP	C5A-C5-C6	2.20	122.95	119.36
2	D	800	PLP	C4A-C4-C5	-2.11	118.77	120.94

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	500	PLP	C5A-O4P-P-O2P
3	D	801	GLY	OXT-C-CA-N
2	A	500	PLP	C5A-O4P-P-O1P
3	D	801	GLY	O-C-CA-N
2	A	500	PLP	C5A-O4P-P-O3P
2	C	700	PLP	C5A-O4P-P-O2P
2	C	700	PLP	C5A-O4P-P-O1P
2	C	700	PLP	C6-C5-C5A-O4P
2	C	700	PLP	C5A-O4P-P-O3P
2	D	800	PLP	C5A-O4P-P-O3P
2	C	700	PLP	C4-C5-C5A-O4P
3	C	701	GLY	OXT-C-CA-N

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	800	PLP	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	PLP	2	0
3	B	502	GLY	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	409/423 (96%)	-0.20	8 (1%) 65 64	10, 26, 40, 57	9 (2%)
1	B	409/423 (96%)	-0.20	8 (1%) 65 64	11, 24, 41, 60	7 (1%)
1	C	414/423 (97%)	0.02	7 (1%) 69 68	12, 30, 46, 70	7 (1%)
1	D	416/423 (98%)	0.22	10 (2%) 59 59	12, 35, 55, 74	2 (0%)
All	All	1648/1692 (97%)	-0.04	33 (2%) 65 64	10, 29, 47, 74	25 (1%)

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	5	ALA	4.5
1	B	7	SER	4.1
1	A	7	SER	4.1
1	B	9	ILE	3.6
1	A	66[A]	TYR	3.4
1	B	355	PHE	3.4
1	D	5	ALA	3.4
1	D	1	MET	3.2
1	C	2	PHE	3.2
1	D	0	HIS	3.1
1	A	9	ILE	2.9
1	C	415	ARG	2.7
1	D	7	SER	2.6
1	C	3	ASP	2.6
1	D	354	PRO	2.5
1	A	355	PHE	2.5
1	B	66[A]	TYR	2.4
1	D	4	ARG	2.4
1	C	6	GLN	2.4
1	A	65	TYR	2.4
1	A	415	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	390	ASN	2.3
1	C	65	TYR	2.3
1	D	32	GLU	2.2
1	B	8	THR	2.2
1	D	355	PHE	2.2
1	B	65	TYR	2.1
1	B	10	ALA	2.1
1	B	63	LYS	2.1
1	C	355	PHE	2.1
1	A	410	ARG	2.0
1	A	8	THR	2.0
1	D	391	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	B	503	4/4	0.70	0.19	51,53,54,56	0
3	GLY	D	801	5/5	0.90	0.10	29,32,33,35	0
3	GLY	A	501	5/5	0.90	0.13	29,33,36,38	0
2	PLP	A	500	15/16	0.92	0.11	23,30,36,38	0
3	GLY	C	701	5/5	0.92	0.10	24,26,29,30	0
2	PLP	B	501	15/16	0.93	0.10	20,25,35,38	0
2	PLP	D	800	15/16	0.94	0.10	17,30,39,39	0
3	GLY	B	502	5/5	0.96	0.07	19,21,27,28	0
2	PLP	C	700	15/16	0.96	0.08	25,29,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.