



wwPDB EM Validation Summary Report ⓘ

Mar 12, 2026 – 09:30 PM UTC

PDB ID : 8OT9 / pdb_00008ot9
EMDB ID : EMD-17173
Title : CTE typeIII tau filament from Guam ALS/PDC
Authors : Qi, C.; Yang, S.; Scheres, S.H.W.; Goedert, M.
Deposited on : 2023-04-20
Resolution : 3.40 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14717 (2.90 - 3.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	441	14% 83%
1	B	441	15% 83%
1	C	441	14% 83%
1	D	441	15% 83%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	441	 14% . 83%
1	F	441	 15% . 83%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3444 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Microtubule-associated protein tau.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	75	Total	C	N	O	S	0	0
			574	360	106	107	1		
1	B	75	Total	C	N	O	S	0	0
			574	360	106	107	1		
1	C	75	Total	C	N	O	S	0	0
			574	360	106	107	1		
1	D	75	Total	C	N	O	S	0	0
			574	360	106	107	1		
1	E	75	Total	C	N	O	S	0	0
			574	360	106	107	1		
1	F	75	Total	C	N	O	S	0	0
			574	360	106	107	1		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Microtubule-associated protein tau



• Molecule 1: Microtubule-associated protein tau



SER	ILE	PRO	SER	THR	HIS	SER	MET
ASP	GLY	GLY	ARG	PRO	VAL	GLU	ALA
MET	GLY	GLY	GLN	PRO	THR	THR	THR
VAL	S205	THR	SER	SER	GLN	ASP	ARG
ASP	V306	ALA	GLY	GLY	ARG	ALA	GLN
SER		PRO	GLU	MET	PRO	LYS	GLU
PRO	V309	VAL	PRO	PRO	VAL	LYS	PHE
GLN	V310	PRO	PRO	PRO	SER	THR	GLU
LEU	K311	MET	MET	LYS	LYS	THR	VAL
ALA		PRO	PRO	SER	SER	THR	MET
THR	S220	ASP	GLY	GLY	LYS	ALA	GLU
LEU	K321	LEU	ASP	ASP	ASP	GLU	ASP
ALA		LYS	ARG	GLY	GLY	ASP	ASP
ASP	K331	ASN	SER	SER	THR	VAL	ALA
GLU	P332	VAL	VAL	GLY	GLY	THR	THR
VAL		LYS	LYS	TYR	SER	ALA	ALA
SER	H362	SER	SER	SER	ASP	PRO	TYR
ALA	V363	LYS	LYS	SER	LEU	GLY	THR
SER	P364	ILE	ILE	PRO	LYS	VAL	LEU
LEU		GLY	GLY	GLY	LYS	ASP	GLY
ALA	L376	SER	SER	SER	ALA	GLU	ASP
ALA		THR	THR	PRO	LYS	GLY	ARG
LYS		GLU	GLY	GLY	GLY	ALA	LYS
GLN	K379	ASN	THR	THR	ALA	PRO	ASP
GLY		LEU	PRO	GLY	ASP	GLY	GLN
LEU	ALA	ALA	LYS	GLY	LYS	LYS	GLY
	ALA	LYS	HIS	SER	LYS	GLN	TYR
	ALA	GLN	GLN	ARG	THR	ALA	THR
	LYS	PRO	PRO	SER	LYS	ALA	THR
	THR	GLY	GLY	ARG	ILE	ALA	MET
	ASP	GLY	THR	THR	ALA	GLN	HIS
	GLY	GLY	PRO	PRO	THR	PRO	GLN
	HIS	LYS	LYS	SER	PRO	HIS	ASP
	GLU	ALA	VAL	LEU	ARG	GLN	GLY
	ILE	ILE	ILE	THR	GLY	ILE	GLY
	VAL	ILE	PRO	PRO	ALA	PRO	ASP
	TVR	ASN	PRO	PRO	PRO	GLU	THR
	LYS	LYS	THR	THR	PRO	GLY	ASP
	SER	LYS	ARG	GLY	GLY	THR	ALA
	PRO	LEU	GLU	GLN	GLN	THR	GLY
	VAL	ASP	PRO	LYS	LYS	ALA	LEU
	VAL	LEU	LYS	GLY	GLY	GLU	LYS
	SER	SER	VAL	VAL	GLN	GLU	SER
	GLY	ASN	VAL	ALA	ASN	GLY	PRO
	ASP	VAL	GLN	VAL	ALA	ILE	LEU
	THR	SER	GLN	VAL	ALA	ILE	GLN
	SER	SER	VAL	VAL	THR	GLY	LEU
	PRO	ARG	LYS	ARG	ILE	THR	THR
	GLY	CYS	THR	THR	ILE	THR	PRO
	HIS	GLY	PRO	PRO	ALA	PRO	THR
	LEU	SER	PRO	LYS	LYS	SER	GLY
	SER	LYS	LYS	THR	LYS	LEU	ASP
	ASN	ASP	SER	SER	THR	GLU	GLY
	VAL	ASN	ASN	PRO	THR	ASP	SER
	SER	ILE	ILE	SER	PRO	GLU	GLU
	SER	SER	SER	SER	ALA	ALA	ALA
	THR	GLY	HIS	LYS	ALA	PRO	PRO
	GLY	THR	VAL	LYS	THR	ALA	GLY

- [illegible]

4 Experimental information

Property	Value	Source
EM reconstruction method	HELICAL	Depositor
Imposed symmetry	HELICAL, twist=-1.19°, rise=4.74 Å, axial sym=C1	Depositor
Number of segments used	13456	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{Å}^2$)	40.0	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.041	Depositor
Minimum map value	-0.022	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.009	Depositor
Map size (Å)	329.6, 329.6, 329.6	wwPDB
Map dimensions	400, 400, 400	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.824, 0.824, 0.824	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.58	0/583	1.13	0/780
1	B	0.66	2/583 (0.3%)	1.14	1/780 (0.1%)
1	C	0.58	0/583	1.13	0/780
1	D	0.67	2/583 (0.3%)	1.14	1/780 (0.1%)
1	E	0.58	0/583	1.14	0/780
1	F	0.67	2/583 (0.3%)	1.14	1/780 (0.1%)
All	All	0.62	6/3498 (0.2%)	1.14	3/4680 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
1	E	0	1
1	F	0	1
All	All	0	6

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	362	HIS	CG-CD2	-5.62	1.29	1.35
1	B	362	HIS	CG-CD2	-5.62	1.29	1.35
1	D	362	HIS	CG-CD2	-5.60	1.29	1.35
1	D	330	HIS	CG-CD2	-5.32	1.30	1.35
1	F	330	HIS	CG-CD2	-5.26	1.30	1.35

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	362	HIS	CB-CG-CD2	-5.46	124.11	131.20
1	F	362	HIS	CB-CG-CD2	-5.46	124.11	131.20
1	D	362	HIS	CB-CG-CD2	-5.44	124.12	131.20

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	306	VAL	Peptide
1	B	326	GLY	Peptide
1	C	306	VAL	Peptide
1	D	326	GLY	Peptide
1	E	306	VAL	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	574	0	597	7	0
1	B	574	0	597	6	0
1	C	574	0	597	6	0
1	D	574	0	597	4	0
1	E	574	0	597	5	0
1	F	574	0	597	4	0
All	All	3444	0	3582	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 25 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:350:VAL:HG13	1:F:350:VAL:HG23	1.72	0.70
1:B:350:VAL:HG23	1:D:350:VAL:HG13	1.72	0.70
1:E:310:TYR:CD1	1:E:376:LEU:HD22	2.35	0.62
1:A:310:TYR:CD1	1:A:376:LEU:HD22	2.35	0.61
1:C:310:TYR:CD1	1:C:376:LEU:HD22	2.35	0.60

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	73/441 (17%)	71 (97%)	2 (3%)	0	100	100
1	B	73/441 (17%)	69 (94%)	4 (6%)	0	100	100
1	C	73/441 (17%)	71 (97%)	2 (3%)	0	100	100
1	D	73/441 (17%)	69 (94%)	4 (6%)	0	100	100
1	E	73/441 (17%)	71 (97%)	2 (3%)	0	100	100
1	F	73/441 (17%)	69 (94%)	4 (6%)	0	100	100
All	All	438/2646 (17%)	420 (96%)	18 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	66/358 (18%)	62 (94%)	4 (6%)	17	43
1	B	66/358 (18%)	61 (92%)	5 (8%)	12	38
1	C	66/358 (18%)	62 (94%)	4 (6%)	17	43
1	D	66/358 (18%)	61 (92%)	5 (8%)	12	38
1	E	66/358 (18%)	62 (94%)	4 (6%)	17	43
1	F	66/358 (18%)	61 (92%)	5 (8%)	12	38

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	396/2148 (18%)	369 (93%)	27 (7%)	16	40

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	321	LYS
1	D	356	SER
1	F	328	ILE
1	D	331	LYS
1	E	305	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	330	HIS
1	E	362	HIS
1	F	327	ASN
1	E	368	ASN
1	C	362	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-17173. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 200




Y Index: 200




Z Index: 200


6.2.2 Raw map



X Index: 200



Y Index: 200




Z Index: 200


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 193




Y Index: 215




Z Index: 207


6.3.2 Raw map



X Index: 193



Y Index: 215




Z Index: 207


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



X




Y




Z

The images above show the 3D surface view of the map at the recommended contour level 0.009. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5.2 Raw map



X



Y



Z

These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_17173_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 375 nm³; this corresponds to an approximate mass of 338 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.294 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of $0.294 \text{ \AA}^{-1}$

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.40	-	-
Author-provided FSC curve	3.35	3.98	3.47
Unmasked-calculated*	4.19	7.47	4.29

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.19 differs from the reported value 3.4 by more than 10 %

9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-17173 and PDB model 8OT9. Per-residue inclusion information can be found in section [3](#) on page [5](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.009 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.009).

9.4 Atom inclusion [i](#)



At the recommended contour level, 80% of all backbone atoms, 75% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.009) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.7510	0.5510
A	0.7500	0.5540
B	0.7450	0.5520
C	0.7630	0.5520
D	0.7450	0.5510
E	0.7540	0.5510
F	0.7490	0.5500

