



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 08:53 AM UTC

PDB ID : 3OUR / pdb_00003our
Title : Crystal structure of complex between EIIA and a novel pyruvate decarboxylase
Authors : Jeong, C.S.; An, Y.J.; Cha, S.S.
Deposited on : 2010-09-15
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	435	5% 59% 28% • 9%
1	C	435	2% 60% 28% • 9%
1	E	435	3% 59% 28% • 9%
1	G	435	59% 29% • 9%
2	B	183	50% 25% 7% 18%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	183	% 52%20%9%•18%
2	F	183	54%23%••18%
2	H	183	51%25%6%18%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 17650 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UPF0255 protein VV1_0328.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	394	Total	C	N	O	S	0	0	0
			3018	1931	509	563	15			
1	C	394	Total	C	N	O	S	0	0	0
			3079	1966	522	576	15			
1	E	396	Total	C	N	O	S	0	0	0
			3048	1947	517	569	15			
1	G	396	Total	C	N	O	S	0	0	0
			3098	1979	525	579	15			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP Q8DF91
A	-18	ARG	-	expression tag	UNP Q8DF91
A	-17	GLY	-	expression tag	UNP Q8DF91
A	-16	SER	-	expression tag	UNP Q8DF91
A	-15	HIS	-	expression tag	UNP Q8DF91
A	-14	HIS	-	expression tag	UNP Q8DF91
A	-13	HIS	-	expression tag	UNP Q8DF91
A	-12	HIS	-	expression tag	UNP Q8DF91
A	-11	HIS	-	expression tag	UNP Q8DF91
A	-10	HIS	-	expression tag	UNP Q8DF91
A	-9	GLY	-	expression tag	UNP Q8DF91
A	-8	SER	-	expression tag	UNP Q8DF91
A	-7	ALA	-	expression tag	UNP Q8DF91
A	-6	CYS	-	expression tag	UNP Q8DF91
A	-5	GLU	-	expression tag	UNP Q8DF91
A	-4	LEU	-	expression tag	UNP Q8DF91
A	-3	GLY	-	expression tag	UNP Q8DF91
A	-2	THR	-	expression tag	UNP Q8DF91
A	-1	PRO	-	expression tag	UNP Q8DF91
A	0	ASN	-	expression tag	UNP Q8DF91
C	-19	MET	-	expression tag	UNP Q8DF91

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-18	ARG	-	expression tag	UNP Q8DF91
C	-17	GLY	-	expression tag	UNP Q8DF91
C	-16	SER	-	expression tag	UNP Q8DF91
C	-15	HIS	-	expression tag	UNP Q8DF91
C	-14	HIS	-	expression tag	UNP Q8DF91
C	-13	HIS	-	expression tag	UNP Q8DF91
C	-12	HIS	-	expression tag	UNP Q8DF91
C	-11	HIS	-	expression tag	UNP Q8DF91
C	-10	HIS	-	expression tag	UNP Q8DF91
C	-9	GLY	-	expression tag	UNP Q8DF91
C	-8	SER	-	expression tag	UNP Q8DF91
C	-7	ALA	-	expression tag	UNP Q8DF91
C	-6	CYS	-	expression tag	UNP Q8DF91
C	-5	GLU	-	expression tag	UNP Q8DF91
C	-4	LEU	-	expression tag	UNP Q8DF91
C	-3	GLY	-	expression tag	UNP Q8DF91
C	-2	THR	-	expression tag	UNP Q8DF91
C	-1	PRO	-	expression tag	UNP Q8DF91
C	0	ASN	-	expression tag	UNP Q8DF91
E	-19	MET	-	expression tag	UNP Q8DF91
E	-18	ARG	-	expression tag	UNP Q8DF91
E	-17	GLY	-	expression tag	UNP Q8DF91
E	-16	SER	-	expression tag	UNP Q8DF91
E	-15	HIS	-	expression tag	UNP Q8DF91
E	-14	HIS	-	expression tag	UNP Q8DF91
E	-13	HIS	-	expression tag	UNP Q8DF91
E	-12	HIS	-	expression tag	UNP Q8DF91
E	-11	HIS	-	expression tag	UNP Q8DF91
E	-10	HIS	-	expression tag	UNP Q8DF91
E	-9	GLY	-	expression tag	UNP Q8DF91
E	-8	SER	-	expression tag	UNP Q8DF91
E	-7	ALA	-	expression tag	UNP Q8DF91
E	-6	CYS	-	expression tag	UNP Q8DF91
E	-5	GLU	-	expression tag	UNP Q8DF91
E	-4	LEU	-	expression tag	UNP Q8DF91
E	-3	GLY	-	expression tag	UNP Q8DF91
E	-2	THR	-	expression tag	UNP Q8DF91
E	-1	PRO	-	expression tag	UNP Q8DF91
E	0	ASN	-	expression tag	UNP Q8DF91
G	-19	MET	-	expression tag	UNP Q8DF91
G	-18	ARG	-	expression tag	UNP Q8DF91
G	-17	GLY	-	expression tag	UNP Q8DF91

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	-16	SER	-	expression tag	UNP Q8DF91
G	-15	HIS	-	expression tag	UNP Q8DF91
G	-14	HIS	-	expression tag	UNP Q8DF91
G	-13	HIS	-	expression tag	UNP Q8DF91
G	-12	HIS	-	expression tag	UNP Q8DF91
G	-11	HIS	-	expression tag	UNP Q8DF91
G	-10	HIS	-	expression tag	UNP Q8DF91
G	-9	GLY	-	expression tag	UNP Q8DF91
G	-8	SER	-	expression tag	UNP Q8DF91
G	-7	ALA	-	expression tag	UNP Q8DF91
G	-6	CYS	-	expression tag	UNP Q8DF91
G	-5	GLU	-	expression tag	UNP Q8DF91
G	-4	LEU	-	expression tag	UNP Q8DF91
G	-3	GLY	-	expression tag	UNP Q8DF91
G	-2	THR	-	expression tag	UNP Q8DF91
G	-1	PRO	-	expression tag	UNP Q8DF91
G	0	ASN	-	expression tag	UNP Q8DF91

- Molecule 2 is a protein called Phosphotransferase system IIA component.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	150	Total	C	N	O	S	0	0	0
			1121	714	177	228	2			
2	D	150	Total	C	N	O	S	0	0	0
			1117	713	178	224	2			
2	F	150	Total	C	N	O	S	0	0	0
			1121	715	178	226	2			
2	H	150	Total	C	N	O	S	0	0	0
			1121	714	177	228	2			

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-13	MET	-	expression tag	UNP Q8DFJ9
B	-12	ARG	-	expression tag	UNP Q8DFJ9
B	-11	GLY	-	expression tag	UNP Q8DFJ9
B	-10	SER	-	expression tag	UNP Q8DFJ9
B	-9	HIS	-	expression tag	UNP Q8DFJ9
B	-8	HIS	-	expression tag	UNP Q8DFJ9
B	-7	HIS	-	expression tag	UNP Q8DFJ9
B	-6	HIS	-	expression tag	UNP Q8DFJ9
B	-5	HIS	-	expression tag	UNP Q8DFJ9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-4	HIS	-	expression tag	UNP Q8DFJ9
B	-3	GLY	-	expression tag	UNP Q8DFJ9
B	-2	SER	-	expression tag	UNP Q8DFJ9
B	-1	ASP	-	expression tag	UNP Q8DFJ9
B	0	THR	-	expression tag	UNP Q8DFJ9
D	-13	MET	-	expression tag	UNP Q8DFJ9
D	-12	ARG	-	expression tag	UNP Q8DFJ9
D	-11	GLY	-	expression tag	UNP Q8DFJ9
D	-10	SER	-	expression tag	UNP Q8DFJ9
D	-9	HIS	-	expression tag	UNP Q8DFJ9
D	-8	HIS	-	expression tag	UNP Q8DFJ9
D	-7	HIS	-	expression tag	UNP Q8DFJ9
D	-6	HIS	-	expression tag	UNP Q8DFJ9
D	-5	HIS	-	expression tag	UNP Q8DFJ9
D	-4	HIS	-	expression tag	UNP Q8DFJ9
D	-3	GLY	-	expression tag	UNP Q8DFJ9
D	-2	SER	-	expression tag	UNP Q8DFJ9
D	-1	ASP	-	expression tag	UNP Q8DFJ9
D	0	THR	-	expression tag	UNP Q8DFJ9
F	-13	MET	-	expression tag	UNP Q8DFJ9
F	-12	ARG	-	expression tag	UNP Q8DFJ9
F	-11	GLY	-	expression tag	UNP Q8DFJ9
F	-10	SER	-	expression tag	UNP Q8DFJ9
F	-9	HIS	-	expression tag	UNP Q8DFJ9
F	-8	HIS	-	expression tag	UNP Q8DFJ9
F	-7	HIS	-	expression tag	UNP Q8DFJ9
F	-6	HIS	-	expression tag	UNP Q8DFJ9
F	-5	HIS	-	expression tag	UNP Q8DFJ9
F	-4	HIS	-	expression tag	UNP Q8DFJ9
F	-3	GLY	-	expression tag	UNP Q8DFJ9
F	-2	SER	-	expression tag	UNP Q8DFJ9
F	-1	ASP	-	expression tag	UNP Q8DFJ9
F	0	THR	-	expression tag	UNP Q8DFJ9
H	-13	MET	-	expression tag	UNP Q8DFJ9
H	-12	ARG	-	expression tag	UNP Q8DFJ9
H	-11	GLY	-	expression tag	UNP Q8DFJ9
H	-10	SER	-	expression tag	UNP Q8DFJ9
H	-9	HIS	-	expression tag	UNP Q8DFJ9
H	-8	HIS	-	expression tag	UNP Q8DFJ9
H	-7	HIS	-	expression tag	UNP Q8DFJ9
H	-6	HIS	-	expression tag	UNP Q8DFJ9
H	-5	HIS	-	expression tag	UNP Q8DFJ9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	-4	HIS	-	expression tag	UNP Q8DFJ9
H	-3	GLY	-	expression tag	UNP Q8DFJ9
H	-2	SER	-	expression tag	UNP Q8DFJ9
H	-1	ASP	-	expression tag	UNP Q8DFJ9
H	0	THR	-	expression tag	UNP Q8DFJ9


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	120	Total	O	0	0
			120	120		
3	B	74	Total	O	0	0
			74	74		
3	C	212	Total	O	0	0
			212	212		
3	D	75	Total	O	0	0
			75	75		
3	E	159	Total	O	0	0
			159	159		
3	F	52	Total	O	0	0
			52	52		
3	G	157	Total	O	0	0
			157	157		
3	H	78	Total	O	0	0
			78	78		


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: UPF0255 protein VV1_0328




• Molecule 1: UPF0255 protein VV1_0328






• Molecule 1: UPF0255 protein VV1_0328




• Molecule 1: UPF0255 protein VV1_0328




• Molecule 2: Phosphotransferase system IIA component







• Molecule 2: Phosphotransferase system IIA component



• Molecule 2: Phosphotransferase system IIA component



• Molecule 2: Phosphotransferase system IIA component



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	322.87Å 62.07Å 126.36Å 90.00° 110.80° 90.00°	Depositor
Resolution (Å)	49.37 – 2.20 49.37 – 2.20	Depositor EDS
% Data completeness (in resolution range)	90.5 (49.37-2.20) 98.6 (49.37-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 2.18Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.201 , 0.248 0.208 , 0.255	Depositor DCC
R_{free} test set	5953 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	29.2	Xtriage
Anisotropy	0.264	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 48.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.020 for -h-2*1,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	17650	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/3092	0.99	15/4211 (0.4%)
1	C	0.53	0/3152	1.00	11/4279 (0.3%)
1	E	0.51	1/3119 (0.0%)	0.99	18/4241 (0.4%)
1	G	0.55	0/3172	0.97	15/4306 (0.3%)
2	B	0.65	1/1135 (0.1%)	1.10	4/1539 (0.3%)
2	D	0.59	0/1131	1.12	9/1533 (0.6%)
2	F	0.61	0/1135	1.13	6/1538 (0.4%)
2	H	0.62	0/1135	1.14	9/1539 (0.6%)
All	All	0.55	2/17071 (0.0%)	1.02	87/23186 (0.4%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	60	MET	SD-CE	-5.98	1.64	1.79
1	E	89	THR	CA-CB	5.03	1.60	1.53

The worst 5 of 87 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	110	GLU	N-CA-C	8.73	121.67	111.02
1	C	89	THR	N-CA-C	8.34	123.26	112.26
2	F	112	GLN	N-CA-C	8.23	122.37	110.10
2	B	40	VAL	N-CA-C	7.67	118.45	110.62
2	H	112	GLN	N-CA-C	7.67	121.36	110.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3018	0	2882	109	0
1	C	3079	0	2994	92	0
1	E	3048	0	2941	103	0
1	G	3098	0	3024	98	0
2	B	1121	0	1137	43	0
2	D	1117	0	1140	54	0
2	F	1121	0	1144	34	0
2	H	1121	0	1137	49	0
3	A	120	0	0	4	0
3	B	74	0	0	1	0
3	C	212	0	0	3	0
3	D	75	0	0	5	0
3	E	159	0	0	2	0
3	F	52	0	0	3	0
3	G	157	0	0	3	0
3	H	78	0	0	3	0
All	All	17650	0	16399	558	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 558 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:127:LEU:HD11	2:H:131:LYS:HE2	1.30	1.11
1:E:410:GLU:OE1	1:E:414:LEU:HD12	1.58	1.04
2:B:37:VAL:HB	2:B:94:ILE:HD11	1.50	0.94
1:C:118:ASN:HD22	1:C:118:ASN:H	1.15	0.94
1:E:97:ASN:HD22	1:E:100:TYR:H	1.21	0.88

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/435 (90%)	341 (87%)	42 (11%)	9 (2%)	5	3
1	C	390/435 (90%)	356 (91%)	29 (7%)	5 (1%)	9	8
1	E	394/435 (91%)	366 (93%)	24 (6%)	4 (1%)	12	11
1	G	394/435 (91%)	368 (93%)	26 (7%)	0	100	100
2	B	148/183 (81%)	139 (94%)	9 (6%)	0	100	100
2	D	148/183 (81%)	138 (93%)	10 (7%)	0	100	100
2	F	148/183 (81%)	138 (93%)	10 (7%)	0	100	100
2	H	148/183 (81%)	142 (96%)	6 (4%)	0	100	100
All	All	2162/2472 (88%)	1988 (92%)	156 (7%)	18 (1%)	16	16

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	46	PHE
1	A	350	SER
1	C	326	LYS
1	C	329	VAL
1	E	241	THR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	310/377 (82%)	296 (96%)	14 (4%)	24	33
1	C	325/377 (86%)	310 (95%)	15 (5%)	24	32
1	E	316/377 (84%)	301 (95%)	15 (5%)	23	31
1	G	328/377 (87%)	305 (93%)	23 (7%)	14	16
2	B	125/153 (82%)	110 (88%)	15 (12%)	5	4
2	D	124/153 (81%)	100 (81%)	24 (19%)	1	1

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	F	125/153 (82%)	108 (86%)	17 (14%)	3	3
2	H	125/153 (82%)	111 (89%)	14 (11%)	6	5
All	All	1778/2120 (84%)	1641 (92%)	137 (8%)	12	13

5 of 137 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	320	LEU
1	G	400	GLN
2	H	96	THR
2	D	58	ASN
2	D	54	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 48 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	94	HIS
2	F	75	ASN
1	E	97	ASN
1	E	286	GLN
1	G	67	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	394/435 (90%)	0.43	23 (5%) 29 25	17, 42, 66, 73	0
1	C	394/435 (90%)	0.06	8 (2%) 65 62	13, 36, 63, 80	0
1	E	396/435 (91%)	0.08	11 (2%) 55 52	15, 38, 61, 73	0
1	G	396/435 (91%)	-0.06	2 (0%) 87 85	15, 34, 54, 61	0
2	B	150/183 (81%)	-0.31	0 100 100	16, 29, 46, 53	0
2	D	150/183 (81%)	-0.22	1 (0%) 84 82	19, 31, 48, 58	0
2	F	150/183 (81%)	-0.35	0 100 100	18, 27, 44, 50	0
2	H	150/183 (81%)	-0.42	0 100 100	16, 26, 45, 61	0
All	All	2180/2472 (88%)	0.00	45 (2%) 63 60	13, 34, 61, 80	0

The worst 5 of 45 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	59	TRP	3.5
1	E	414	LEU	3.4
1	E	401	SER	3.4
1	C	260	TYR	3.3
1	C	390	SER	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.